


# WORLD MALARIA REPORT 2020



YEARS OF GLOBAL PROGRESS & CHALLENGES



World Health  
Organization



# WORLD MALARIA REPORT 2020



YEARS OF GLOBAL PROGRESS & CHALLENGES



**World Health  
Organization**

World malaria report 2020: 20 years of global progress and challenges

ISBN 978-92-4-001579-1 (electronic version)

ISBN 978-92-4-001580-7 (print version)

© **World Health Organization 2020**

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: "This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition".

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<https://www.wipo.int/amc/en/mediation/rules/>).

**Suggested citation.** World malaria report 2020: 20 years of global progress and challenges. Geneva: World Health Organization; 2020. Licence: CC BY-NC-SA 3.0 IGO.

**Cataloguing-in-Publication (CIP) data.** CIP data are available at <https://apps.who.int/iris>.

**Sales, rights and licensing.** To purchase WHO publications, see <https://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <https://www.who.int/about/licensing>.

**Third-party materials.** If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

**General disclaimers.** The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Map production: WHO Global Malaria Programme and WHO Public Health Information and Geographic Systems.

Layout: Claude Cardot/[designisgood.info](http://designisgood.info)

Cover design: Lushomo (Cape Town, South Africa)

Please consult the WHO Global Malaria Programme website for the most up-to-date version of all documents (<https://www.who.int/teams/global-malaria-programme>).

# Contents

Foreword	vi
Acknowledgements	ix
Abbreviations and acronyms	xii
This year's report at a glance	xiv
<b>1. Introduction</b>	<b>1</b>
<b>2. Malaria milestones, 2000–2020</b>	<b>2</b>
2.1 Laying the foundations	2
2.2 2000–2004	4
2.3 2005–2010	6
2.4 2011–2015	8
2.5 2016–2019	12
<b>3. Global trends in the burden of malaria</b>	<b>18</b>
3.1 Global estimates of malaria cases and deaths, 2000–2019	18
3.2 Estimated malaria cases and deaths in the WHO African Region, 2000–2019	22
3.3 Estimated malaria cases and deaths in the WHO South-East Asia Region, 2000–2019	24
3.4 Estimated malaria cases and deaths in the WHO Eastern Mediterranean Region, 2000–2019	26
3.5 Estimated malaria cases and deaths in the WHO Western Pacific Region, 2000–2019	28
3.6 Estimated malaria cases and deaths in the WHO Region of the Americas, 2000–2019	30
3.7 Estimated malaria cases and deaths in the WHO European Region, 2000–2019	31
3.8 Cases and deaths averted since 2000, globally and by WHO region	32
3.9 Burden of malaria in pregnancy	34
<b>4. Elimination</b>	<b>38</b>
4.1 Malaria elimination certification	38
4.2 E-2020 initiative	38
4.3 The Greater Mekong subregion	40
4.4 Prevention of re-establishment	41
<b>5. High burden to high impact approach</b>	<b>42</b>
5.1 Galvanizing political will, mobilizing resources and mobilizing community response	42
5.2 Using strategic information to drive impact	45
5.3 Improving WHO's malaria policy-making and dissemination processes	48
5.4 Coordinated response	48
5.5 Malaria in HBHI countries since 2018	48
5.6 Reported malaria cases in HBHI countries since 2018 and comparisons with estimated cases	48
<b>6. Investments in malaria programmes and research</b>	<b>52</b>
6.1 Funding trends for malaria control and elimination	52
6.2 Investments in malaria-related R&D	56
<b>7. Distribution and coverage of malaria prevention, diagnosis and treatment</b>	<b>58</b>
7.1 Distribution and coverage of ITNs	58
7.2 Population protected with IRS	62
7.3 Scale-up of SMC	63
7.4 Coverage of IPTp use by dose	64
7.5 Malaria diagnosis and treatment	65
<b>8. Global progress towards the GTS milestones</b>	<b>70</b>
8.1 Global progress	70
8.2 WHO African Region	74
8.3 WHO Region of the Americas	76
8.4 WHO Eastern Mediterranean Region	78
8.5 WHO South-East Asia Region	79
8.6 WHO Western Pacific Region	80

<b>9. Biological threats</b>	82
9.1 Deletions in <i>P. falciparum</i> histidine-rich protein 2 and protein 3 genes	82
9.2 Therapeutic efficacy of ACTs	83
9.3 The global prevalence of <i>PfKelch13</i> molecular mutations	88
9.4 Vector resistance to insecticides	88
<b>10. Malaria response during the COVID-19 pandemic</b>	92
10.1 The 2020 COVID-19 pandemic	92
10.2 Global workstreams on sustaining the malaria response during the COVID-19 pandemic	95
10.3 Global highlights in the malaria response during the COVID-19 pandemic	96
10.4 Country responses to mitigate global service disruptions	98
10.5 Levels of service disruption by country and implications for delivery of interventions	100
10.6 The consequences of service disruptions during the COVID-19 pandemic	104
<b>11. Key results, context and conclusion</b>	106
11.1 Key results	106
11.2 The enabling environment and threats to the malaria progress	107
11.3 Consequences of the COVID-19 pandemic	112
11.4 Building a more prosperous future	113
11.5 Concluding remarks	115
<b>References</b>	116
<b>Annexes</b>	123



# Foreword



**Dr Tedros Adhanom Ghebreyesus**  
Director-General  
World Health Organization

In this year's *World malaria report*, WHO reflects on key milestones that have shaped the global response to the disease over the last 2 decades – a period of unprecedented success in malaria control that saw 1.5 billion cases averted and 7.6 million lives saved.

Following the end of the Global Malaria Eradication Programme in 1969, reduced political commitment and funding for malaria control led to resurgences of the disease in many parts of the world – particularly in Africa. While reliable data are scarce, hundreds of millions of people were likely infected with malaria, and tens of millions died.

Beginning in the 1990s, senior health leaders and scientists charted a course for a renewed response to malaria. Stepped-up investment in research and innovation led to the development of new disease-cutting tools, such as insecticide-treated nets, rapid diagnostic tests and more effective medicines.

The creation of new financing mechanisms – notably the Global Fund to Fight AIDS, Tuberculosis and Malaria and the US President's Malaria Initiative – coupled with a steep increase in malaria funding, enabled the wide-scale deployment of these tools, contributing to reductions in disease and death on a scale that had never been seen before.

Robust political commitment in Africa was key to success. Through the landmark 2000 Abuja Declaration, African leaders pledged to reduce malaria mortality on the continent by 50% over a 10-year timeframe.

According to our report, global malaria mortality fell by 60% over the period 2000 to 2019. The African Region achieved impressive reductions in its annual malaria death toll – from 680 000 in 2000 to 384 000 in 2019.

Countries in South-East Asia made particularly strong progress, with reductions in cases and deaths of 73% and 74%, respectively. India contributed to the largest drop in cases region-wide – from approximately 20 million to about 6 million.

Twenty-one countries have eliminated malaria over the last 2 decades and, of these, 10 countries were officially certified by WHO as malaria free. Countries of the Greater Mekong continue to make major gains, with a staggering 97% reduction in cases of *P. falciparum* malaria seen since 2000 – a primary target in view of the ongoing threat posed by antimalarial drug resistance.



## A plateau in progress

Progress made since the beginning of the millennium has been truly astonishing. However, as seen in this report, the gains have levelled off – a trend observed over recent years.

In 2017, WHO warned that the global response to malaria had reached a “crossroads”, and that key targets of WHO’s global malaria strategy would likely be missed. Three years on, we continue to see a plateau in progress; according to our latest report, the strategy’s 2020 targets for reductions in disease and death will be missed by 37% and 22%, respectively.

In 2020, COVID-19 emerged as an added – and formidable – challenge to malaria responses worldwide. In line with WHO guidance, many countries have adapted the way they deliver nets, diagnostics and medicines to ensure the safety of frontline health workers and communities. I wholeheartedly applaud these efforts, without which we would have likely seen much higher levels of mortality.

However, according to new WHO projections, even moderate disruptions in access to effective treatment could lead to a considerable loss of life. The report finds, for example, that a 25% disruption in access to effective antimalarial treatment in sub-Saharan Africa could lead to 46 000 additional deaths.

## Reigniting progress

To reinvigorate progress, WHO catalysed the “high burden to high impact” (HBHI) approach in 2018, together with the RBM Partnership to End Malaria. The response is led by 11 countries – including 10 in sub-Saharan Africa – that account for approximately 70% of the world’s malaria burden.

HBHI countries are moving away from a one-size-fits-all approach to malaria control – choosing instead to implement tailored responses based on local data and intelligence. While it is too early to evaluate the impact of this approach on malaria burden, important groundwork has been laid.

A recent analysis from Nigeria, for example, found that through an optimized mix of interventions the country could avert tens of millions of additional cases and thousands of additional deaths by the year 2023, compared with a business-as-usual approach.

A better targeting of malaria interventions and resources – particularly in countries like Nigeria, where the disease strikes hardest – will help speed the pace of progress towards our global malaria targets. Increased funding is also needed at domestic and international levels, together with innovations in new tools and approaches.

Crucially, efforts to combat malaria must be integrated with broader efforts to build strong health systems based on people-centred primary health care, as part of every country’s journey towards universal health coverage.

It is time for leaders across Africa – and the world – to rise once again to the challenge of malaria – just as they did when they laid the foundation for the progress made since the beginning of this century. Through joint action, and a commitment to leaving no one behind, we can achieve our shared vision of a world free of malaria.





# Acknowledgements

We are very grateful to the numerous people who contributed to the production of the *World malaria report 2020*. The following people collected and reviewed data from both malaria endemic and malaria free countries and areas:

Ahmad Mureed Muradi, Lutfullah Noori and Mohammad Shoaib Tamim (Afghanistan); Lammali Karima and Houria Khelifi (Algeria); Fernanda Francisco Guimarães and Fernanda Isabel Martins Da Graça Do Espirito Santo Alves (Angola); Malena Basilio and Yael Provecho (Argentina); Raja Alsaloom and Hasan Shuaib (Bahrain); Afsana Khan, Mya Ngon and Anjan K. Saha (Bangladesh); Kim Bautista (Belize); Telesphore Houansou and Aurore Ogouyomi-Hounto (Benin); Tobgyel Drukpa, Phurpa Tenzin and Sonam Wangdi (Bhutan); Raúl Marcelo Manjón Tellería (Bolivia [Plurinational State of]); Kentse Moakofhi, Mpho Mogopa, Davis Ntebela and Godira Segoea (Botswana); Cristianne Aparecida Costa Haraki, Franck Cardoso de Souza, Keyty Costa Cordeiro, Anderson Coutinho da Silva, Poliana de Brito Ribeiro Reis, Paloma Dias de Sousa, Francisco Edilson Ferreira de Lima Júnior, Klauss Kleydmann Sabino Garcia, Gilberto Gilmar Moresco, Marcela Lima Dourado, Paola Barbosa Marchesini, Márcia Helena Maximiano de Almeida, Joyce Mendes Pereira, Ronan Rocha Coelho, Edília Sâmela Freitas Santos, Pablo Sebastian Tavares Amaral and Marcelo Yoshito Wada (Brazil); Cheick S. Compaore and Laurent Moyenga (Burkina Faso); Dismas Baza and Juvénal Manirampa (Burundi); Carolina Cardoso da Silva Leite Gomes and Antonio Lima Moreira (Cabo Verde); Tol Bunkea (Cambodia); Abomabo Moise Hugue Rene and Etienne Nnomzo'o (Cameroon); Aristide Désiré Komangoya-Nzonzo and Christophe Ndoua (Central African Republic); Mahamat Idriss Djaskano and Honoré Djimrassengar (Chad); Wei Ding and Li Zhang (China); Eduin Pachón Abril (Colombia); Affane Bacar, Mohamed Issa Ibrahim and Ahamada Nassuri (Comoros); Hermann Ongouo and Jean-Mermoz Youndouka (Congo); Teresita Solano Chinchilla (Costa Rica); Tanoh Méa Antoine and N'goran Raphaël N'dri (Côte d'Ivoire); Kim Yun Chol, Nam Ju O and Gagan Sonal (Democratic People's Republic of Korea); Patrick Bahizi Bizozo, Hyacinthe Kaseya Ilunga, Bacary Sambou and Eric Mukomena Sompwe (Democratic Republic of the Congo); Basimike Mulenda (Djibouti); Keyla Urena (Dominican Republic); Monica Cañas Benavides and Julio Rivera (Ecuador); Angela Katherine Lao Seoane and Mathilde Riloha Rivas (Equatorial Guinea); Selam Mihreteab and Assefash Zehaie (Eritrea); Quinton Dhlamini, Kevin Makadzange and Zulisile Zulu (Eswatini); Henock Ejigu, Mebrahtom Haile and Bekele Worku (Ethiopia); Alice Sanna (French Guiana); Ghislaine Nkone Asseko and Okome Nze Gyslaine (Gabon); Momodou Kalleh and Sharmila Lareef-Jah (Gambia); Keziah Malm and Felicia Owusu-Antwi (Ghana); Ericka Lidia Chávez Vásquez (Guatemala); Siriman Camara and Nouman Diakite (Guinea); Inacio Alveranga and Paulo Djabatá (Guinea-Bissau); Helen Imhoff (Guyana); Antoine Darlie (Haiti); Engels Banegas, Jessica Henriquez, Carlos Miranda, Jose Orlander Nicolas, Raoul O'Connor and Nely Romero (Honduras); Neeraj Dhingra and Roop Kumari (India); Guntur Argana, Sri Budi Fajariyani, Herdiana Hasan Basri and M. Kez (Indonesia); Leila Faraji, Fatemeh Nikpoor and Ahmad Raeisi (Iran [Islamic Republic of]); James Kiarie and James Otieno (Kenya); Viengxay Vanisaveth (Lao People's Democratic Republic); Najib Achi (Lebanon); Moses Jeuronlon and Oliver J. Pratt (Liberia); Mauricette Andrianamanjara, Henintsoa Rabarijaona and Urbain Rabibizaka (Madagascar); Wilfred Dodoli, Austin Gumbo and Michael Kayange (Malawi); Jenarun Jelip (Malaysia); Sidibe Boubacar and Idrissa Cisse (Mali); Lemlih Baba and Sidina Mohamed Ghoulam (Mauritania); Frédéric Pages (Mayotte); Santa Elizabeth Ceballos Liceaga and Gustavo Sánchez Tejeda (Mexico); Balthazar Candrinho, Eva de Carvalho and Guidion Mathe (Mozambique); Md Rahman, Badri Thapa, Aung Thi and Tet Toe Tun (Myanmar); Rauha Jacob, Wilma Soroses and Petrina Uusiku (Namibia); Basu Dev Pandey, Subhash Lakhe and Prakash Prasad Shah (Nepal); Holvin Martin Gutierrez Perez (Nicaragua); Fatima Aboubakar and Hadiza Jackou (Niger); Audu Bala-Mohammed and Lynda Ozor (Nigeria); Hammad Habib (Pakistan); Lizbeth Cerezo and Santiago Cherigo (Panama); John Deli (Papua New Guinea); Cynthia Viveros (Paraguay); Cesar Bueno Cuadra (Peru); Raffy Deray, Kate Lopez and Maria Santa Portillo (Philippines); Jeong-Ran Kwon (Republic of Korea); Michee Kabera Semugunzu (Rwanda);

Claudina Augusto da Cruz and Jose Alvaro Leal Duarte (Sao Tome and Principe); Mohammed Hassan Al-Zahrani (Saudi Arabia); Ndella Diakhate and Medoune Ndiop (Senegal); Louisa Ganda and Samuel Juana Smith (Sierra Leone); John Leaburi (Solomon Islands); Abdi Abdillahi Ali, Ali Abdulrahmann, Abdikarim Hussein Hassan and Fahim Yusuf (Somalia); Mary Anne Groepe, Patrick Moonasar and Mbavhalelo Bridget Shandukani (South Africa); Harriet Akello Pasquale, Moses Jeuronlon and Moses Nganda (South Sudan); Navaratnasingam Janakan, Sumudu Karunaratna, Prasad Ranaweera and Preshila Samaraweera (Sri Lanka); Mariam Adam, Doha Elnazir and Abdalla Ibrahim (Sudan); Loretta Hardjopawiro (Suriname); Deyer Gopinath and Suravadee Kitchakarn (Thailand); Maria do Rosario de Fatima Mota, Rajesh Pandav and Manel Yapabandara (Timor-Leste); Kokou Mawule Davi and Tchassama Tchadjobo (Togo); Bayo Fatunmbi, Charles Katureebe, Paul Mbaka, John Opigo and Damian Rutazaana (Uganda); Abdullah Ali, Mohamed Haji Ali, Jovin Kitau, Anna Mahendeka, Ally Mohamed, Irene Mwoga and Ritha Njau (United Republic of Tanzania); Wesley Donald (Vanuatu); Licenciada América Rivero (Venezuela [Bolivarian Republic of]); Nguyen Quy Anh (Viet Nam); Moamer Mohammed Badi and Ryboon Saeed Al-Amoudi (Yemen); Japhet Chiwaula, Freddie Masaninga and Mutinta Mudenda (Zambia); and Anderson Chimusoro, Joseph Mberikunashe, Jasper Pasipamire and Ottias Tapfumanei (Zimbabwe).

We are grateful to the following people for their contribution:

Patrick Walker (Imperial College) contributed to the analysis of exposure to malaria infection during pregnancy and attributable low birthweight. Andre Marie Tchouatieu and Celine Audibert (Medicines for Malaria Venture [MMV]), and Paul Milligan (London School of Hygiene & Tropical Medicine) contributed to updating the section on seasonal malaria chemoprevention with the most up-to-date information on implementation and coverage. Manjiri Bhawalkar and Lisa Regis (Global Fund to Fight AIDS, Tuberculosis and Malaria [Global Fund]) supplied information on financial disbursements from the Global Fund. Adam Aspden and Nicola Wardrop (United Kingdom of Great Britain and Northern Ireland [United Kingdom] Department for International Development), and Adam Wexler and Julie Wallace (Kaiser Family Foundation) provided information on financial contributions for malaria control from the United Kingdom and the United States of America, respectively. Policy Cures Research used its G-FINDER data in the analysis of financing for malaria research and development, and wrote the associated section. John Milliner (Milliner Global Associates) provided information on long-lasting insecticidal nets delivered by manufacturers. The estimates of *Plasmodium falciparum* parasite prevalence and incidence in sub-Saharan Africa were produced by Daniel Weiss and Ewan Cameron of Malaria Atlas Project (MAP, led by Peter Gething, Curtin University and Telethon Kids Institute) and Samir Bhatt (Imperial College). Samir Bhatt, Amelia Bertozzi-Villa (Institute for Disease Modelling) and MAP collaborated to produce the estimates of insecticide-treated mosquito net (ITN) coverage for African countries using data from household surveys, ITN deliveries by manufacturers and ITNs distributed by national malaria programmes (NMPs). This research was funded by the Bill & Melinda Gates Foundation. Modelling of the impact of COVID-19 was contributed to by Peter Gething and Daniel Weiss with inputs from Samir Bhatt, Susan Rumisha (MAP) and Amelia Bertozzi-Villa. This research was funded by the Bill & Melinda Gates Foundation. Victor Alegana and Laurissa Suiyanka of Kenya Medical Research Institute (KEMRI) - Wellcome Trust Research Programme provided results of subnational analysis of concentration indices for socioeconomic equity in coverage of interventions, ITNs and treatment seeking. Tom McLean and Jason Richardson (Innovative Vector Control Consortium [IVCC]) provided national indoor residual spraying coverage and implementation data complementary to reported country information. Melanie Renshaw (African Leaders Malaria Alliance) and Marcy Erskine (Alliance for Malaria Prevention) provided information on the status of national ITN campaigns during the COVID-19 pandemic. Richard Steketee (United States President's Malaria Initiative [PMI]) reviewed the section on the malaria response during the COVID-19 pandemic. George Jagoe (MMV), Lisa Hare (PMI) and Andrea Bosman (World Health Organization [WHO] Global Malaria Programme [GMP]) contributed to the documentation of the global efforts to mitigate disruptions to diagnostics and antimalarials. Jennifer Armistead (PMI) provided insecticide resistance data on behalf of PMI. Gildas Yahouedo (WHO consultant) assisted with data compilation from publications. Colin Mathers (WHO consultant) and Bochen Cao (WHO Division of Data, Analytics and Delivery for

Impact [DDI]) prepared estimates of malaria mortality in children aged under 5 years, on behalf of the Child Health Epidemiology Reference Group. Yonas Tegegn (WHO Country Representative to Uganda), Bayo Segun Fatunmbi (WHO Uganda Country Office) and Jimmy Opigo (NMP, Uganda) contributed to the documentation on the Mass Action Against Malaria (MAAM) initiative in Uganda.

The following WHO staff in regional and subregional offices assisted in the design of data collection forms; the collection and validation of data; and the review of epidemiological estimates, country profiles, regional profiles and sections:

- Ebenezer Sheshi Baba, Emmanuel Chanda, Akpaka A. Kalu, Steve Kubenga Banza and Jackson Sillah (WHO Regional Office for Africa [AFRO]);
- Spes Ntabangana (AFRO/Inter-country Support Team [IST] Central Africa);
- Khoti Gausi (AFRO/IST East and Southern Africa);
- Abderrahmane Kharchi Tfeil (AFRO/IST West Africa);
- Maria Paz Ade, Janina Chavez, Rainier Escalada, Blanca Escribano, Roberto Montoya, Dennis Navaroo Costa, Eric Ndofor and Prabhjot Singh (WHO Regional Office for the Americas);
- Samira Al-Eryani and Ghasem Zamani (WHO Regional Office for the Eastern Mediterranean);
- Elena Chulkova and Elkhan Gasimov (WHO Regional Office for Europe);
- Risintha Premaratne and Neena Valecha (WHO Regional Office for South-East Asia); and
- James Kelley (WHO Regional Office for the Western Pacific).

The maps for country and regional profiles were produced by MAP's Data Engineering team funded by the Bill & Melinda Gates Foundation. The map production was led and coordinated by Jen Rozier, with help from Joe Harris and Suzanne Keddie. Tolu Okitika coordinated MAP's contribution to this report.

We are also grateful to Kevin Marsh (University of Oxford), Emelda Okiro (Kenya Medical Research Institute – Wellcome Trust Research Programme) and Larry Slutsker (PATH) who graciously reviewed all sections and provided substantial comments for improvement; Nelly Biondi, Diana Estevez Fernandez and Jessica Chi Ying Ho (WHO) for statistics review; Tessa Edejer and Agnès Soucat (WHO) for review of economic evaluation and analysis; Egle Granziera and Claudia Nannini (WHO) for legal review; Martha Quiñones (WHO consultant) and Beatriz Galatas (WHO) for the translation of the foreword and key points into Spanish, and Amélie Latour (WHO consultant) and Laurent Bergeron (WHO) for the translation into French; and Hilary Cadman and the Cadman Editing Services team for technical editing of the report.

On behalf of the WHO Global Malaria Programme (GMP), the publication of the *World malaria report 2020* was coordinated by Abdisalan Noor. Significant contributions were made by Pedro Alonso, Laura Anderson, John Aponte, Maru Aregawi, Amy Barrette, Yuen Ching Chan, Tamara Ehler, Lucia Fernandez Montoya, Beatriz Galatas, Mwalenga Nghipumbwa, Peter Olumese, Edith Patouillard, Alastair Robb, David Schellenberg and Ryan Williams. Laurent Bergeron (WHO GMP) provided programmatic support for overall management of the project. The editorial committee for the report comprised Pedro Alonso, Andrea Bosman, Jan Kolaczinski, Kimberly Lindblade, Leonard Ortega, Pascal Ringwald, Alastair Robb and David Schellenberg from the WHO GMP. Additional reviews were received from colleagues in the GMP: Jane Cunningham, Xiao Hong Li, Charlotte Rasmussen, Silvia Schwarte, Erin Shutes and Saira Stewart. Report layout, design and production were coordinated by Laurent Bergeron.

Funding for the production of this report was gratefully received from the Bill & Melinda Gates Foundation; the Global Fund; the Government of China; the Spanish Agency for International Development Cooperation; Unitaid; the United Nations Office for Project Services (UNOPS); and the United States Agency for International Development (USAID).

# Abbreviations and acronyms

ACT	artemisinin-based combination therapy	GMP	Global Malaria Programme
AIDS	acquired immunodeficiency syndrome	GMS	Greater Mekong subregion
AIM	<i>Action and investment to defeat malaria 2016–2030</i>	GPARC	Global Plan for Artemisinin Resistance Containment
AL	artemether-lumefantrine	GTS	<i>Global technical strategy for malaria 2016–2030</i>
ALMA	African Leaders Malaria Alliance	HBHI	high burden to high impact
AMFm	Affordable Medicines Facility-malaria	HCQ	hydroxychloroquine
An.	<i>Anopheles</i>	HIV	human immunodeficiency virus
ANC	antenatal care	HRP	histidine-rich protein
AQ	amodiaquine	IPTi	intermittent preventive treatment in infants
AS	artesunate	IPTp	intermittent preventive treatment in pregnancy
BAU	business as usual	IQR	interquartile range
CDC	Centers for Disease Control and Prevention	IRS	indoor residual spraying
CI	confidence interval	IST	Inter-country Support Team
CQ	chloroquine	ITN	insecticide-treated mosquito net
CRS	creditor reporting system	IVCC	Innovative Vector Control Consortium
DAC	Development Assistance Committee	LBW	low birthweight
DHA-PPQ	dihydroartemisinin-piperaquine	LGA	local government authority
DHIS2	District Health Information Software 2	LLIN	long-lasting insecticidal net
DHS	demographic and health survey	LMIC	low- and middle-income countries
E-2020	eliminating countries for 2020	LSHTM	London School of Hygiene & Tropical Medicine
EDCTP	European and Developing Countries Clinical Trials Partnership	MAAM	Mass Action Against Malaria
FIND	Foundation for Innovative New Diagnostics	MAP	Malaria Atlas Project
GDP	gross domestic product	MCEE	Maternal and Child Health Epidemiology Estimation Group
Global Fund	Global Fund to Fight AIDS, Tuberculosis and Malaria	MDG	Millennium Development Goal
GMAP	Global Malaria Action Plan for a malaria free world	MEDB	Malaria Elimination Database
		MIS	malaria indicator survey
		MME	Mekong Malaria Elimination
		MMV	Medicines for Malaria Venture

MPAC	Malaria Policy Advisory Committee	TES	therapeutic efficacy studies
MQ	mefloquine	UHC	universal health coverage
NMEP	National Malaria Elimination Programme	UN	United Nations
NMP	national malaria programme	UNDP	United Nations Development Programme
NMSP	national malaria strategic plan	UNICEF	United Nations Children's Fund
OECD	Organisation for Economic Co-operation and Development	United Kingdom	United Kingdom of Great Britain and Northern Ireland
<i>P.</i>	<i>Plasmodium</i>	US	United States
PBO	piperonyl butoxide	USA	United States of America
<i>pfhrp</i>	<i>Plasmodium falciparum</i> histidine-rich protein	USAID	United States Agency for International Development
pLDH	<i>Plasmodium</i> lactate dehydrogenase	WHO	World Health Organization
PMI	President's Malaria Initiative	WHO-CHOICE	WHO-CHOosing Interventions that are Cost-Effective
PPE	personal protective equipment		
PQ	primaquine		
PY	pyronaridine		
R&D	research and development		
RAI	Regional Artemisinin-resistance Initiative		
RDT	rapid diagnostic test		
SAGme	Strategic Advisory Group for Malaria Eradication		
SARS-CoV2	severe acute respiratory syndrome coronavirus 2		
SDG	Sustainable Development Goal		
SMC	seasonal malaria chemoprevention		
SP	sulfadoxine-pyrimethamine		
TDR	Special Programme for Research and Training in Tropical Diseases		

# This year's report at a glance

## TRENDS IN THE BURDEN OF MALARIA

---

### Malaria cases

- Globally, there were an estimated 229 million malaria cases in 2019 in 87 malaria endemic countries, declining from 238 million in 2000. At the *Global technical strategy for malaria 2016–2030* (GTS) baseline of 2015, there were 218 million estimated malaria cases.
- The proportion of cases due to *Plasmodium vivax* reduced from about 7% in 2000 to 3% in 2019.
- Malaria case incidence (i.e. cases per 1000 population at risk) reduced from 80 in 2000 to 58 in 2015 and 57 in 2019 globally. Between 2000 and 2015, global malaria case incidence declined by 27%, and between 2015 and 2019 it declined by less than 2%, indicating a slowing of the rate of decline since 2015.
- Twenty-nine countries accounted for 95% of malaria cases globally. Nigeria (27%), the Democratic Republic of the Congo (12%), Uganda (5%), Mozambique (4%) and Niger (3%) accounted for about 51% of all cases globally.
- The World Health Organization (WHO) African Region, with an estimated 215 million cases in 2019, accounted for about 94% of cases.
- Although there were fewer malaria cases in 2000 (204 million) than in 2019 in the WHO African Region, malaria case incidence reduced from 363 to 225 cases per 1000 population at risk in this period, reflecting the complexity of interpreting changing disease transmission in a rapidly increasing population. The population living in the WHO African Region increased from about 665 million in 2000 to 1.1 billion in 2019.
- The WHO South-East Asia Region accounted for about 3% of the burden of malaria cases globally. Malaria cases reduced by 73%, from 23 million in 2000 to about 6.3 million in 2019. Malaria case incidence in this region reduced by 78%, from about 18 cases per 1000 population at risk in 2000 to about four cases in 2019.
- India contributed to the largest absolute reductions in the WHO South-East Asia Region, from about 20 million cases in 2000 to about 5.6 million in 2019. Sri Lanka was certified malaria free in 2015, and Timor-Leste reported zero malaria cases in 2018 and 2019.
- Malaria cases in the WHO Eastern Mediterranean Region reduced by 26%, from about 7 million cases in 2000 to about 5 million in 2019. About a quarter of the cases in 2019 were due to *P. vivax*, mainly in Afghanistan and Pakistan.
- Over the period 2000–2019, malaria case incidence in the WHO Eastern Mediterranean Region declined from 20 to 10. Sudan is the leading contributor to malaria in this region, accounting for about 46% of cases. The Islamic Republic of Iran had no indigenous malaria cases in 2018 and 2019.
- The WHO Western Pacific Region had an estimated 1.7 million cases in 2019, a decrease of 43% from the 3 million cases in 2000. Over the same period, malaria case incidence reduced from five to two cases per 1000 population at risk. Papua New Guinea accounted for nearly 80% of all cases in this region in 2019. China has had no indigenous malaria cases since 2017. Malaysia had no cases of human malaria in 2018 and 2019.
- In the WHO Region of the Americas, malaria cases reduced by 40% (from 1.5 million to 0.9 million) and case incidence by 57% (from 14 to 6). The region's progress in recent years has suffered from the major increase in malaria in Venezuela (Bolivarian Republic of), which had about 35 500 cases



in 2000, rising to over 467 000 by 2019. Brazil, Colombia and Venezuela (Bolivarian Republic of) account for over 86% of all cases in this region.

- Since 2015, the WHO European Region has been free of malaria.

## **Malaria deaths**

- Globally, malaria deaths have reduced steadily over the period 2000–2019, from 736 000 in 2000 to 409 000 in 2019. The percentage of total malaria deaths among children aged under 5 years was 84% in 2000 and 67% in 2019. The global estimate of deaths in 2015, the GTS baseline, was about 453 000.
- Globally, the malaria mortality rate (i.e. deaths per 100 000 population at risk) reduced from about 25 in 2000 to 12 in 2015 and 10 in 2019, with the slowing of the rate of decline in the latter years.
- About 95% of malaria deaths globally were in 31 countries. Nigeria (23%), the Democratic Republic of the Congo (11%), the United Republic of Tanzania (5%), Mozambique (4%), Niger (4%) and Burkina Faso (4%) accounted for about 51% of all malaria deaths globally in 2019.
- Malaria deaths in the WHO African Region reduced by 44%, from 680 000 in 2000 to 384 000 in 2019, and the malaria mortality rate reduced by 67% over the same period, from 121 to 40 deaths per 100 000 population at risk.
- In the WHO South-East Asia Region, malaria deaths reduced by 74%, from about 35 000 in 2000 to 9 000 in 2019.
- India accounted for about 86% of all malaria deaths in the WHO South-East Asia Region.
- In the WHO Eastern Mediterranean Region, malaria deaths reduced by 16%, from about 12 000 in 2000 to 10 100 in 2019, and the malaria mortality rate reduced by 50%, from four to two deaths per 100 000 population at risk.
- In the WHO Western Pacific Region, malaria deaths reduced by 52%, from about 6600 cases in 2000 to 3200 in 2019, and the mortality rate reduced by 60%, from one to 0.4 malaria deaths per 100 000 population at risk. Papua New Guinea accounted for over 85% of malaria deaths in 2019.
- In the WHO Region of the Americas, malaria deaths reduced by 39% (from 909 to 551) and mortality rate by 50% (from 0.8 to 0.4). Over 70% of malaria deaths in 2019 in this region were in Venezuela (Bolivarian Republic of).

## **Malaria cases and deaths averted**

- Globally, an estimated 1.5 billion malaria cases and 7.6 million malaria deaths have been averted in the period 2000–2019.
- Most of the cases (82%) and deaths (94%) averted were in the WHO African Region, followed by the WHO South-East Asia Region (cases 10% and deaths 3%).

## **Burden of malaria in pregnancy**

- In 2019, in 33 moderate to high transmission countries in the WHO African Region, there were an estimated 33 million pregnancies, of which 35% (12 million) were exposed to malaria infection during pregnancy.
- By WHO subregion, Central Africa had the highest prevalence of exposure to malaria during pregnancy (40%), closely followed by West Africa (39%), while prevalence was 24% in East and Southern Africa.
- It is estimated that malaria infection during pregnancy in these 33 countries resulted in 822 000 children with low birthweight.
- If up to 80% of pregnant women who reported using antenatal care (ANC) services once were to receive one dose of intermittent preventive treatment in pregnancy (IPTp), an additional 56 000 low birthweights would be averted in these 33 countries.

## MALARIA ELIMINATION AND PREVENTION OF RE-ESTABLISHMENT

---

- Globally, the number of countries that were malaria endemic in 2000 and that reported fewer than 10 000 malaria cases increased from 26 in 2000 to 46 in 2019.
- In the same period, the number of countries with fewer than 100 indigenous cases increased from six to 27.
- In the period 2010–2019, total malaria cases in the 21 E-2020 countries reduced by 79%.
- There were more cases in 2019 than in 2018 in Comoros, Costa Rica, Ecuador and Suriname, which reported 1986, 25, 150 and 66 additional cases, respectively.
- Iran (Islamic Republic of), Malaysia and Timor-Leste reported zero indigenous malaria cases in 2018 and 2019. In 2019, Belize and Cabo Verde reported zero indigenous malaria cases for the first time since 2000.
- China and El Salvador had no indigenous malaria cases for a third consecutive year and have made a formal request for certification.
- Between 2000 and 2019, in the six countries of the Greater Mekong subregion (GMS) – Cambodia, China (Yunnan Province), Lao People’s Democratic Republic, Myanmar, Thailand and Viet Nam – *P. falciparum* malaria cases fell by 97%, while all malaria cases fell by 90%. Of the 239 000 malaria cases reported in 2019, 65 000 were *P. falciparum* cases.
- The rate of decline has been fastest since 2012, when the Mekong Malaria Elimination (MME) programme was launched. During this period, malaria cases reduced sixfold, while *P. falciparum* cases reduced by a factor of nearly 14.
- Overall, Cambodia (58%) and Myanmar (31%) accounted for most cases of malaria in the GMS.
- This accelerated decrease in *P. falciparum* is especially critical because of increasing drug resistance; in the GMS, *P. falciparum* parasites have developed partial resistance to artemisinin, the core compound of the best available antimalarial drugs.
- Between 2000 and 2019, no country that was certified malaria free has been found to have malaria transmission re-established.

## HIGH BURDEN TO HIGH IMPACT APPROACH

---

- Since November 2018, the high burden to high impact (HBHI) approach has been launched in 10 of the 11 countries (it has not yet been launched in Mali owing to disruptions due to the COVID-19 pandemic). However, all 11 countries have implemented HBHI-related activities across the four response elements.
- In each HBHI country initiation, there has been high-level political engagement and support. The Mass Action Against Malaria initiative in Uganda is presented as an example of a country-led process of political engagement at all levels, and multisectoral and community mobilization.
- Analysis for subnational tailoring of interventions has been completed in all countries except Mali, where this work is in progress. The example of Nigeria is presented in the report.
- All countries have committed to conduct a comprehensive exercise of urban microstratification to better target interventions and improve efficiencies given the increasing rate of urbanization.
- The WHO Global Malaria Programme (GMP) updated its technical brief to support countries to better prioritize resources, while adhering to the evidence-based recommendations that have been developed through WHO’s standard, stringent processes.
- Because the HBHI response was launched in November 2018, when countries were coming to the end of their funding cycles, it is too soon to determine the impact of the response. The numbers of malaria cases in the 11 HBHI countries in 2019 were similar to 2018 (156 million versus 155 million).

## PROGRESS TOWARDS THE GTS MILESTONES OF 2020

---

- The GTS aims for a reduction in malaria case incidence and mortality rate of at least 40% by 2020, 75% by 2025 and 90% by 2030 from a 2015 baseline.
- The 2000–2019 trends in malaria cases and deaths were used to make annual projections from 2020 to 2030, to track progress towards the targets and milestones of the GTS.
- The projections presented in this report *do not* account for potential disruptions due to the COVID-19 pandemic, which – despite commendable global and national efforts to maintain essential malaria services – is likely to lead to higher than expected malaria morbidity and mortality.
- Despite the considerable progress made since 2000, the GTS 2020 milestones for morbidity and mortality will not be achieved globally.
- Malaria case incidence of 56 cases per 1000 population at risk in 2020 instead of the expected 35 cases per 1000 if the world was on track for the 2020 GTS morbidity milestone means that, globally, we are off track by 37% at the current trajectory.
- Although relative progress in the mortality rate is greater than that in case incidence, globally projected malaria deaths per 100 000 population at risk in 2020 was 9.8, reducing from 11.9 in 2015, implying that the world was off track for the 2020 GTS mortality milestone by 22%.
- Of the 92 countries that were malaria endemic globally in 2015, 31 (34%) were estimated to be on track for the GTS morbidity milestone for 2020, having achieved 40% or more reduction in case incidence or reported zero malaria cases.
- Twenty-one countries (23%) had made progress in reducing malaria case incidence but were not on track for the GTS milestone.
- Thirty-one countries (34%) are estimated to have increased incidence, with 15 countries (16%) estimated to have an increase of 40% or more in malaria case incidence in 2020 compared with 2015.
- Malaria case incidence in nine countries (10%) in 2020 was estimated to be at levels similar to those of 2015.
- Thirty-nine countries (42%) that were malaria endemic in 2015 were on track for the GTS mortality milestone for 2020, with 28 of them reporting zero malaria cases.
- Thirty-four countries (37%) were estimated to have achieved reductions in malaria mortality rates but progress was below the 40% target.
- Malaria mortality rates remained at the same level in 2020 as 2015 in seven countries (8%), whereas there were estimated increases in another 12 countries (13%), six of which had increases of 40% or more.
- All countries in the WHO South-East Asia Region were on track for both the morbidity and mortality 2020 GTS milestones.

## INVESTMENTS IN MALARIA PROGRAMMES AND RESEARCH

---

- The GTS sets out estimates of the funding required to achieve milestones for 2020, 2025 and 2030. Total annual resources needed were estimated at US\$ 4.1 billion in 2016, rising to US\$ 6.8 billion in 2020. An additional US\$ 0.72 billion is estimated to be required annually for global malaria research and development (R&D).
- Total funding for malaria control and elimination in 2019 was estimated at US\$ 3.0 billion, compared with US\$ 2.7 billion in 2018 and US\$ 3.2 billion in 2017. The amount invested in 2019 falls short of the US\$ 5.6 billion estimated to be required globally to stay on track towards the GTS milestones.

- The funding gap between the amount invested and the resources needed has continued to widen dramatically over recent years, increasing from US\$ 1.3 billion in 2017 to US\$ 2.3 billion in 2018, and to US\$ 2.6 billion in 2019.
- Over the period 2010–2019, international sources provided 70% of the total funding for malaria control and elimination, led by the United States of America (USA), the United Kingdom of Great Britain and Northern Ireland (United Kingdom) and France.
- Of the US\$ 3.0 billion invested in 2019, US\$ 2.1 billion came from international funders. The highest contributions in 2019 were from the government of the USA, which provided a total of US\$ 1.1 billion through planned bilateral funding and contributions to multilateral funding agencies.
- This was followed by bilateral and multilateral disbursements from the United Kingdom of US\$ 0.2 billion, contributions of over US\$ 0.1 billion from each of France, Germany and Japan (totalling US\$ 0.4 billion), and a combined US\$ 0.4 billion from other countries that are members of the Development Assistance Committee and from private sector contributors.
- Governments of malaria endemic countries continued to contribute about 30% of the total funding, with investments nearing US\$ 0.9 billion in 2019. Of this amount, an estimated US\$ 0.2 billion was spent on malaria case management in the public sector and US\$ 0.7 billion on other malaria control activities.
- Of the US\$ 3.0 billion invested in 2019, nearly US\$ 1.2 billion (39%) was channelled through the Global Fund to Fight AIDS, Tuberculosis and Malaria (Global Fund). Compared with 2018, the Global Fund's disbursements to malaria endemic countries increased by about US\$ 0.2 billion in 2019.
- Of the US\$ 3.0 billion invested in 2019, about 73% went to the WHO African Region, 9% to the WHO South-East Asia Region, 5% each to the WHO Region of the Americas and the WHO Western Pacific Region, and 4% to the WHO Eastern Mediterranean Region.
- Between 2007 and 2018, almost US\$ 7.3 billion was invested in basic research and product development for malaria.
- The malaria R&D funding landscape has been led by investment in drugs (US\$ 2.6 billion, 36% of malaria funding between 2007 and 2018), followed by relatively similar shares for basic research (US\$ 1.9 billion, 26%) and vaccines R&D (US\$ 1.8 billion, 25%). Investments in vector control products and diagnostics were notably lower, reaching overall totals of US\$ 453 million (6.2%) and US\$ 185 million (2.5%), respectively.
- Between 2007 and 2018, the public sector held a leading role in malaria R&D funding, growing from US\$ 246 million in 2007 to a peak of US\$ 365 million in 2017. Within the public sector and among all malaria R&D funders, the US National Institutes of Health was the largest contributor, focusing just over half of its US\$ 1.9 billion investment into basic research (US\$ 1.02 billion, 54% of its overall malaria investment between 2007 and 2018).
- The Bill & Melinda Gates Foundation has been another instrumental player, investing US\$ 1.8 billion (25% of all malaria R&D funding) between 2007 and 2018, and supporting the clinical development of key innovations such as the RTS,S vaccine.

## **DISTRIBUTION AND COVERAGE OF MALARIA PREVENTION**

---

- Manufacturers' delivery data for 2004–2019 show that nearly 2.2 billion insecticide-treated mosquito nets (ITNs) were supplied globally in that period, of which 1.9 billion (86%) were supplied to sub-Saharan Africa.
- Manufacturers delivered about 253 million ITNs to malaria endemic countries in 2019, an increase of 56 million ITNs compared with 2018. About 84% of these ITNs were delivered to countries in sub-Saharan Africa.

- By 2019, 68% of households in sub-Saharan Africa had at least one ITN, increasing from about 5% in 2000. The percentage of households owning at least one ITN for every two people increased from 1% in 2000 to 36% in 2019. In the same period, the percentage of the population with access to an ITN within their household increased from 3% to 52%.
- The percentage of the population sleeping under an ITN also increased considerably between 2000 and 2019, for the whole population (from 2% to 46%), for children aged under 5 years (from 3% to 52%) and for pregnant women (from 3% to 52%).
- The most recent household survey data from demographic and health surveys (DHS) and malaria indicator surveys (MIS) from 24 countries in sub-Saharan Africa from 2015 to 2019 were used to analyse socioeconomic equity in the use of ITNs. In most West African countries, ITN use was generally pro-poor or close to perfect equality. In contrast, ITN use was higher in wealthier households in many parts of Central and East Africa.
- Globally, the percentage of the population at risk protected by indoor residual spraying (IRS) in malaria endemic countries declined from 5% in 2010 to 2% in 2019. The percentage of the population protected by IRS decreased in all WHO regions.
- The number of people protected globally fell from 180 million in 2010 to 115 million in 2015, but declined to 97 million in 2019.
- The number of children reached with at least one dose of seasonal malaria chemoprevention (SMC) steadily increased, from about 0.2 million in 2012 to about 21.5 million in 2019.
- In the 13 countries that implemented SMC, a total of about 21.7 million children were targeted in 2019. On average, 21.5 million children received treatment.
- Using data from 33 African countries, the percentage of IPTp use by dose was computed. In 2019, 80% of pregnant women used ANC services at least once during their pregnancy. About 62% of pregnant women received IPTp1 and 49% received IPTp2. There was a slight increase in IPTp3 coverage, from 31% in 2018 to 34% in 2019.

## **DISTRIBUTION AND COVERAGE OF MALARIA DIAGNOSIS AND TREATMENT**

---

- Globally, 2.7 billion rapid diagnostic tests (RDTs) for malaria were sold by manufacturers in 2010–2019, with nearly 80% of these sales being to sub-Saharan African countries. In the same period, national malaria programmes (NMPs) distributed 1.9 billion RDTs – 84% in sub-Saharan Africa.
- In 2019, 348 million RDTs were sold by manufacturers and 267 million distributed by NMPs. RDT sales and distributions in 2019 were lower than those reported in 2018, by 63 million and 24 million, respectively, with most decreases being in sub-Saharan Africa.
- More than 3.1 billion treatment courses of artemisinin-based combination therapy (ACT) were sold globally by manufacturers in 2010–2019. About 2.1 billion of these sales were to the public sector in malaria endemic countries, and the rest were sold through either public or private sector co-payments (or both), or exclusively through the private retail sector.
- National data reported by NMPs show that, in the same period, 1.9 billion ACTs were delivered to health service providers to treat malaria patients in the public health sector.
- In 2019, some 190 million ACTs were sold by manufacturers for use in the public health sector; in that same year, 183 million ACTs were distributed to this sector by NMPs, of which 90% were in sub-Saharan Africa.
- Aggregated data from household surveys conducted in sub-Saharan Africa between 2005 and 2019 in 21 countries with at least two surveys (baseline 2005–2011, and most recent 2015–2019) in this period were used to analyse coverage of treatment seeking, diagnosis and use of ACTs in children aged under 5 years.

- Comparing the baseline and latest surveys, there was little change in prevalence of fever within the 2 weeks preceding the surveys (median 24% versus 21%) and treatment seeking for fever (median 64% versus 69%).
- Comparisons of the source of treatment between the baseline and more recent surveys show that a median 63% versus 71% received care from public health facilities, and a median 39% versus 30% received care from the private sector. Use of community health workers was low in both periods, at a median of less than 2%.
- The rate of diagnosis among children aged under 5 years for whom care was sought increased considerably, from a median of 15% at baseline to 38% in the latest household surveys.
- Use of ACTs also increased more than threefold, from 39% at baseline to 81% in the latest surveys when all children with fever for whom care was sought were considered.
- Among those who received a finger or heel prick, use of ACTs was 42% in the most recent survey, suggesting that many children received ACTs without parasitological diagnosis.
- Analysis of equity of fever prevalence and treatment seeking at subnational level shows that in most countries, children in poorer households had a higher prevalence of fever in the 2 weeks preceding the household surveys.
- In contrast, treatment seeking was higher in febrile children from wealthier households in all subnational units, although in some units that difference was small.

## BIOLOGICAL THREATS

---

### Parasite deletions of *pfhrp2/3* genes

- Deletions in the *pfhrp2* and *pfhrp3* (*pfhrp2/3*) genes of the parasite renders parasites undetectable by RDTs based on histidine-rich protein 2 (HRP2).
- WHO has recommended that countries with reports of *pfhrp2/3* deletions or neighbouring countries should conduct representative baseline surveys among suspected malaria cases to determine whether the prevalence of *pfhrp2/3* deletions causing false negative RDT results has reached a threshold for RDT change (>5% *pfhrp2* deletions causing false negative RDT results).
- Alternative RDT options (e.g. based on detection of the *Plasmodium* lactate dehydrogenase [pLDH]) are limited; in particular, there are currently no WHO-prequalified non-HRP2 combination tests that can detect and distinguish between *P. falciparum* and *P. vivax*.
- WHO is tracking published reports of *pfhrp2/3* deletions using the Malaria Threats Map mapping tool, and is encouraging a harmonized approach to mapping and reporting of *pfhrp2/3* deletions through publicly available survey protocols.
- Among the 39 reports published by 39 countries, 32 (82%) reported *pfhrp2* deletions; however, variable methods in sample selection and laboratory analysis mean that the scale and scope of clinically significant *pfhrp2/3* deletions is still unclear.
- Between 2019 and September 2020, investigations for *pfhrp2/3* deletions were reported in 16 publications from 15 countries. *Pfhrp2/3* deletions were confirmed in 12 reports from 11 countries: China, Equatorial Guinea, Ethiopia, Ghana, Myanmar, Nigeria, Sudan, Uganda, United Kingdom (imported from various malaria endemic countries), the United Republic of Tanzania and Zambia. No deletions were identified in France (among returning travellers), Haiti, Kenya and Mozambique.

## Parasite resistance to antimalarial drugs

- *PfKelch13* mutations have been identified as molecular markers of partial artemisinin resistance.
- In the WHO African Region, the first-line treatments for *P. falciparum* include artemether-lumefantrine (AL), artesunate–amodiaquine (AS–AQ) and dihydroartemisinin–piperaquine (DHA–PPQ). The overall average efficacy rates for *P. falciparum* – 98.0% for AL, 98.4% for AS–AQ and 99.4% for DHA–PPQ – remained consistent over time. Treatment failure rates of more than 10% were observed in four studies of AL but can be considered statistical outliers. There is no evidence of confirmed lumefantrine resistance in Africa. For all other medicines, treatment failure rates remain below 10%.
- The first-line treatments for *P. falciparum* in the WHO Region of the Americas include AL, artesunate–mefloquine (AS–MQ) and chloroquine (CQ). Efficacy of AL and AS–MQ remains high. One study of CQ from Bolivia (Plurinational State of) in 2011 detected a treatment failure rate of 10.4%.
- The first-line treatments for *P. falciparum* in the WHO South-East Asia Region include AL, artesunate–sulfadoxine–pyrimethamine (AS+SP), and DHA–PPQ. Therapeutic efficacy studies (TES) of AL demonstrated high treatment efficacy in Bhutan, India, Myanmar, Nepal and Timor-Leste. AL treatment failure rates exceeded 10% in three studies, one in Thailand and two in Bangladesh. Following high rates of AS+SP treatment failure in the north-eastern provinces, in 2013, India changed its treatment policy in those provinces to AL; AS+SP remains effective elsewhere in the country. TES findings in Thailand led to the adoption of DHA–PPQ as the first-line treatment in 2015. In Thailand, moderate to high rates of treatment failure were observed with DHA–PPQ in the eastern part of the country; thus, Thailand is currently recommending treatment with artesunate–pyronaridine (AS–PY) in this area.
- AL and AS+SP remain efficacious in the countries that use them as first-line treatment in the WHO Eastern Mediterranean Region.
- The first-line treatments for *P. falciparum* in the WHO Western Pacific Region are AL in all malaria endemic countries except China, where AS–AQ is used. AL treatment failure rates were 10% or less in four studies in Lao People’s Democratic Republic, but those studies did not have the recommended sample sizes. A study with an adequate number of patients is currently underway to further investigate these high rates of treatment failure.
- Artemisinin partial resistance emerged independently in several foci in the GMS. WHO continues to monitor the situation, which has evolved rapidly since the first detections of *PfKelch13* mutations in the GMS. Some mutations have disappeared, whereas the prevalence of others has increased.
- Currently, the most prevalent markers west of Bangkok (western Thailand and Myanmar) are F446I, M476I and R561H. The most prevalent markers east of Bangkok (eastern Thailand, Cambodia, Lao People’s Democratic Republic and Viet Nam) are Y493H and P553L. Two markers, R539T and C580Y, are also highly prevalent in both areas. The change in treatment policy in Cambodia from DHA–PPQ to AS–MQ resulted in a reduction in the prevalence of strains carrying both C580Y and PPQ resistance.
- Rwanda has detected an increasing prevalence of the R561H mutation, a validated marker that emerged independently in the GMS between 2012 and 2015. The presence of this mutation was confirmed in Rwanda in 2018; however, so far it seems that delayed clearance associated with this mutation has not affected the efficacy of the ACTs that are currently among those tested and used in Rwanda.
- The R622I mutation seems to be appearing independently in Africa, having been found in Eritrea, Ethiopia, Somalia and Sudan, and with increasing frequency in the Horn of Africa. The ACTs used in these four countries remain effective, despite the presence of the mutation. Further investigation of delayed parasite clearance is needed in this region.
- In Guyana, the C580Y mutation also emerged independently between 2010 and 2017. However, in recent studies (including surveys and TES), 100% of samples were found to be wild type, indicating that the mutation may be disappearing in Guyana.

## Vector resistance to insecticides

- From 2010 to 2019, some 81 countries reported data on standard insecticide resistance monitoring to WHO.
- Concerningly, between 2010 and 2019, 57% of the countries that reported using IRS did not report the status of insecticide resistance for every insecticide class used in the year of implementation or the preceding one, and 14% did not report the status of resistance for any insecticide class used. Malaria endemic countries are highly encouraged to ensure adequate monitoring of insecticide resistance to classes that are in use or under consideration for use in malaria vector control interventions, and to prioritize monitoring these classes.
- Of the 82 malaria endemic countries that provided data for 2010–2019, 28 have detected resistance to all four of the most commonly used insecticide classes in at least one malaria vector and one collection site, and 73 have detected resistance to at least one insecticide class. Only eight countries have not detected resistance to any insecticide class so far.
- Globally, resistance to pyrethroids – the only insecticide class currently used in ITNs – continues to be widespread. It was detected in at least one malaria vector in 69.9% of the sites for which data were available. Resistance to organochlorines was reported in 63.4% of the sites. Resistance to carbamates and organophosphates was less prevalent, being detected in 31.7% and 24.9% of the sites that reported monitoring data, respectively.
- Based on insecticide resistance monitoring data reported to WHO by Member States, a total of 330 areas in 33 countries currently meet the WHO–recommended criteria for the deployment of pyrethroid–piperonyl butoxide nets.
- Although WHO Member States and their implementing partners have started to report insecticide resistance monitoring data for neonicotinoids and pyrroles, Member States are discouraged from using data generated by means of non–validated procedures to arrive at conclusions about the resistance status of their local vector populations to these insecticide classes. A formal WHO process to establish discriminating dosages and test procedures for these two insecticide classes is ongoing. The data reported to WHO will be evaluated according to these dosages and procedures as they become available.
- To guide resistance management, countries should develop and implement a national plan for insecticide resistance monitoring and management, drawing on the WHO *Framework for a national plan for monitoring and management of insecticide resistance in malaria vectors*. In 2019, the number of countries that had completed such plans rose to 53, and 29 countries were in the process of developing them.
- Standard insecticide resistance data reported to WHO are included in the WHO global database on insecticide resistance in malaria vectors and are available for exploration via the Malaria Threats Map. A new version of this tool with enhanced functionality and data download options was released in 2020.



## MALARIA RESPONSE DURING THE COVID-19 PANDEMIC

---

- By April 2020, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV2), causing COVID-19, had spread to all malaria endemic countries, and by the end of the second week of November 2020, about 22 million cases and 600 000 deaths had been reported in these countries.
- The COVID-19 pandemic and restrictions related to the response have caused disruptions in essential malaria services.
- Furthermore, early messaging targeted at reducing coronavirus transmission advised the public to stay at home if they had fever, potentially disrupting treatment seeking for febrile diseases such as malaria.
- In March 2020, as the COVID-19 pandemic spread rapidly around the globe, WHO convened a cross-partner effort to mitigate the negative impact of the coronavirus in malaria-affected countries and contribute to the COVID-19 response.
- The work was carried out in close collaboration with the RBM Partnership to End Malaria, the Global Fund, the US President's Malaria Initiative (PMI), several implementation and advocacy partners, and research institutions.
- The cross-partner effort led to a strong partnership alignment that resulted in various outcomes:
  - publication of technical guidance on how to safely maintain malaria control services in the context of the COVID-19 pandemic;
  - publication of a modelling analysis to quantify the potential impact of service disruptions due to the COVID-19 pandemic, to reinforce the consequences of service disruption; the analysis suggested that malaria mortality in sub-Saharan Africa was likely to double by the end of 2020, relative to a 2018 baseline, if extreme disruption in prevention and treatment occurred;
  - mitigating the pressure to shift diagnostic production away from malaria to the detection of SARS-CoV2;
  - success in resolving major global manufacturing bottlenecks for malaria medicines;
  - mitigating the disruptions in the shipment and delivery of malaria commodities;
  - resource mobilization for personal protective equipment (PPE) and other commodities to help with the implementation of prevention campaigns, diagnosis and treatment; and
  - tracking of disruptions in countries to help guide the response.
- The collective effort has led to impressive efforts by countries to complete malaria prevention campaigns involving long-lasting insecticidal nets (LLINs), IRS and SMC, and to minimize disruptions to diagnosis and treatment.
- All countries that had planned SMC campaigns were on track to complete them, despite moderate delays in some areas.
- Of the 47 countries that had IRS campaigns planned in 2020, 23 had completed them, 13 were on track to complete them, and 11 were off track or at risk of not completing them.
- Several countries have completed their LLIN campaigns and many are in the process of distributing LLINs. However, as of the third week of November, of the 222 million LLINs planned for distribution in 2020, about 105 million had been distributed.
- Many countries have also reported moderate levels of disruptions, and modelling analysis shows that reductions in access to effective antimalarial treatment of 10%, 15%, 25% and 50% in sub-Saharan Africa in 2020 could lead to an additional 19 000, 28 000, 46 000 and 100 000 malaria deaths, respectively, by the end of 2020, even if all prevention campaigns are completed.

# Avant-propos



**Dr Tedros Adhanom Ghebreyesus**  
Directeur général  
de l'Organisation mondiale de la Santé (OMS)

Dans le *Rapport sur le paludisme dans le monde* de cette année, l'OMS se penche sur les principales étapes ayant marqué la riposte mondiale contre cette maladie au cours des deux dernières décennies et qui ont abouti à une période de succès sans précédent permettant d'éviter 1,5 milliard de cas et 7,6 millions de décès associés.

À l'issue du Programme mondial d'éradication du paludisme en 1969, le désengagement politique et la baisse des financements ont entraîné une résurgence de la maladie dans de nombreuses régions du monde, en particulier en Afrique. Même si les données fiables sont rares, des centaines de millions de personnes ont vraisemblablement été infectées par le paludisme et des dizaines de millions en sont mortes.

Au début des années 1990, les principaux dirigeants des services de santé et experts scientifiques ont tracé les grandes lignes d'une nouvelle réponse au paludisme. Des investissements accrus dans la recherche et l'innovation ont conduit au développement de nouveaux outils de lutte contre la maladie, notamment des moustiquaires imprégnées d'insecticide, des tests de diagnostic rapide et des médicaments plus efficaces.

Associée à une nette augmentation des investissements dans la lutte contre le paludisme, la création de nouveaux mécanismes de financement, notamment le Fonds mondial de lutte contre le sida, la tuberculose et le paludisme, et l'Initiative du Président américain contre le paludisme (PMI), a permis le déploiement à grande échelle de ces nouveaux outils, et a contribué à réduire morbidité et mortalité liées au paludisme dans des proportions inédites jusqu'alors.

Un engagement politique ferme dans les pays d'endémie palustre a constitué la clé du succès. En signant la Déclaration d'Abuja en 2000, une étape historique, les dirigeants des pays africains se sont engagés à réduire de 50 % la mortalité due au paludisme sur le continent en dix ans.

D'après notre rapport, la mortalité associée au paludisme a diminué de 60 % au niveau mondial entre 2000 et 2019. La région Afrique a enregistré une impressionnante baisse du nombre de décès annuels, passant de 680 000 en 2000 à 384 000 en 2019.

Les pays de la région Asie du Sud-Est ont également accompli de sérieux progrès, en réduisant les nombres de cas et de décès de 73 % et 74 %, respectivement. Dans cette région, l'Inde a contribué à la plus forte baisse du nombre de cas, passant de quasiment 20 millions à 6 millions de cas pendant cette période.

Vingt-un pays ont éliminé le paludisme au cours des deux dernières décennies et dix d'entre eux ont été officiellement certifiés exempts de paludisme par l'OMS. Les pays de la sous-région du Grand Mékong continuent à réaliser des avancées majeures, avec un recul de 97 % des infections à *P. falciparum* depuis 2000, un objectif prioritaire compte tenu de la menace permanente que fait peser la résistance aux médicaments antipaludiques.

## Stagnation des progrès

Les progrès enregistrés depuis le début du millénaire sont vraiment stupéfiants. Toutefois, comme le décrit ce rapport, ils stagnent depuis plusieurs années.

En 2017, l'OMS avait souligné que la lutte contre le paludisme au niveau mondial était arrivée à la « croisée des chemins » et que les cibles essentielles de la stratégie mondiale contre le paludisme de l'OMS ne seraient probablement pas atteintes. Trois ans plus tard, les progrès stagnent toujours. Selon notre dernier rapport, les cibles en matière de baisse de l'incidence et de la mortalité liée au paludisme, telles que définies par la stratégie pour 2020, seront respectivement manquées de 37 % et de 22 %.

En 2020, la COVID-19 est venue s'ajouter aux obstacles de taille que la riposte contre le paludisme doit affronter au niveau mondial. Conformément aux orientations de l'OMS, de nombreux pays ont adapté leurs méthodes de distribution de moustiquaires, diagnostics et médicaments afin d'assurer la sécurité des agents de santé et des communautés en première ligne. Je salue du fond du cœur ces efforts, sans lesquels nous aurions sans doute observé des taux de mortalité beaucoup plus élevés.

Les nouvelles projections de l'OMS montrent néanmoins que des dysfonctionnements, même modérés, de l'accès aux traitements antipaludiques efficaces pourraient entraîner un nombre considérable de décès. Le rapport insiste, par exemple, sur le fait qu'un dysfonctionnement à hauteur de 25 % de l'accès au traitement antipaludique efficace en Afrique subsaharienne pourrait entraîner 46 000 décès supplémentaires.

## Relance des progrès

Afin de redynamiser les progrès, l'OMS et le Partenariat RBM pour en finir avec le paludisme ont initié, en 2018, l'approche « *high burden to high impact* » (HBHI, « D'une charge élevée à un fort impact »). Cette approche est menée par 11 pays, dont 10 en Afrique subsaharienne, qui concentrent près de 70 % des cas et décès dus au paludisme dans le monde.

Les pays de l'approche HBHI ont abandonné l'idée d'une démarche « universelle », choisissant au contraire d'utiliser des données et informations collectés localement pour mettre en œuvre des réponses adaptées. Même s'il est trop tôt pour évaluer l'impact de cette approche sur la charge palustre, un important travail préparatoire a été réalisé.

Une récente analyse menée au Nigéria a révélé, par exemple, que le pays pourrait éviter des dizaines de millions de cas et des milliers de décès supplémentaires d'ici 2023 en optant pour une combinaison optimisée d'interventions plutôt qu'en recourant à une approche habituelle.

Un meilleur ciblage des ressources et des interventions antipaludiques, notamment dans des pays où la maladie sévit le plus, comme au Nigéria, va aider à accélérer le rythme des progrès vers les cibles de la stratégie mondiale de lutte contre le paludisme. Il est indispensable d'accroître les financements nationaux et internationaux, et d'innover dans le domaine des outils et des approches.

Sur la voie d'une couverture de santé universelle dans chaque pays, il est aussi essentiel d'intégrer les efforts de lutte contre le paludisme aux initiatives plus larges visant à mettre en place des systèmes de santé solides, basés sur des soins de santé primaires axés sur la personne.

Il est temps pour les dirigeants de toute l'Afrique, mais aussi du monde entier, de relever le défi du paludisme une fois encore, comme ils l'avaient fait lorsqu'ils ont jeté les bases des avancées réalisées depuis le début de ce siècle. À travers une action commune et un engagement à n'oublier personne, nous pourrons concrétiser notre vision partagée d'un monde sans paludisme.



# Le rapport de cette année en un clin d'œil

## POIDS DU PALUDISME : ÉVOLUTION DU NOMBRE DE CAS ET DE DÉCÈS

---

### Cas de paludisme

- Au niveau mondial, le nombre de cas de paludisme est estimé à 229 millions en 2019 dans 87 pays d'endémie palustre, soit une baisse par rapport aux 238 millions de 2000. Lors de la définition de la *Stratégie technique mondiale de lutte contre le paludisme 2016-2030* ([le] GTS) en 2015, le nombre de cas de paludisme était estimé à 218 millions.
- Le pourcentage des infections à *Plasmodium vivax* a diminué, passant de 7 % en 2000 à 3 % en 2019.
- L'incidence du paludisme (i.e. nombre de cas pour 1 000 habitants exposés au risque de paludisme) a reculé au niveau mondial, passant de 80 en 2000 à 58 en 2015, puis 57 en 2019. De 2000 à 2015, l'incidence du paludisme au niveau mondial a donc diminué de 27 %, mais de 2 % seulement entre 2015 et 2019, ce qui reflète un net ralentissement depuis 2015.
- Vingt-neuf pays ont concentré 95 % du nombre total de cas de paludisme dans le monde. Le Nigéria (27 %), la République démocratique du Congo (12 %), l'Ouganda (5 %), le Mozambique (4 %) et le Niger (3 %) ont enregistré, à eux seuls, près de 51 % des cas.
- La région Afrique de l'Organisation mondiale de la Santé (OMS) représente à elle seule 94 % (215 millions) des cas estimés en 2019.
- Dans la région Afrique de l'OMS, même si le nombre de cas de paludisme était moins élevé (204 millions) en 2000 qu'en 2019, l'incidence du paludisme a baissé de 363 à 225 cas pour 1 000 habitants exposés au risque de paludisme sur cette période, ce qui traduit la complexité d'interpréter l'évolution de la transmission de la maladie au sein d'une population qui ne cesse de croître. La population vivant dans la région Afrique de l'OMS est passée de 665 millions en 2000 à 1,1 milliard en 2019.
- La région Asie du Sud-Est de l'OMS a concentré près de 3 % des cas de paludisme dans le monde. Le nombre de cas y a chuté de 73 %, passant de 23 millions en 2000 à près de 6,3 millions en 2019. De même, l'incidence du paludisme dans cette région a diminué de 78 %, avec quelque 18 cas pour 1 000 habitants exposés au risque de paludisme en 2000, contre 4 en 2019.
- Dans la région Asie du Sud-Est de l'OMS, l'Inde a enregistré la baisse la plus prononcée, en valeur absolue, avec près de 20 millions de cas en 2000, contre 5,6 millions environ en 2019. Le Sri Lanka a été certifié exempt de paludisme en 2015, et le Timor-Leste a rapporté zéro cas de paludisme en 2018 et 2019.
- Dans la région Méditerranée orientale de l'OMS, le nombre de cas de paludisme a baissé de 26 %, passant de près de 7 millions en 2000 à quelque 5 millions en 2019. Près d'un quart de ces cas en 2019 étaient dus à *P. vivax*, principalement en Afghanistan et au Pakistan.
- Sur la période 2000-2019, l'incidence du paludisme dans la région Méditerranée orientale de l'OMS a diminué de 20 à 10. Avec quasiment 46 % des cas, le Soudan est le pays le plus touché dans cette région. La République islamique d'Iran a rapporté zéro cas de paludisme indigène en 2018 et 2019.
- Dans la région Pacifique occidental de l'OMS, 1,7 million de cas ont été estimés en 2019, soit une baisse de 43 % par rapport aux 3 millions de 2000. Sur la même période, l'incidence du paludisme est passée de cinq à deux cas pour 1 000 habitants exposés au risque de paludisme. La Papouasie-Nouvelle-Guinée a enregistré près de 80 % des cas dans cette région en 2019. Depuis 2017, la Chine rapporte zéro cas de paludisme indigène. La Malaisie n'a rapporté aucun cas de paludisme humain en 2018 et 2019.
- Dans la région Amériques de l'OMS, le nombre de cas de paludisme a diminué de 40 % (passant de 1,5 million à 0,9 million) et l'incidence du paludisme de 57 % (de 14 à 6). Les progrès réalisés dans cette région ces dernières années ont souffert de la forte hausse du paludisme au Venezuela (République bolivarienne du), qui avait recensé près de 35 500 cas en 2000 contre plus de 467 000 en 2019. Le

Brésil, la Colombie et le Venezuela (République bolivarienne du) concentrent plus de 86 % des cas dans cette région.

- Depuis 2015, la région Europe de l'OMS est exempte de paludisme.

### **Mortalité associée**

- Au niveau mondial, le nombre de décès dus au paludisme a baissé de façon régulière sur la période 2000-2019, passant de 736 000 en 2000 à 409 000 en 2019. Les enfants de moins de 5 ans représentaient 84 % des décès associés au paludisme en 2000, contre 67 % en 2019. L'estimation du nombre de décès dans le monde en 2015, la référence du GTS, avoisinait les 453 000.
- La mortalité associée au paludisme (à savoir le nombre de décès pour 100 000 habitants exposés au risque de paludisme) a baissé au niveau mondial, passant de 25 en 2000 à 12 en 2015, puis 10 en 2019, ce qui traduit un ralentissement de tendance ces dernières années.
- Au niveau mondial, près de 95 % des décès dus au paludisme ont été enregistrés dans 31 pays. Le Nigéria (23 %), la République démocratique du Congo (11 %), la République-Unie de Tanzanie (5 %), le Mozambique (4 %), le Niger (4 %) et le Burkina Faso (4 %) ont concentré près de 51 % de tous les décès dus au paludisme dans le monde en 2019.
- Dans la région Afrique de l'OMS, le nombre de décès dus au paludisme a diminué de 44 %, passant de 680 000 en 2000 à 384 000 en 2019. Sur la même période, la mortalité associée a baissé de 67 %, chutant de 121 à 40 décès pour 100 000 habitants exposés au risque de paludisme.
- Dans la région Asie du Sud-Est de l'OMS, le nombre de décès dus au paludisme a diminué de 74 %, avec 35 000 décès en 2000 contre 9 000 en 2019.
- L'Inde a concentré près de 86 % des décès dus au paludisme dans la région Asie du Sud-Est de l'OMS.
- Dans la région Méditerranée orientale de l'OMS, le nombre de décès dus au paludisme a diminué de 16 %, passant de 12 000 en 2000 à 10 100 en 2019. Dans le même temps, la mortalité associée a baissé de moitié, passant de quatre à deux décès pour 100 000 habitants exposés au risque de paludisme.
- Dans la région Pacifique occidentale de l'OMS, le nombre de décès dus au paludisme a diminué de 52 %, passant de 6 600 en 2000 à 3 200 en 2019. Sur la même période, la mortalité associée a baissé de 60 %, chutant de 1 à 0,4 décès pour 100 000 habitants exposés au risque de paludisme. Dans cette région, la Papouasie-Nouvelle-Guinée a enregistré près de 85 % des décès dus au paludisme en 2019.
- Dans la région Amériques de l'OMS, le nombre de décès dus au paludisme a diminué de 39 % (909 contre 551) et la mortalité associée de 50 % (0,8 contre 0,4). Plus de 70 % des décès dus au paludisme en 2019 dans cette région ont été enregistrés au Venezuela (République bolivarienne du).

### **Nombre de cas de paludisme et de décès évités**

- Selon les estimations, 1,5 milliard de cas de paludisme et 7,6 millions de décès associés ont été évités dans le monde entre 2000 et 2019.
- La plupart des cas (82 %) et des décès (94 %) évités l'auraient été dans la région Afrique de l'OMS, suivie par la région Asie du Sud-Est (10 % des cas et 3 % des décès).

### **Poids du paludisme pendant la grossesse**

- En 2019, sur les 33 millions de femmes enceintes vivant dans 33 pays de la région Afrique de l'OMS où la transmission est modérée à élevée, 35 % (soit 12 millions) ont été exposées à une infection palustre durant leur grossesse.
- En détaillant les sous-régions de l'OMS, l'Afrique centrale a affiché la plus forte prévalence d'exposition au paludisme durant la grossesse (40 %), suivie de près par l'Afrique de l'Ouest (39 %), alors que la prévalence était de 24 % en Afrique de l'Est et en Afrique australe.
- Conséquence de ces infections pendant la grossesse, 822 000 enfants ont présenté un faible poids à la naissance dans ces 33 pays.
- Si 80 % des femmes enceintes ayant reçu des soins prénataux avaient reçu une dose de traitement préventif intermittent pendant la grossesse (TPIp), 56 000 cas de faible poids à la naissance auraient été évités dans ces 33 pays.

## ÉLIMINATION DU PALUDISME ET PRÉVENTION DE SA RÉAPPARITION

---

- Au niveau mondial, le nombre de pays où le paludisme était endémique en 2000 et qui ont rapporté moins de 10 000 cas a augmenté, passant de 26 en 2000 à 46 en 2019.
- Au cours de la même période, les pays comptant moins de 100 cas de paludisme indigène sont passés de 6 à 27.
- Sur la période 2010–2019, le nombre total de cas de paludisme dans les 21 pays de l’initiative « E-2020 » a diminué de 79 %.
- Les Comores, le Costa Rica, l’Équateur et le Suriname ont signalé plus de cas en 2019 qu’en 2018, avec respectivement 1 986, 25, 150 et 66 cas supplémentaires en 2019.
- La République islamique d’Iran, la Malaisie et le Timor-Leste ont rapporté zéro cas de paludisme indigène en 2018 et 2019. En 2019, le Belize et le Cabo Verde n’ont signalé aucun cas de paludisme indigène pour la première fois depuis 2000.
- La Chine et El Salvador ont rapporté zéro cas de paludisme indigène pour la troisième année consécutive et ont donc déposé une demande formelle de certification.
- Dans les six pays de la sous-région du Grand Mékong (Cambodge, Chine [province du Yunnan], République démocratique populaire lao, Myanmar, Thaïlande et Viet Nam), le nombre de cas de paludisme à *P. falciparum* a diminué de 97 % entre 2000 et 2019, alors que le nombre total de cas a chuté de 90 %. Sur les 239 000 cas de paludisme rapportés en 2019, 65 000 étaient dus à *P. falciparum*.
- Ce recul s’est accéléré depuis 2012, date à laquelle le programme « Mekong Malaria Elimination » (MME) a été lancé. Durant cette période, le nombre de cas de paludisme a été divisé par six et les cas dus à *P. falciparum* par 14 ou presque.
- Dans l’ensemble, le Cambodge (58 %) et le Myanmar (31 %) ont concentré une large majorité des cas de paludisme dans la sous-région du Grand Mékong.
- Cette accélération de la baisse des cas dus à *P. falciparum* est particulièrement importante du fait de la résistance accrue aux médicaments. En effet, dans la sous-région du Grand Mékong, les parasites *P. falciparum* ont développé une résistance partielle à l’artémisinine, le composant principal des meilleurs médicaments antipaludiques disponibles.
- De 2000 à 2019, la transmission du paludisme n’est réapparue dans aucun des pays préalablement certifiés exempts de paludisme.

## APPROCHE « HIGH BURDEN TO HIGH IMPACT »

---

- Depuis novembre 2018, l’approche « high burden to high impact » (HBHI) a été lancée dans 10 des 11 pays concernés (elle n’a pas encore été lancée au Mali en raison des dysfonctionnements liés à la pandémie de COVID-19). Toutefois, ces 11 pays ont déjà mis en place des activités HBHI en rapport avec les quatre éléments de riposte définis.
- Dans chaque pays HBHI, le lancement a fait l’objet d’un engagement politique à haut niveau et de soutien important. L’initiative « Mass Action Against Malaria » en Ouganda est citée à titre d’exemple de processus mené par un pays avec un engagement politique à tous les niveaux, ainsi qu’une mobilisation communautaire et multisectorielle.
- L’analyse de l’adaptation infranationale des interventions a été réalisée dans tous les pays, sauf au Mali où elle est en cours. L’exemple du Nigéria est présenté dans le rapport.
- Tous les pays se sont engagés à conduire un exercice exhaustif de microstratification urbaine afin de mieux cibler les interventions et d’améliorer leur efficacité en tenant compte de l’urbanisation croissante.
- Le programme mondial de lutte antipaludique de l’OMS a actualisé son dossier technique pour aider les pays à mieux prioriser les ressources, tout en respectant les recommandations développées dans le cadre des processus normalisés et rigoureux de l’OMS.
- Comme l’approche HBHI a été lancée en novembre 2018, à une période où les pays arrivaient à la fin de leurs cycles de financement, il est trop tôt pour déterminer l’impact de la réponse. En 2019, le nombre de cas de paludisme dans les 11 pays HBHI était similaire à celui de 2018 (156 millions contre 155 millions).

## PROGRÈS VERS L'ATTEINTE DES OBJECTIFS DU GTS POUR 2020

---

- Le GTS vise à réduire l'incidence du paludisme et la mortalité associée d'au moins 40 % d'ici 2020, 75 % d'ici 2025 et 90 % d'ici 2030 en se basant sur les données de référence de 2015.
- Les tendances 2000-2019 concernant le nombre de cas de paludisme et de décès associés ont servi à établir des projections annuelles de 2020 à 2030, afin de suivre les progrès sur la voie des cibles et des objectifs intermédiaires du GTS.
- Les projections présentées dans le rapport *ne tiennent pas compte* des éventuels dysfonctionnements dus à la pandémie de COVID-19, lesquels risquent d'entraîner une morbidité et une mortalité liées au paludisme plus élevées que prévu, malgré les efforts remarquables consentis au niveau national et international pour préserver les services de base en matière de lutte contre le paludisme.
- En dépit des progrès considérables accomplis depuis 2000, les objectifs intermédiaires du GTS pour 2020 en matière de morbidité et de mortalité ne seront pas atteints au niveau mondial.
- En 2020, l'incidence du paludisme s'est établie à 56 cas pour 1 000 habitants à risque, au lieu des 35 cas représentés par l'objectif intermédiaire de morbidité fixé dans le GTS. En d'autres termes, nous sommes à 37 % en deçà de notre objectif.
- Même si la baisse de la mortalité est plus nette que la baisse de l'incidence, la projection du nombre de décès pour 100 000 habitants exposés au risque de paludisme a été établie au niveau mondial à 9,8 en 2020 contre 11,9 en 2015, soit un écart de 22 % par rapport à l'objectif intermédiaire de mortalité défini dans le GTS pour 2020.
- Sur les 92 pays où le paludisme était endémique en 2015, 31 (34 %) étaient en passe d'atteindre l'objectif intermédiaire pour 2020 en matière de morbidité. En effet, selon les estimations, ils ont réduit leur incidence de 40 % ou plus, ou ont rapporté zéro cas de paludisme.
- Vingt-deux pays (23 %) ont réalisé des progrès en termes de baisse de l'incidence, mais pas suffisamment pour atteindre l'objectif intermédiaire du GTS.
- Trente-et-un pays (34 %) ont enregistré une hausse de l'incidence, et elle était supérieure ou égale à 40 % dans 15 (16 %) d'entre eux par rapport à 2015.
- Dans neuf pays (10 %), l'incidence du paludisme en 2020 a été estimée à un niveau équivalent à celui de 2015.
- Trente-neuf pays (42 %) où le paludisme était endémique en 2015 étaient en passe d'atteindre l'objectif intermédiaire du GTS pour 2020 en matière de mortalité, et 28 d'entre eux ont rapporté zéro cas de paludisme.
- Selon les estimations, trente-quatre pays (37 %) ont réduit la mortalité due au paludisme, mais leurs progrès sont restés en-deçà de l'objectif de 40 %.
- En 2020, la mortalité due au paludisme est restée au même niveau qu'en 2015 dans sept pays (8 %), alors que 12 autres pays (13 %) semblent avoir enregistré des hausses, et même de 40 % ou plus dans six pays.
- Tous les pays de la région Asie du Sud-Est de l'OMS étaient en passe d'atteindre les objectifs intermédiaires du GTS à la fois en matière de morbidité et de mortalité pour 2020.

## INVESTISSEMENTS DANS LES PROGRAMMES ET LA RECHERCHE ANTIPALUDIQUES

---

- Le GTS estime les fonds requis pour atteindre les objectifs intermédiaires de 2020, 2025 et 2030. Au total, les ressources annuelles nécessaires ont été estimées à US\$ 4,1 milliards en 2016, avec une hausse à US\$ 6,8 milliards en 2020. Toujours selon les estimations, US\$ 720 000 millions supplémentaires seront requis chaque année pour la recherche et le développement (R&D) sur le paludisme au niveau mondial.
- En 2019, US\$ 3 milliards ont été investis au total pour le contrôle et l'élimination du paludisme, contre US\$ 2,7 milliards en 2018 et US\$ 3,2 milliards en 2017. Les investissements de 2019 sont bien inférieurs aux US\$ 5,6 milliards estimés nécessaires au niveau mondial pour rester sur la voie des objectifs du GTS.

- L'écart entre investissements et ressources nécessaires a continué à augmenter de façon spectaculaire au cours de ces dernières années, passant de US\$ 1,3 milliard en 2017 à US\$ 2,3 milliards en 2018, puis US\$ 2,6 milliards en 2019.
- Les partenaires internationaux ont représenté 70 % du financement total pour le contrôle et l'élimination du paludisme sur la période 2010-2019, avec les États-Unis en tête, suivis par le Royaume-Uni de Grande-Bretagne et d'Irlande du Nord (Royaume-Uni), et la France.
- Sur les US\$ 3 milliards investis en 2019, US\$ 2,1 milliards provenaient de bailleurs de fonds internationaux. Le gouvernement des États-Unis a été le premier bailleur de fonds en 2019, apportant US\$ 1,1 milliard au travers de financements bilatéraux planifiés et de contributions à des agences de financement multilatérales.
- Des décaissements bilatéraux et multilatéraux du Royaume-Uni à hauteur de US\$ 200 millions sont venus s'ajouter à ces financements, des contributions de plus de US\$ 100 millions de la part de la France, de l'Allemagne et du Japon (pour un total de US\$ 400 millions), ainsi que US\$ 400 millions supplémentaires de la part d'autres pays membres du Comité d'aide au développement et de bailleurs de fonds du secteur privé.
- En 2019, les gouvernements des pays d'endémie ont contribué à hauteur de 30 % du financement total, soit près de US\$ 900 millions. Sur ce montant, US\$ 200 millions ont été investis dans la prise en charge des cas de paludisme dans le secteur public et US\$ 700 millions dans d'autres activités de lutte contre le paludisme.
- Sur les US\$ 3 milliards investis en 2019, près de US\$ 1,2 milliard (39 %) ont transité par le Fonds mondial de lutte contre le sida, la tuberculose et le paludisme (Fonds mondial). Par rapport à 2018, les décaissements du Fonds mondial en faveur des pays d'endémie ont augmenté de près de US\$ 200 millions en 2019.
- Sur les US\$ 3 milliards investis en 2019, près de 73 % ont été dirigés vers la région Afrique de l'OMS, 9 % vers la région Asie du Sud-Est, 5 % vers les régions Amériques et Pacifique occidental (chacune), et 4 % vers la région Méditerranée orientale.
- De 2007 à 2018, près de US\$ 7,3 milliards ont été investis dans la recherche fondamentale et le développement de produits contre le paludisme.
- Les fonds dédiés à la recherche-développement ont surtout été investis dans les médicaments (US\$ 2,6 milliards, soit 36 % des fonds investis entre 2007 et 2018), suivis à parts relativement proches par la recherche fondamentale (US\$ 1,9 milliard, soit 26 %) et la recherche-développement dans le domaine des vaccins (US\$ 1,8 milliard, soit 25 %). Les investissements dans les produits de lutte antivectorielle et les outils de diagnostic ont été nettement plus modérés, atteignant globalement US\$ 453 millions (6,2 %) et US\$ 185 millions (2,5 %), respectivement.
- Entre 2007 et 2018, le secteur public a tenu un rôle majeur dans le financement de la recherche-développement antipaludique, passant de US\$ 246 millions en 2007 à un pic de US\$ 365 millions en 2017. Au sein du secteur public et parmi tous les bailleurs de fonds engagés dans la recherche-développement antipaludique, les US National Institutes of Health ont apporté la contribution la plus importante, en concentrant un peu plus de la moitié de leurs investissements de US\$ 1,9 milliard dans la recherche fondamentale (soit US\$ 1,02 milliard ou 54 % de leurs investissements totaux dans la lutte contre le paludisme entre 2007 et 2018).
- La Fondation Bill & Melinda Gates a également tenu un rôle important, en investissant US\$ 1,8 milliard (soit 25 % de tous les financements de recherche-développement antipaludique) entre 2007 et 2018, ainsi qu'en soutenant le développement clinique d'innovations essentielles, comme le vaccin RTS,S.

## **DISTRIBUTION ET COUVERTURE DES OUTILS DE PRÉVENTION DU PALUDISME**

- Les fabricants de moustiquaires imprégnées d'insecticide (MII) ont indiqué en avoir livré près de 2,2 milliards dans le monde entre 2004 et 2019, dont 1,9 milliard (86 %) en Afrique subsaharienne.
- En 2019, ces fabricants ont livré près de 253 millions de MII à des pays d'endémie, soit une augmentation de 56 millions par rapport à 2018. Près de 84 % de ces MII ont été livrées dans des pays d'Afrique subsaharienne.



- En 2019, 68 % des ménages vivant en Afrique subsaharienne disposaient d'au moins une MII, soit une hausse d'environ 5 % par rapport à 2000. Le pourcentage des ménages disposant d'au moins une MII pour 2 membres du foyer est passé de 1 % en 2000 à 36 % en 2019. Durant la même période, le pourcentage de la population ayant accès à une MII dans son foyer a augmenté, passant de 3 % à 52 %.
- Le pourcentage de la population dormant sous MII a aussi considérablement augmenté entre 2000 et 2019, qu'il s'agisse de la population dans son ensemble (de 2 % à 46 %), des enfants de moins de 5 ans (de 3 % à 52 %) ou des femmes enceintes (de 3 % à 52 %).
- Les données les plus récentes, issues d'enquêtes démographiques et de santé et d'autres enquêtes sur les indicateurs du paludisme réalisées au sein des ménages dans 24 pays d'Afrique subsaharienne entre 2015 et 2019, ont servi à analyser l'équité socio-économique concernant l'utilisation des MII. Dans la plupart des pays d'Afrique de l'Ouest, l'utilisation des MII a été d'une manière générale plus importante parmi les plus démunis, ou alors homogène parmi les différents quintiles de richesse. À l'inverse, dans de nombreuses régions d'Afrique centrale et d'Afrique de l'Est, l'utilisation des MII a été supérieure au sein des ménages les moins démunis.
- Au niveau mondial, la part de la population à risque protégée par pulvérisation intradomiciliaire d'insecticides à effet rémanent (PID) dans les pays d'endémie a reculé de 5 % en 2010 à 2 % en 2019. Le pourcentage de la population protégée par PID a diminué dans toutes les régions de l'OMS.
- Au niveau mondial, le nombre de personnes protégées par cette intervention a chuté de 180 millions en 2010 à 115 millions en 2015, puis à 97 millions en 2019.
- Le nombre d'enfants ayant reçu au moins une dose de chimioprévention du paludisme saisonnier (CPS) n'a cessé d'augmenter, passant de quelque 0,2 million en 2012 à près de 21,5 millions en 2019.
- Dans les 13 pays ayant mis en œuvre la CPS, quelque 21,7 millions d'enfants au total ont été ciblés en 2019. En moyenne, 21,5 millions d'enfants ont reçu un traitement.
- Le pourcentage d'utilisation du TPIp par dose a été calculé sur la base des données provenant de 33 pays d'Afrique. En 2019, 80 % des femmes enceintes ont reçu des soins prénataux au moins une fois durant leur grossesse. Environ 62 % des femmes enceintes ont reçu une dose de TPIp, et 49 % ont reçu deux doses. La couverture en TPIp par trois doses a légèrement augmenté, passant de 31 % en 2018 à 34 % en 2019.

## **DISTRIBUTION ET COUVERTURE DES OUTILS DE DIAGNOSTIC ET DE TRAITEMENT DU PALUDISME**

- De 2010 à 2019, 2,7 milliards de tests de diagnostic rapide (TDR) du paludisme ont été vendus dans le monde, dont 80 % à destination des pays d'Afrique subsaharienne. Durant la même période, 1,9 milliard de TDR ont été distribués par les programmes nationaux de lutte contre le paludisme (PNLP), dont 84 % en Afrique subsaharienne.
- En 2019, 348 millions de TDR ont été vendus et 267 millions distribués par les PNL. Les ventes et les distributions de TDR en 2019 ont été inférieures aux chiffres rapportés pour 2018, de 63 millions et 24 millions respectivement, avec les plus fortes baisses enregistrées en Afrique subsaharienne.
- Entre 2010 et 2019, plus de 3,1 milliards de traitements par combinaison thérapeutique à base d'artémisinine (ACT) ont été vendus dans le monde. Sur ces ventes, près de 2,1 milliards de traitements ont été à destination du secteur public dans des pays d'endémie, alors que le reste correspond à des co-paiements publics ou privés (voire les deux), ou exclusivement au secteur des détaillants privés.
- Les données nationales rapportées par les PNL montrent que, durant la même période, 1,9 milliard de traitements par ACT ont été livrés à des prestataires de santé chargés de traiter des patients atteints de paludisme dans un établissement public.
- En 2019, quelque 190 millions de traitements par ACT ont été vendus par les fabricants au secteur public. Cette même année, les PNL ont distribué 183 millions de traitements par ACT dans ce secteur, dont 90 % en Afrique subsaharienne.
- Les données compilées à partir d'enquêtes réalisées auprès des ménages entre 2005 et 2019 dans 21 pays d'Afrique subsaharienne (ayant mené au moins deux enquêtes sur cette période, l'une entre 2005-2011 pour servir de référence et l'autre entre 2015-2019 pour les plus récentes) ont permis

d'analyser le taux de sollicitation de traitement, la couverture en diagnostic et l'utilisation des ACT chez les enfants de moins de 5 ans.

- En comparant enquêtes de référence et enquêtes plus récentes, peu de différences sont apparues concernant la prévalence de la fièvre dans les 2 semaines précédant les enquêtes (médiane de 24 % contre 21 %) et la sollicitation de traitement en cas de fièvre (médiane de 64 % contre 69 %).
- Les comparaisons de source du traitement entre enquêtes de référence et enquêtes plus récentes indiquent une médiane de 63 % contre 71% pour les soins reçus dans des établissements de santé publics, et une médiane de 39 % contre 30 % pour les soins administrés dans le secteur privé. Le recours aux agents de santé communautaires a été faible sur ces deux périodes, avec une médiane de moins de 2 %.
- Le taux de diagnostic chez les enfants de moins de 5 ans pour lesquels des soins ont été sollicités a largement progressé, d'une médiane de 15 % au départ à 38 % dans les dernières enquêtes.
- L'utilisation des ACT a également triplé, passant de 39 % à 81 % si l'on prend en compte tous les enfants fiévreux pour lesquels des soins ont été sollicités.
- Parmi les enfants fiévreux ayant subi un prélèvement sanguin au doigt ou au talon, le recours aux ACT a atteint 42 % d'après l'enquête la plus récente, suggérant que de nombreux enfants ont reçu des ACT sans diagnostic parasitologique.
- L'analyse de l'équité de la prévalence de la fièvre et de la sollicitation de soins à des niveaux infranationaux montre que, dans la plupart des pays, la prévalence de la fièvre dans les 2 semaines précédant les enquêtes était plus importante chez les enfants issus des ménages les plus démunis.
- En revanche, dans toutes les collectivités infranationales, la sollicitation de traitement était plus importante chez les enfants fiévreux issus des foyers les moins démunis et ce, même si la différence était parfois minime.

## MENACES BIOLOGIQUES

---

### Suppression des gènes *pfhrp2/3* du parasite

- La suppression des gènes *pfhrp2* et *pfhrp3* (*pfhrp2/3*) du parasite rendent ces derniers indétectables par les TDR basés sur la protéine riche en histidine 2 (HRP2).
- L'OMS a recommandé aux pays rapportant des suppressions des gènes *pfhrp2/3* ou à leurs pays voisins de mener des études de référence représentatives sur les cas suspectés de paludisme, afin de déterminer si la prévalence des suppressions *pfhrp2/3* causant des résultats de TDR négatifs avait atteint un seuil qui nécessite un changement de TDR (suppressions du gène *pfhrp2* > 5 % causant des faux résultats de TDR négatifs).
- Les alternatives aux TDR (par exemple, basées sur la détection du lactate déshydrogénase du parasite [pLDH]) sont limitées. Il n'existe à l'heure actuelle aucune combinaison de tests non-HRP2 préqualifiée par l'OMS, capable de faire la distinction entre *P. falciparum* et *P. vivax*.
- L'OMS effectue un suivi des rapports publiés sur les suppressions des gènes *pfhrp2/3* par le biais de l'outil de cartographie Carte des menaces du paludisme, et encourage une approche harmonisée de cartographie et de signalement des suppressions des gènes *pfhrp2/3* grâce à des protocoles d'enquête accessibles au public.
- Sur les 39 rapports publiés par 39 pays, 32 (82 %) ont rapporté une suppression du gène *pfhrp2* ; toutefois, les méthodes différentes de sélection des échantillons et d'analyse en laboratoire signifient que l'échelle et l'envergure d'une suppression des gènes *pfhrp2/3* significative sur le plan clinique restent à clarifier.
- Entre 2019 et septembre 2020, des enquêtes sur la suppression des gènes *pfhrp2/3* ont été rapportées dans 16 publications émanant de 15 pays. La suppression des gènes *Pfhrp2/3* a été confirmée dans 12 rapports provenant de 11 pays : Chine, Guinée équatoriale, Éthiopie, Ghana, Myanmar, Nigéria, Soudan, Ouganda, Royaume-Uni (par importation depuis divers pays d'endémie), République-Unie de Tanzanie et Zambie. Aucune suppression n'a été identifiée en France (parmi les voyageurs qui y reviennent), à Haïti, au Kenya et au Mozambique.

## Résistance des parasites aux antipaludiques

- Des mutations du gène *PfKelch13* ont été identifiées en tant que marqueurs moléculaires de résistance partielle à l'artémisinine.
- Dans la région Afrique de l'OMS, les traitements de première intention contre les infections à *P. falciparum* sont à base d'artéméter-luméfántrine (AL), d'artésunate-amodiaquine (AS-AQ) et de dihydroartémisinine-pipéraqúine (DHA-PPQ). Les taux d'efficacité contre les infections à *P. falciparum*, à savoir 98 % pour AL, 98,4 % pour AS-AQ et 99,4 % pour DHA-PPQ, n'ont jamais faibli au fil du temps. Des taux d'échec au traitement de plus de 10 % ont été observés dans quatre études sur l'AL, mais ils peuvent être considérés comme des aberrations statistiques. Il n'existe aucune preuve d'une résistance confirmée à la luméfántrine en Afrique. Pour tous les autres médicaments, les taux d'échec au traitement restent inférieurs à 10 %.
- Les traitements de première intention contre les infections à *P. falciparum* dans la région Amériques de l'OMS sont à base d'AL, d'artésunate-méfloquine (AS-MQ) et de chloroquine (CQ). L'efficacité de l'AL et de l'AS-MQ reste élevée. Une étude sur la CQ réalisée en Bolivie (État plurinational de) en 2011 a détecté un taux d'échec au traitement de 10,4 %.
- Les traitements de première intention contre les infections à *P. falciparum* dans la région Asie du Sud-Est de l'OMS sont à base d'AL, d'artésunate-sulfadoxine-pyriméthamine (AS+SP) et de DHA-PPQ. Les études relatives à l'efficacité thérapeutique de l'AL ont prouvé la très grande efficacité de ce traitement au Bhoutan, en Inde, au Myanmar, au Népal et au Timor-Leste. Des taux d'échec au traitement par AL de plus de 10 % ont été observés dans trois études, dont une en Thaïlande et deux au Bangladesh. À la suite de forts taux d'échec au traitement par AS+SP dans les provinces du nord-est en 2013, l'Inde a modifié sa politique de traitement dans ces provinces et est passée à un traitement à base d'AL. Le traitement par AS+SP reste efficace partout ailleurs dans le pays. Les résultats des études menées en Thaïlande sur l'efficacité des traitements ont conduit, en 2015, à l'adoption de la DHA-PPQ comme traitement de première intention. Des taux d'échec au traitement modérés à élevés ont été observés avec la DHA-PPQ dans l'est de la Thaïlande. De ce fait, le pays recommande actuellement un traitement à base d'artésunate-pyronaridine (AS-PY) dans cette région.
- Dans la région Méditerranée orientale de l'OMS, les traitements à base d'AL et d'AS+SP restent efficaces dans les pays qui les utilisent en tant que traitement de première intention.
- Les traitements de première intention contre *P. falciparum* dans la région Pacifique occidental de l'OMS sont à base d'AL dans tous les pays d'endémie, hormis la Chine qui utilise l'AS-AQ. Des taux d'échec au traitement par AL de 10 % ou moins ont été observés dans quatre études en République démocratique populaire lao, mais ces études ne reposaient pas sur les tailles d'échantillons recommandées. Une étude avec un nombre de patients adéquat est actuellement en cours pour examiner de plus près ces taux élevés d'échec au traitement.
- Une résistance partielle à l'artémisinine s'est développée indépendamment dans plusieurs foyers de la sous-région du Grand Mékong. L'OMS continue de surveiller la situation, qui a évolué rapidement depuis les premières détections de mutations du gène *PfKelch13* dans la sous-région du Grand Mékong. Certaines mutations ont disparu, alors que la prévalence d'autres mutations s'est accrue.
- À présent, les marqueurs affichant la prévalence la plus élevée à l'ouest de Bangkok (Thaïlande occidentale et Myanmar) sont les marqueurs F446I, M476I et R561H. Quant aux marqueurs affichant la prévalence la plus élevée à l'est de Bangkok (Thaïlande orientale, Cambodge, République démocratique populaire lao et Viet Nam), il s'agit de Y493H et P553L. Deux marqueurs, R539T et C580Y, sont également extrêmement prévalents dans ces deux zones. Le changement de politique de traitement au Cambodge, de DHA-PPQ à AS-MQ, a provoqué la réduction de la prévalence des souches portant une résistance au marqueur C580Y et à la PPQ.
- Le Rwanda a détecté une prévalence en hausse de la mutation R561H, un marqueur validé, apparu indépendamment dans la sous-région du Grand Mékong entre 2012 et 2015. La présence de cette mutation a été confirmée au Rwanda en 2018. Toutefois, il apparaît pour l'instant que l'élimination retardée associée à cette mutation n'a pas affecté l'efficacité des ACT utilisés parmi les traitements en cours de test et de déploiement au Rwanda.
- La mutation R622I semble être apparue indépendamment en Afrique et a été détectée en Érythrée, en Éthiopie, en Somalie et au Soudan, avec une fréquence en hausse dans la Corne de l'Afrique. Les ACT utilisés dans ces quatre pays restent efficaces en dépit de la présence de cette mutation. D'autres études sur l'élimination retardée du parasite sont nécessaires dans cette région.
- Au Guyana, la mutation C580Y est également apparue indépendamment entre 2010 et 2017. Cependant, des études récentes (y compris des enquêtes et des études sur l'efficacité thérapeutique) ont découvert que 100 % des échantillons sont de souche sauvage, ce qui indique que cette mutation risque de disparaître au Guyana.

## Résistance des vecteurs aux insecticides

- De 2010 à 2019, quelque 81 pays ont transmis à l'OMS des données standard de surveillance sur la résistance aux insecticides.
- Il est préoccupant de constater que 57 % des pays ayant rapporté recourir à des campagnes de PID de 2010 à 2019 n'ont pas communiqué de rapport de résistance aux insecticides pour chaque classe d'insecticides utilisés dans le courant de l'année de la mise en œuvre ou l'année précédente. De plus, 14 % n'ont pas rapporté sur la résistance aux insecticides de l'une ou l'autre classe d'insecticides utilisés. Les pays d'endémie sont vivement encouragés à assurer une surveillance adéquate de la résistance aux insecticides concernant les classes qui sont utilisées ou qui sont envisagées dans le cadre des interventions de lutte antivectorielle, ainsi qu'à donner la priorité à la surveillance de ces classes.
- Sur les 82 pays d'endémie ayant fourni des données pour la période 2010-2019, 28 ont détecté une résistance aux quatre classes d'insecticides les plus couramment utilisés chez au moins un des vecteurs du paludisme et sur un site de collecte. Par ailleurs, 73 de ces pays ont constaté une résistance à une des classes d'insecticides au moins. Seuls huit pays n'ont détecté jusqu'à présent aucune résistance à une quelconque classe d'insecticides.
- Au niveau mondial, la résistance aux pyréthoïdes, la seule classe d'insecticides actuellement utilisés dans les MII, continue de se répandre. Elle a été détectée chez au moins un des vecteurs du paludisme sur 69,9 % des sites pour lesquels des données sont disponibles. La résistance aux organochlorés a été détectée sur 63,4 % des sites. La résistance aux carbamates et aux organophosphorés a été moins prévalente, mais a été détectée, respectivement, sur 31,7 % et 24,9 % des sites disposant de données de surveillance.
- En se basant sur les données de surveillance de la résistance aux insecticides transmises à l'OMS par les États Membres, 330 zones situées dans 33 pays remplissent actuellement les critères recommandés par l'OMS pour le déploiement des moustiquaires imprégnées de butoxyde de pipéronyle (PBO).
- Même si les États Membres de l'OMS et leurs partenaires de mise en œuvre commencent à rapporter des données de surveillance sur la résistance aux néonicotinoïdes et aux pyrazoles, les États Membres sont dissuadés d'utiliser les données générées par le biais de procédures non validées pour tirer des conclusions sur l'état de résistance de leurs populations vectorielles locales face à ces classes d'insecticides. Un processus formel de l'OMS visant à établir des dosages discriminants et des procédures de tests pour ces deux classes d'insecticides est en cours de développement. Les données rapportées à l'OMS seront évaluées en tenant compte de ces dosages et procédures au fur et à mesure de leur disponibilité.
- Pour orienter la gestion de la résistance, les pays doivent développer et mettre en œuvre des plans nationaux de suivi et de gestion de la résistance aux insecticides, en se basant sur le *Cadre conceptuel d'un plan national de suivi et de gestion de la résistance aux insecticides chez les vecteurs du paludisme* élaboré par l'OMS. En 2019, le nombre de pays ayant établi un tel plan a augmenté pour atteindre 53, alors que 29 pays en étaient encore à la phase de développement.
- Les données standard sur la résistance aux insecticides rapportées à l'OMS sont intégrées à la base de données mondiales de l'OMS sur la résistance aux insecticides chez les vecteurs du paludisme, et leur accès à des fins d'exploration est possible via la Carte des menaces du paludisme. Une nouvelle version de cet outil, enrichie de fonctionnalités avancées et d'options de téléchargement, a été lancée en 2020.

## LUTTE CONTRE LE PALUDISME DURANT LA PANDÉMIE DE COVID-19

---

- En avril 2020, le coronavirus 2 associé au syndrome respiratoire aigu sévère (SARS-CoV2), virus responsable de la COVID-19, s'est propagé dans tous les pays d'endémie palustre et, à la fin de la deuxième semaine du mois de novembre 2020, près de 22 millions de cas et 600 000 décès avaient été signalés dans ces pays.
- La pandémie de COVID-19 et les restrictions imposées par la riposte ont provoqué des dysfonctionnements des services de base pour la lutte contre le paludisme.
- De plus, les premiers messages visant à réduire la transmission du coronavirus conseillaient au public de rester à la maison en cas de fièvre, ce qui a pu nuire à la sollicitation des soins en cas de survenue de fièvres, telles que celles liées au paludisme.
- En mars 2020, comme la pandémie de COVID-19 se propageait rapidement dans le monde entier, l'OMS a appelé à un effort conjoint des partenaires en vue d'atténuer l'impact négatif du coronavirus dans les pays touchés par le paludisme et de contribuer à la riposte contre la COVID-19.
- Ce travail a été mené en étroite collaboration avec le Partenariat RBM pour en finir avec le paludisme, le Fonds mondial, l'Initiative du Président américain contre le paludisme (PMI), plusieurs partenaires de mise en œuvre et de plaidoyer, ainsi que des instituts de recherche.
- Cet effort conjoint des partenaires a permis un alignement de tous et a produit des résultats, notamment :
  - la publication d'orientations techniques sur le maintien sécurisé des services de lutte contre le paludisme dans le contexte de la pandémie de COVID-19 ;
  - la publication d'une analyse par modélisation ayant pour but de quantifier l'impact potentiel des dysfonctionnements des services liés à la pandémie de COVID-19, ainsi que d'insister sur les conséquences de ces dysfonctionnements : cette analyse a souligné le risque que la mortalité due au paludisme en Afrique subsaharienne double d'ici la fin de 2020 par rapport à la référence de 2018 en cas de dysfonctionnements sévères des services de prévention et de traitement ;
  - la baisse de la pression pour orienter la production d'outils de détection du virus SARS-CoV2 au détriment de la production d'outils de diagnostic du paludisme ;
  - la suppression des goulots d'étranglement majeurs congestionnant la fabrication mondiale de médicaments antipaludiques ;
  - une limitation des dysfonctionnements dans le transport et la livraison des produits antipaludiques ;
  - la mobilisation des ressources pour les équipements de protection individuelle (EPI) et d'autres produits, afin d'aider à la mise en œuvre des campagnes de prévention, de diagnostic et de traitement ; et
  - le suivi des dysfonctionnements dans les pays pour aider à orienter la riposte.
- Cet effort collectif a donné lieu à des efforts impressionnants dans les pays, avec pour objectif de terminer les campagnes de prévention du paludisme par le biais des moustiquaires imprégnées d'insecticide longue durée (MILD), de la PID et de la CPS, et de minimiser les dysfonctionnements des services de diagnostic et de traitement.
- Tous les pays qui avaient programmé des campagnes de CPS étaient en passe de les terminer, malgré de légers retards dans certaines régions.
- Sur les 47 pays ayant planifié des campagnes de PID en 2020, 23 les ont terminées, 13 sont en passe de les terminer, et 11 sont mal partis ou risquent de ne pas les terminer.
- Plusieurs pays ont terminé leurs campagnes de distribution de MILD et un certain nombre sont encore en train de les distribuer. Pourtant, à la fin de la troisième semaine de novembre, environ 105 millions environ de MILD avaient été distribués sur les 222 millions prévues en 2020.
- Plusieurs pays ont également rapporté des niveaux de dysfonctionnements modérés. L'analyse par modélisation montre que la baisse de l'accès à un traitement antipaludique efficace, qu'elle soit de 10 %, de 15 %, de 25 % ou de 50 % en Afrique subsaharienne en 2020 pourrait respectivement entraîner 19 000, 28 000, 46 000 et 100 000 décès supplémentaires d'ici la fin de 2020 et ce, même si toutes les campagnes de prévention sont menées à bien.

# Prefacio



**Dr Tedros Adhanom Ghebreyesus**  
Director General  
Organización Mundial de la Salud (OMS)

En el *Informe mundial sobre la malaria* de este año, la OMS reflexiona sobre los hitos clave que han dado forma a la respuesta mundial contra la enfermedad durante las últimas dos décadas –un período de éxito sin precedentes en el control de la malaria en el que se evitaron 1.500 millones de casos y se salvaron 7,6 millones de vidas.

Tras la finalización del Programa Mundial de Erradicación de la Malaria en 1969, la reducción del compromiso político y de la financiación para el control de malaria provocaron el resurgimiento de ésta enfermedad en muchas partes del mundo, especialmente en África. Si bien los datos confiables son escasos, es probable que cientos de millones de personas se hayan infectado con malaria y decenas de millones hayan muerto.

A partir de la década de 1990, líderes de alto nivel en el sector de la salud y científicos trazaron un rumbo para una respuesta renovada contra la malaria. El aumento de la inversión en investigación e innovación condujo al desarrollo de nuevas herramientas para eliminar la enfermedad, como mosquiteros tratados con insecticidas, pruebas de diagnóstico rápido y medicamentos más eficaces.

La creación de nuevos mecanismos de financiación, en particular del Fondo Mundial de Lucha contra el SIDA, la Tuberculosis y la Malaria y la Iniciativa contra la Malaria del Presidente de los Estados Unidos, junto con un fuerte aumento de la financiación para malaria, permitió la distribución a gran escala de estas herramientas, contribuyendo a la reducción de la enfermedad y de las muertes en una escala que nunca antes se había visto.

El firme compromiso político en África fue clave para el éxito. A través de la histórica Declaración de Abuja del año 2000, los líderes africanos se comprometieron a reducir la mortalidad por malaria en el continente en un 50% durante un período de 10 años.

Según nuestro informe, la mortalidad mundial por malaria se redujo en un 60% durante el período 2000 a 2019. La Región de África logró reducciones impresionantes en su número anual de muertes por malaria – de 680 000 en el año 2000 a 384 000 en el 2019.

Los países del sudeste asiático lograron avances particularmente importantes, con reducciones en el número de casos y muertes del 73% y 74%, respectivamente. India contribuyó a la mayor reducción de casos en toda la región, de aproximadamente 20 millones a cerca de 6 millones.

Veintiún países han eliminado la malaria en las últimas dos décadas y, de ellos, 10 países se han certificado oficialmente por la OMS como libres de malaria. Los países del Gran Mekong continúan obteniendo importantes avances, con una asombrosa reducción del 97% en los casos de malaria por *P. falciparum* desde el año 2000, un objetivo primordial en vista de la amenaza constante que representa la resistencia a los medicamentos antimaláricos.

## Una meseta en el progreso

Los progresos obtenidos desde el comienzo del milenio han sido verdaderamente asombrosos. Sin embargo, como se ve en este informe, las ganancias se han estabilizado, tendencia observada en los últimos años.

En 2017, la OMS advirtió que la respuesta mundial contra la malaria había llegado a una “encrucijada” y que probablemente no se alcanzarían los objetivos clave de la estrategia mundial contra la malaria de la OMS. Tres años después, seguimos viendo una meseta en el progreso; según nuestro último informe, los objetivos de la estrategia para el año 2020 de reducción de la enfermedad y las muertes no se alcanzará por un 37% y 22%, respectivamente.

En 2020, COVID-19 surgió como un desafío adicional, y formidable, para las respuestas contra la malaria en todo el mundo. De acuerdo con la orientación de la OMS, muchos países han adaptado la forma en que distribuyen mosquiteros, medicamentos y realizan el diagnóstico para garantizar la seguridad de los trabajadores de salud de primera línea y las comunidades. Aplaudo de todo corazón estos esfuerzos, sin los cuales probablemente habríamos visto niveles mucho más altos en la mortalidad.

Sin embargo, según las nuevas proyecciones de la OMS, incluso alteraciones moderadas en el acceso a un tratamiento eficaz podrían provocar una considerable pérdida de vidas. El informe encuentra, por ejemplo, que una interrupción del 25% en el acceso al tratamiento antimalárico eficaz en África subsahariana podría provocar 46 000 muertes adicionales.

## Reavivar el progreso

Para revitalizar el progreso, la OMS impulsó el enfoque de “Alta carga a alto impacto” (ACAI) en 2018, junto con la Alianza para Hacer Retroceder la Malaria para Ponerle Fin. La respuesta está liderada por 11 países, incluidos 10 del África subsahariana, que representan aproximadamente el 70% de la carga mundial de malaria.

Los países de ACAI se están alejando de un enfoque de “talla única” para el control de la malaria, optando, en cambio, por implementar respuestas más particulares basadas en datos e inteligencia locales. Si bien es demasiado pronto para evaluar el impacto de este enfoque en la carga de la malaria, se han sentado bases importantes.

Un análisis reciente de Nigeria, por ejemplo, encontró que a través de una combinación optimizada de intervenciones, el país podría evitar decenas de millones de casos adicionales y miles de muertes adicionales para el año 2023, en comparación con el enfoque habitual.

Una mejor focalización de las intervenciones y los recursos contra la malaria, particularmente en países como Nigeria, donde la enfermedad golpea con más fuerza, ayudará a acelerar el ritmo del progreso hacia nuestras metas mundiales contra la malaria. También se necesita una mayor financiación a nivel nacional e internacional, junto con innovaciones en nuevas herramientas y enfoques.

Es fundamental que los esfuerzos para combatir la malaria se integren con esfuerzos más amplios para construir sistemas de salud sólidos basados en la atención primaria en salud centrada en las personas, como parte del camino de cada país hacia una cobertura universal de salud.

Es hora de que los líderes de África, y del mundo, se enfrenten una vez más al desafío de la malaria, tal como lo hicieron cuando sentaron las bases para el progreso logrado desde principios de este siglo. Mediante una acción conjunta y el compromiso de no dejar a nadie atrás, podemos lograr nuestra visión compartida de un mundo libre de malaria.

# El informe de este año de un vistazo

## TENDENCIAS EN LA CARGA DE MALARIA

---

### Casos de malaria

- A nivel mundial, hubo 229 millones de casos estimados de malaria en 2019 en 87 países donde la malaria es endémica, una disminución comparado con los 238 millones en el año 2000. En la línea de base del 2015 de la *Estrategia técnica mundial contra la malaria 2016–2030*, se estimaron 218 millones de casos de malaria.
- La proporción de casos debidos a *Plasmodium vivax* se redujo de cerca del 7% en el año 2000 a un 3% en 2019.
- La incidencia de casos de malaria (es decir, casos por 1000 habitantes en riesgo) se redujo a nivel mundial de 80 en el año 2000 a 58 en 2015 y 57 en 2019. Entre los años 2000 y 2015, la incidencia mundial de casos de malaria disminuyó en un 27%, y entre el 2015 y 2019 disminuyó en menos del 2%, lo que indica una desaceleración de la tasa de disminución desde el 2015.
- En 29 países se concentra el 95% de los casos de malaria a nivel mundial. En Nigeria (27%), la República Democrática del Congo (12%), Uganda (5%), Mozambique (4%) y Níger (3%) se presentan alrededor del 51% de todos los casos a nivel mundial.
- La Región de África de la Organización Mundial de la Salud (OMS) presentó alrededor del 94% de los casos en 2019, con un estimado de 215 millones de casos.
- Aunque hubo menos casos de malaria en el año 2000 (204 millones) que en 2019 en la Región de África de la OMS, la incidencia de casos de malaria se redujo de 363 a 225 casos por 1 000 habitantes en riesgo en este período, lo que refleja la complejidad de interpretar la transmisión cambiante de la enfermedad en una población en rápido aumento. La población que vive en la Región de África de la OMS aumentó de unos 665 millones en el año 2000 a 1.100 millones en 2019.
- La Región de Asia Sudoriental de la OMS representó alrededor del 3% de la carga de casos de malaria a nivel mundial. Los casos de malaria se redujeron en un 73%, de 23 millones en el año 2000 a aproximadamente 6.3 millones en 2019. La incidencia de casos de malaria en esta región se redujo en un 78%, de aproximadamente 18 casos por 1000 habitantes en riesgo en el año 2000 a aproximadamente cuatro casos en 2019.
- India contribuyó a las mayores reducciones absolutas en la Región de Asia Sudoriental de la OMS, de cerca de 20 millones de casos en el año 2000 a unos 5.6 millones en 2019. Sri Lanka fue certificado como libre de malaria en 2015, y Timor-Leste notificó cero casos de malaria en 2018 y 2019.
- Los casos de malaria en la Región del Mediterráneo Oriental de la OMS se redujeron en un 26%, de aproximadamente 7 millones de casos en el año 2000 a cerca de 5 millones en 2019. Aproximadamente una cuarta parte de los casos en 2019 se debieron a *P. vivax*, principalmente en Afganistán y Pakistán.
- Durante el período 2000–2019, la incidencia de casos de malaria en la Región del Mediterráneo Oriental de la OMS disminuyó de 20 a 10. Sudán es el principal contribuyente a la malaria en esta región y representa alrededor del 46% de los casos. La República Islámica del Irán no tuvo casos autóctonos de malaria en 2018 y 2019.
- La Región del Pacífico Occidental de la OMS tuvo un estimado de 1,7 millones de casos en 2019, una disminución del 43% de los 3 millones de casos en el año 2000. Durante el mismo período, la incidencia de casos de malaria se redujo de cinco a dos casos por 1 000 habitantes en riesgo. Papua Nueva Guinea representó casi el 80% de todos los casos en esta región en 2019. China no ha tenido casos autóctonos de malaria desde 2017. Malasia no tuvo casos de malaria humana en 2018 y 2019.
- En la Región de las Américas de la OMS, los casos de malaria se redujeron en un 40% (de 1,5 millones a 0,9 millones) y la incidencia de casos en un 57% (de 14 a 6) entre los años 2000 y 2019. El progreso de la región en los últimos años se ha visto afectado por el importante aumento de la malaria en Venezuela (República Bolivariana de), que registró alrededor de 35 500 casos en el año 2000, llegando a más



de 467 000 en 2019. En Brasil, Colombia y Venezuela (República Bolivariana de) se presentan más del 86% de todos los casos de esta región.

- Desde el 2015, la Región de Europa de la OMS está libre de malaria.

## Muertes por malaria

- A nivel mundial, las muertes por malaria han disminuido continuamente durante el período 2000–2019, de 736 000 en el año 2000 a 409 000 en 2019. El porcentaje del total de muertes por malaria en niños menores de 5 años fue del 84% en el año 2000 y del 67% en 2019. La estimación mundial de muertes que se realizó en el año 2015, línea de base de la Estrategia técnica mundial, fue de alrededor de 453 000.
- A nivel mundial, la tasa de incidencia de la mortalidad por malaria (es decir, muertes por cada 100 000 habitantes en riesgo) se redujo de alrededor de 25 en el año 2000 a 12 en 2015 y 10 en 2019, con una desaceleración en la tasa de disminución en los últimos años.
- Aproximadamente el 95% de las muertes por malaria de todo el mundo sucedieron en 31 países. En Nigeria (23%), la República Democrática del Congo (11%), la República Unida de Tanzania (5%), Mozambique (4%), Níger (4%) y Burkina Faso (4%) sucedieron alrededor del 51% de todas las muertes por malaria a nivel mundial en 2019.
- Las muertes por malaria en la Región de África de la OMS se redujeron en un 44%, de 680 000 en el año 2000 a 384 000 en 2019, y la tasa de incidencia de mortalidad por malaria se redujo en un 67% durante el mismo período, de 121 a 40 muertes por 100 000 habitantes en riesgo.
- En la Región de Asia Sudoriental de la OMS, las muertes por malaria se redujeron en un 74%, de cerca de 35 000 en el año 2000 a 9 000 en 2019.
- En India sucedieron aproximadamente el 86% de todas las muertes por malaria de la Región de Asia Sudoriental de la OMS.
- En la Región del Mediterráneo Oriental de la OMS, las muertes por malaria se redujeron en un 16%, de alrededor de 12 000 en el año 2000 a 10 100 en 2019, y la tasa de incidencia de mortalidad por malaria se redujo en un 50%, de cuatro a dos muertes por 100 000 habitantes en riesgo.
- En la Región del Pacífico Occidental de la OMS, las muertes por malaria se redujeron en un 52%, de aproximadamente 6 600 casos en el año 2000 a 3 200 en 2019, y la tasa de incidencia de mortalidad se redujo en un 60%, pasando de una a 0,4 muertes por malaria por 100 000 habitantes en riesgo. En Papua Nueva Guinea sucedieron más del 85% de las muertes por malaria en 2019.
- En la Región de las Américas de la OMS, las muertes por malaria se redujeron en un 39% (de 909 a 551) y la tasa de incidencia de mortalidad en un 50% (de 0,8 a 0,4) entre el 2000 y 2019. Más del 70% de las muertes por malaria en 2019 en esta región sucedieron en Venezuela (República Bolivariana de).

## Casos de malaria y muertes evitadas

- A nivel mundial, se estima que se han evitado 1.500 millones de casos de malaria y 7,6 millones de muertes por malaria en el período 2000–2019.
- La mayoría de los casos (82%) y muertes (94%) evitados fueron en la Región de África de la OMS, seguida de la Región de Asia Sudoriental de la OMS (10% de los casos y 3% de las muertes).

## Carga de la malaria en el embarazo

- En 2019, en 33 países con transmisión moderada y alta en la Región de África de la OMS, hubo aproximadamente 33 millones de mujeres embarazadas, de las cuales el 35% (12 millones) estuvieron expuestas a la infección por malaria durante el embarazo.
- Por subregión de la OMS, África Central tuvo la mayor prevalencia de exposición a la malaria durante el embarazo (40%), seguida de cerca por África Occidental (39%), mientras que la prevalencia fue del 24% en África Oriental y en África del Sur.
- Se estima que la infección por malaria durante el embarazo en estos 33 países resultó en 822 000 niños con bajo peso al nacer.
- Si el 80% de las mujeres embarazadas que informaron haber utilizado los servicios de atención prenatal alguna vez, hubieran recibido una dosis de tratamiento preventivo intermitente durante el embarazo, se habrían evitado adicionalmente 56 000 nacimientos de bajo peso en estos 33 países.

## ELIMINACIÓN DE LA MALARIA Y PREVENCIÓN DE SU RESTABLECIMIENTO

---

- A nivel mundial, el número de países que eran endémicos de malaria en el 2000 y que notificaron menos de 10 000 casos de malaria aumentó de 26 en el año 2000 a 46 en 2019.
- En el mismo período, el número de países con menos de 100 casos autóctonos aumentó de seis a 27.
- En el período 2010-2019, el total de casos de malaria en los 21 países de la E-2020 se redujo en un 79%.
- Hubo más casos en 2019 que en 2018 en Comoras, Costa Rica, Ecuador y Surinam, los cuales informaron de 1986, 25, 150 y 66 casos adicionales, respectivamente, en 2019.
- Irán (República Islámica del), Malasia y Timor-Leste notificaron cero casos autóctonos de malaria en 2018 y 2019. En 2019, Belice y Cabo Verde notificaron cero casos autóctonos de malaria por primera vez desde el año 2000.
- China y El Salvador no tuvieron casos autóctonos de malaria por tercer año consecutivo y han presentado una solicitud formal de certificación.
- Entre 2000 y 2019, en los seis países de la subregión del Gran Mekong (SGM) - Camboya, China (provincia de Yunnan), República Democrática Popular Lao, Myanmar, Tailandia y Vietnam - los casos de malaria por *P. falciparum* disminuyeron en un 97%, mientras que todos los casos de malaria se redujeron en un 90%. De los 239 000 casos de malaria notificados en 2019, 65 000 fueron casos de *P. falciparum*.
- La tasa de disminución ha sido más rápida desde 2012, cuando se lanzó el programa de Eliminación de la Malaria del Mekong. Durante este período, los casos de malaria se redujeron seis veces, mientras que los casos de *P. falciparum* se redujeron en un factor de casi 14.
- En general, Camboya (58%) y Myanmar (31%) representaron la mayoría de los casos de malaria en la SGM.
- Esta disminución acelerada de *P. falciparum* es especialmente crítica debido al aumento de la resistencia a los medicamentos; en la SGM, los parásitos *P. falciparum* han desarrollado una resistencia parcial a la artemisinina, el compuesto principal de los mejores fármacos antimaláricos disponibles.
- Entre los años 2000 y 2019, no se ha restablecido la transmisión de la malaria en ningún país certificado como libre de malaria.

## ENFOQUE “DE ALTA CARGA A ALTO IMPACTO”

---

- Desde noviembre de 2018, el enfoque de alta carga a alto impacto (ACAI) se ha lanzado en 10 de los 11 países (aún no se ha lanzado en Mali debido a las alteraciones por la pandemia de COVID-19). Sin embargo, los 11 países han implementado actividades relacionadas con ACAI en los cuatro elementos de la respuesta.
- En cada país donde se ha iniciado el enfoque ACAI, ha habido un alto nivel de compromiso y apoyo político. La iniciativa de Acción en Masa Contra la Malaria en Uganda se presenta como un ejemplo de un proceso liderado por un país de participación política en todos los niveles y de movilización multisectorial y comunitaria.
- Se ha completado el análisis para la adaptación sub-nacional de las intervenciones en todos los países excepto en Malí, donde este trabajo está en progreso. El ejemplo de Nigeria se presenta en el informe.
- Todos los países se han comprometido a realizar un ejercicio integral de micro-estratificación urbana para orientar mejor las intervenciones y mejorar la eficiencia dada la creciente tasa de urbanización.
- El Programa Mundial de Malaria (PMM) de la OMS actualizó su informe técnico para ayudar a los países a priorizar mejor los recursos, al tiempo que se adhieren a las recomendaciones basadas en evidencia que se han desarrollado a través de los rigurosos procesos estándar de la OMS.
- Debido a que la respuesta de ACAI se lanzó en noviembre de 2018, cuando los países estaban llegando al final de sus ciclos de financiamiento, es demasiado pronto para determinar el impacto de la respuesta. El número de casos de malaria en los 11 países de ACAI en 2019 fue similar al de 2018 (156 millones frente a 155 millones).

## PROGRESO HACIA LOS OBJETIVOS DE LA ESTRATEGIA TÉCNICA MUNDIAL (ETM) DE 2020

---

- La ETM apunta a una reducción en la incidencia de casos de malaria y la tasa de mortalidad de al menos un 40% para 2020, un 75% para 2025 y un 90% para 2030, comparados con la línea de base de 2015.
- Las tendencias de 2000-2019 en casos y muertes por malaria se utilizaron para hacer proyecciones anuales para 2020 a 2030, con el fin de hacer un seguimiento del progreso hacia las metas y los hitos de la ETM.
- Las proyecciones presentadas en este informe no tienen en cuenta las posibles alteraciones debidas a la pandemia COVID-19, la cual, a pesar de los encomiables esfuerzos mundiales y nacionales para mantener los servicios esenciales contra la malaria, es probable que provoque una morbilidad y mortalidad por malaria más altas de lo esperado.
- A pesar de los considerables avances realizados desde el año 2000, los objetivos para la morbilidad y mortalidad de la ETM 2020 no se alcanzarán a nivel mundial.
- La incidencia de casos de malaria de 56 casos por 1 000 habitantes en riesgo en 2020 en lugar de los 35 casos por 1 000 esperados si el mundo estuviera encaminado hacia el objetivo de morbilidad de la ETM 2020 significa que, a nivel mundial, la trayectoria actual se ha desviado en un 37% de lo esperado.
- Aunque el progreso relativo en la tasa de mortalidad es mayor que en la incidencia de casos, las muertes por malaria proyectadas a nivel mundial por cada 100 000 habitantes en riesgo en 2020 fue de 9,8, comparado con 11.9 en 2015. Esto implica que el mundo está un 22% fuera de la trayectoria establecida por el ETM para el 2020.
- De los 92 países que eran endémicos de malaria a nivel mundial en 2015, se estimó que 31 (34%) estaban en camino de alcanzar el objetivo de morbilidad de la ETM para el año 2020, habiendo logrado una reducción del 40% o más en la incidencia de casos o informado de cero casos de malaria.
- Veintiún países (23%) han progresado en la reducción de la incidencia de casos de malaria, pero no están en camino de alcanzar el objetivo de la ETM.
- Se estima que 31 países (34%) tienen una mayor incidencia, y se estima que 15 países (16%) tienen un aumento del 40% o más en la incidencia de casos de malaria en 2020 en comparación con 2015.
- Se estimó que la incidencia de casos de malaria en nueve países (10%) en 2020 se encuentra en niveles similares a los de 2015.
- Treinta y nueve países (42%) que eran endémicos de malaria en 2015 estaban en camino de alcanzar el objetivo de mortalidad de la ETM para 2020, y 28 de ellos notificaron cero casos de malaria.
- Se estimó que 34 países (37%) habían logrado reducciones en las tasas de incidencia de la mortalidad por malaria, pero el progreso estuvo por debajo de la meta del 40%.
- Las tasas de incidencia de mortalidad por malaria se mantuvieron al mismo nivel en 2020 que en 2015 en siete países (8%), mientras que se estimaron aumentos en otros 12 países (13%), seis de los cuales tuvieron aumentos del 40% o más.
- Todos los países de la Región de Asia Sudoriental de la OMS están en camino de alcanzar los objetivos de la ETM 2020, tanto en morbilidad como en mortalidad.

## INVERSIONES EN PROGRAMAS E INVESTIGACIÓN SOBRE MALARIA

---

- La ETM establece estimaciones de la financiación necesaria para alcanzar los objetivos para los años 2020, 2025 y 2030. Los recursos anuales totales necesarios se estimaron en 4.100 millones de dólares estadounidenses en 2016, y ascendieron a 6 800 millones de dólares estadounidenses en 2020. Se estima que se necesitarán otros 700 millones de dólares anuales para investigación y desarrollo (I & D) a nivel mundial en malaria.
- El financiamiento total para el control y la eliminación de la malaria en 2019 se estimó en \$ 3.000 millones, en comparación con \$ 2 700 millones en 2018 y \$ 3 200 millones en 2017. La cantidad

invertida en 2019 no alcanza los \$ 5 600 millones de dólares estimados como necesarios a nivel mundial para mantenerse encaminado hacia los hitos de la ETM.

- La brecha de financiamiento entre el monto invertido y los recursos necesarios ha seguido ampliándose drásticamente en los últimos años, pasando de 1 300 millones de dólares en 2017 a 2 300 millones de dólares en 2018 y a 2.600 millones de dólares en 2019.
- Durante el período 2010–2019, fuentes internacionales proporcionaron el 70% de la financiación total para el control y la eliminación de la malaria, encabezadas por los Estados Unidos de América (EE UU.), el Reino Unido de Gran Bretaña e Irlanda del Norte (Reino Unido) y Francia.
- De los \$ 3 000 millones de dólares invertidos en 2019, \$ 2 100 millones provinieron de financiadores internacionales. Las mayores contribuciones en 2019 fueron del gobierno de los EE. UU., quien proporcionó un total de \$ 1,1 mil millones de dólares a través de fondos bilaterales planificados y contribuciones a agencias de financiamiento multilaterales.
- A esto le siguieron desembolsos bilaterales y multilaterales del Reino Unido por 200 millones de dólares, contribuciones de más de 100 millones de dólares de cada uno de los países de Francia, Alemania y Japón (por un total de 400 millones de dólares estadounidenses) y un total combinado de 400 millones de otros países que son miembros del Comité de Asistencia para el Desarrollo y de contribuyentes del sector privado.
- Los gobiernos de los países donde la malaria es endémica continuaron aportando alrededor del 30% del financiamiento total, con inversiones cercanas a los \$ 900 millones de dólares en 2019. De esta cantidad, se estima que \$ 200 millones se gastaron en el manejo de los casos de malaria en el sector público y \$ 700 millones en otras actividades de control de la malaria.
- De los \$ 3 000 millones de dólares invertidos en 2019, casi \$ 1 200 millones (39%) se canalizaron a través del Fondo Mundial de Lucha contra el SIDA, la Tuberculosis y la Malaria (Fondo Mundial). En comparación con 2018, los desembolsos del Fondo Mundial a los países donde la malaria es endémica aumentaron en alrededor de 200 millones de dólares en 2019.
- De los \$ 3 000 millones invertidos en 2019, alrededor del 73% se destinó a la Región de África de la OMS, el 9% a la Región de Asia Sudoriental de la OMS, el 5% a la Región de las Américas de la OMS y a la Región del Pacífico Occidental de la OMS, y 4% a la Región del Mediterráneo Oriental de la OMS.
- Entre 2007 y 2018, se invirtieron casi \$ 7 300 millones de dólares en investigación básica y desarrollo de productos para la malaria.
- El panorama de la financiación de la I & D contra la malaria ha sido liderado por la inversión en medicamentos (\$ 2,6 mil millones, 36% de la financiación contra la malaria entre 2007 y 2018), seguida de proporciones relativamente similares para la investigación básica (\$ 1,9 mil millones, 26%) y la I & D sobre vacunas (\$ 1.8 mil millones, 25%). Las inversiones en productos de control de vectores y diagnóstico fueron notablemente menores, alcanzando un total general de \$ 453 millones (6,2%) y \$ 185 millones (2,5%), respectivamente.
- Entre 2007 y 2018, el sector público ocupó un papel de liderazgo en la financiación de I & D contra la malaria, pasando de 246 millones de dólares estadounidenses en 2007 a un máximo de 365 millones de dólares estadounidenses en 2017. Dentro del sector público y entre todos los financiadores de I & D contra la malaria, los Institutos Nacionales de Salud de Estados Unidos fue el mayor contribuyente, y centró poco más de la mitad de su inversión de 1 900 millones de dólares en investigación básica (1.020 millones de dólares, el 54% de su inversión total en malaria entre 2007 y 2018).
- La Fundación Bill y Melinda Gates ha sido otro actor fundamental, invirtiendo 1.800 millones de dólares (el 25% de todos los fondos para I & D contra la malaria) entre 2007 y 2018, y apoyando el desarrollo clínico de innovaciones clave como la vacuna RTS, S.

## **DISTRIBUCIÓN Y COBERTURA DE LA PREVENCIÓN DE LA MALARIA**

---

- Los datos de entrega de los fabricantes muestran que se suministraron en todo el mundo casi 2.200 millones de mosquiteros tratados con insecticidas (MTI) entre 2004 y 2019, de los cuales 1.900 millones (86%) se suministraron al África subsahariana.
- Los fabricantes entregaron alrededor de 253 millones de mosquiteros tratados con insecticidas a países endémicos de malaria en 2019, un aumento de 56 millones de mosquiteros tratados con insecticidas

en comparación con 2018. Aproximadamente el 84% de estos mosquiteros tratados con insecticidas se entregaron a países del África subsahariana.

- Para el año 2019, el 68% de los hogares en África subsahariana tenían al menos un MTI, en comparación con aproximadamente un 5% en el año 2000. El porcentaje de hogares que poseen al menos un MTI por cada dos personas aumentó del 1% en el 2000 al 36% en el 2019. En el mismo período, el porcentaje de población con acceso a un MTI dentro de su hogar aumentó de 3% a 52%.
- El porcentaje de la población que duerme bajo un MTI también aumentó considerablemente entre el año 2000 y el 2019, para toda la población (del 2% al 46%), para los niños menores de 5 años (del 3% al 52%) y para las mujeres embarazadas (del 3% al 52%).
- Los datos de encuestas de hogares más recientes de las encuestas demográficas y de salud y las encuestas de indicadores de malaria de 24 países de África subsahariana de 2015 a 2019 se utilizaron para analizar la equidad socioeconómica en el uso de MTI. En la mayoría de los países de África occidental, el uso de mosquiteros tratados con insecticidas fue generalmente a favor de los pobres o estuvo cerca de una equitatividad perfecta. Por el contrario, el uso de MTI fue mayor en los hogares más ricos de muchas partes de África central y oriental.
- A nivel mundial, el porcentaje de población en riesgo protegida con rociado residual intradomiciliario (RRI) en países endémicos para malaria disminuyó del 5% en 2010 al 2% en 2019. El porcentaje de la población protegida con RRI disminuyó en todas las regiones de la OMS.
- La cantidad de personas protegidas en todo el mundo se redujo de 180 millones en 2010 a 115 millones en 2015, y disminuyó a 97 millones en 2019.
- El número de niños a los que se llegó con al menos una dosis de quimio-prevencción estacional de la malaria (QPE) aumentó constantemente, de aproximadamente 0,2 millones en 2012 a aproximadamente 21,5 millones en 2019.
- En los 13 países que implementaron QPE, la intervención fue dirigida a alrededor de 21,7 millones de niños en 2019. En promedio, 21,5 millones de niños recibieron tratamiento.
- Utilizando datos de 33 países africanos, se calculó el porcentaje de uso del Tratamiento Preventivo Intermitente de la malaria durante el Embarazo (TPI) por dosis. En 2019, el 80% de las mujeres embarazadas utilizaron los servicios de atención prenatal al menos una vez durante el embarazo. Aproximadamente el 62% de las mujeres embarazadas recibió una dosis de TPI y el 49% recibió 2 dosis TPI. Hubo un ligero aumento en la cobertura de 3 dosis de TPI, del 31% en 2018 al 34% en 2019.

## **DISTRIBUCIÓN Y COBERTURA DEL DIAGNÓSTICO Y TRATAMIENTO DE LA MALARIA**

- A nivel mundial, los fabricantes vendieron 2.700 millones de pruebas de diagnóstico rápido (PDR) para la malaria entre 2010 y 2019, y casi el 80% de estas ventas se realizaron a países del África subsahariana. En el mismo período, los programas nacionales de malaria (PNM) distribuyeron 1.900 millones de PDR, el 84% en África subsahariana.
- En 2019, los fabricantes vendieron 348 millones de PDR y los PNM distribuyeron 267 millones. La venta y distribución de PDR en 2019 fueron inferiores a las informadas en 2018, en 63 millones y 24 millones, respectivamente, y la mayoría de las disminuciones se produjeron en África subsahariana.
- Los fabricantes vendieron a nivel mundial más de 3.100 millones de tratamientos de terapia combinada con derivados de la artemisinina (TCA) en 2010-2019. Aproximadamente 2,1 mil millones de estas ventas fueron al sector público en países donde la malaria es endémica, y el resto se vendió a través de copagos del sector público o privado (o ambos), o exclusivamente a través del sector minorista privado.
- Los datos nacionales informados por los PNM muestran que, en el mismo período, se entregaron 1.900 millones de TCA a los proveedores de servicios de salud para tratar a los pacientes con malaria en el sector de la salud pública.
- En 2019, los fabricantes vendieron unos 190 millones de TCA para su uso en el sector de la salud pública; En ese mismo año, los PNM distribuyeron 183 millones de TCA a este sector, de los cuales el 90% estaban en África subsahariana.

- Los datos agregados de las encuestas de hogares realizadas en África subsahariana entre 2005 y 2019 en 21 países con al menos dos encuestas en este período (línea de base 2005-2011 y más reciente 2015-2019) se utilizaron para analizar la cobertura de la búsqueda de tratamiento, el diagnóstico y uso de TCA en niños menores de 5 años.
- Comparando las encuestas de línea de base con las más recientes, hubo pocos cambios en la prevalencia de fiebre dentro de las 2 semanas anteriores a las encuestas (mediana 24% versus 21%) y búsqueda de tratamiento para la fiebre (mediana 64% versus 69%).
- Las comparaciones de la fuente de tratamiento entre la línea de base y las encuestas más recientes muestran que una mediana del 63% frente al 71% recibió atención en instalaciones de salud pública y una mediana del 39% frente al 30% recibió atención del sector privado. El uso de trabajadores de salud comunitarios fue bajo en ambos períodos, con una mediana de menos del 2%.
- La tasa de diagnóstico entre los niños menores de 5 años para quienes se buscó atención aumentó considerablemente, de una mediana del 15% al inicio, al 38% en las últimas encuestas de hogares.
- El uso de TCA también se multiplicó por más de tres, del 39% al inicio, al 81% en las últimas encuestas, cuando se consideraron todos los niños con fiebre para quienes se buscó atención.
- Entre los que recibieron un pinchazo en el dedo o el talón, el uso de TCA fue del 42% en la encuesta más reciente, lo que sugiere que muchos niños recibieron TCA sin diagnóstico parasitológico.
- Analizando la equidad en la prevalencia de la fiebre y la búsqueda de tratamiento a nivel sub-nacional, se muestra que en la mayoría de los países, los niños de los hogares más pobres tenían una mayor prevalencia de fiebre en las 2 semanas anteriores a las encuestas de hogares.
- En contraste, la búsqueda de tratamiento fue mayor en los niños febriles de hogares más ricos en todas las unidades sub-nacionales, aunque en algunas unidades esa diferencia fue pequeña.

## AMENAZAS BIOLÓGICAS

### Deleciones en los genes *pfhrp2* / *3* de los parásitos

- Las deleciones en los genes *pfhrp2* y *pfhrp3* (*pfhrp2* / *3*) del parásito hacen que los parásitos sean indetectables por las PDR basadas en la proteína 2 rica en histidina (HRP2).
- La OMS ha recomendado que los países con informes de deleciones de *pfhrp2* / *3* o los países vecinos deben realizar encuestas de línea de base representativas en los casos sospechosos de malaria para determinar si la prevalencia de deleciones de *pfhrp2* / *3* que causan resultados de falsos negativos de la PDR ha alcanzado un umbral para el cambio de PDR (> 5 % de deleciones de *pfhrp2* que causan resultados de falsos negativos en PDR).
- Las opciones alternativas de PDR (por ejemplo, basadas en la detección de la lactato deshidrogenasa [pLDH] del parásito) son limitadas; en particular, actualmente no existen pruebas precalificadas de las OMS que no sean pruebas de combinación de HRP2 que puedan detectar y distinguir entre *P. falciparum* y *P. vivax*.
- La OMS está rastreando reportes publicados de deleciones de *pfhrp2* / *3* utilizando la herramienta de mapeo *Mapa de los Desafíos de la Malaria* (Malaria Threats Map) y está fomentando un enfoque armonizado para mapear y notificar las deleciones de *pfhrp2* / *3* a través de protocolos de encuestas disponibles públicamente.
- Entre los 39 informes publicados por 39 países, 32 (82%) reportaron deleciones de *pfhrp2*; sin embargo, la variabilidad en los métodos de selección de muestras y análisis de laboratorio significan que la escala y el alcance de la significancia clínica de las deleciones de *pfhrp2* / *3* aún no está claro.
- Entre 2019 y septiembre de 2020, se informaron investigaciones de deleciones de *pfhrp2* / *3* en 16 publicaciones de 15 países. Se confirmaron deleciones de *Pfhrp2* / *3* en 12 informes de 11 países: China, Guinea Ecuatorial, Etiopía, Ghana, Myanmar, Nigeria, Sudán, Uganda, Reino Unido (importados de varios países endémicos de malaria), República Unida de Tanzania y Zambia. No se identificaron deleciones en Francia (entre los viajeros que regresan), Haití, Kenia y Mozambique.

## Resistencia de los parásitos a los medicamentos antimaláricos

- Se han identificado mutaciones de Pfk13 como marcadores moleculares de resistencia parcial a la artemisinina.
- En la Región de África de la OMS, los tratamientos de primera línea para *P. falciparum* incluyen arteméter-lumefantrina (AL), artesunato-amodiaquina (AS-AQ) y dihidroartemisinina-piperaquina (DHA-PPQ). Las tasas de eficacia promedio general para *P. falciparum* (98,0% para AL, 98,4% para AS-AQ y 99,4% para DHA-PPQ) se mantuvieron constantes a lo largo del tiempo. Se observaron tasas de fallas del tratamiento de más del 10% en cuatro estudios de AL, pero pueden considerarse valores estadísticos atípicos. No hay evidencia de resistencia confirmada a la lumefantrina en África. Para todos los demás medicamentos, las tasas de fallas terapéuticas permanecen por debajo del 10%.
- Los tratamientos de primera línea para *P. falciparum* en la Región de las Américas de la OMS incluyen AL, artesunato-mefloquina (AS-MQ) y cloroquina (CQ). La eficacia de AL y AS-MQ sigue siendo alta. Un estudio de CQ en Bolivia (Estado Plurinacional de) en 2011 detectó una tasa de falla terapéutica del 10,4%.
- Los tratamientos de primera línea para *P. falciparum* en la Región de Asia Sudoriental de la OMS incluyen AL, artesunato-sulfadoxina-pirimetamina (AS + SP) y DHA-PPQ. Los estudios de eficacia terapéutica (EET) de AL demostraron una alta eficacia del tratamiento en Bután, India, Myanmar, Nepal y Timor-Leste. Las tasas de falla terapéutica para AL superaron el 10% en tres estudios, uno en Tailandia y dos en Bangladesh. Tras las altas tasas de falla terapéutica a AS + SP en las provincias del noreste, en 2013, India cambió su política de tratamiento en esas provincias a AL; AS + SP sigue siendo eficaz en otras partes del país. Los hallazgos de EET en Tailandia llevaron a la adopción de DHA-PPQ como tratamiento de primera línea en 2015. En Tailandia, se observaron tasas moderadas a altas de falla terapéutica con DHA-PPQ en la parte oriental del país; por lo tanto, Tailandia recomienda actualmente el tratamiento con artesunato-pironaridina (AS-PY) en esta área.
- AL y AS + SP siguen siendo eficaces en los países que los utilizan como tratamiento de primera línea en la Región del Mediterráneo Oriental de la OMS.
- Los tratamientos de primera línea para *P. falciparum* en la Región del Pacífico Occidental de la OMS son AL en todos los países donde la malaria es endémica, excepto China, donde se usa AS-AQ. Las tasas de falla terapéutica a AL fueron del 10% o menos en cuatro estudios en la República Democrática Popular Lao, pero esos estudios no tenían los tamaños de muestra recomendados. Actualmente se está realizando un estudio con un número adecuado de pacientes para investigar más a fondo estas altas tasas de falla terapéutica.
- La resistencia parcial a la artemisinina surgió de forma independiente en varios focos de la sub-región del gran Mekong (SGM). La OMS continúa monitoreando la situación, que ha evolucionado rápidamente desde las primeras detecciones de mutaciones de *Pfk13* en la SGM. Algunas mutaciones han desaparecido, mientras que la prevalencia de otras ha aumentado.
- Actualmente, los marcadores más frecuentes al oeste de Bangkok (oeste de Tailandia y Myanmar) son F446I, M476I y R561H. Los marcadores más frecuentes al este de Bangkok (este de Tailandia, Camboya, República Democrática Popular Lao y Viet Nam) son Y493H y P553L. Dos marcadores, R539T y C580Y, también son muy prevalentes en ambas áreas. El cambio en la política de tratamiento en Camboya de DHA-PPQ a AS-MQ resultó en una reducción en la prevalencia de cepas portadoras de resistencia tanto a C580Y como a PPQ.
- Ruanda ha detectado una prevalencia creciente de la mutación R561H, un marcador validado que surgió de forma independiente en la SGM entre 2012 y 2015. La presencia de esta mutación se confirmó en Ruanda en 2018; sin embargo, hasta ahora parece que el retraso en curar la parasitemia asociado con esta mutación no ha afectado la eficacia de la terapia combinada con derivados de la artemisinina (TCA) que se encuentran actualmente entre los evaluados y utilizados en Ruanda.
- La mutación R622I parece estar apareciendo de forma independiente en África, habiéndose encontrado en Eritrea, Etiopía, Somalia y Sudán, y con una frecuencia cada vez mayor en el Cuerno de África. La TCA utilizada en estos cuatro países sigue siendo eficaz, a pesar de la presencia de la mutación. Se necesita una mayor investigación sobre las demoras en curar la parasitemia en esta región.
- En Guyana, la mutación C580Y también surgió de forma independiente entre 2010 y 2017. Sin embargo, en estudios recientes (incluidas encuestas y EET), se encontró que el 100% de las muestras contenían en el gen de tipo silvestre, lo que indica que la mutación puede estar desapareciendo en Guyana.

## Resistencia de los vectores a los insecticidas

- De 2010 a 2019, unos 81 países reportaron datos a la OMS sobre la vigilancia regular de la resistencia a los insecticidas.
- De manera preocupante, entre 2010 y 2019, el 57% de los países que informaron usar el RRI no informaron el estado de resistencia a los insecticidas para cada clase de insecticida utilizada en el año de implementación o el anterior, y el 14% no informó sobre el estado de resistencia para cualquier clase de insecticida utilizado. Se exhorta encarecidamente a los países donde la malaria es endémica a garantizar un seguimiento adecuado de la resistencia a los insecticidas de las clases que están en uso o que se están considerando para su uso en las intervenciones de control del vector de la malaria, y a priorizar el seguimiento de estas clases.
- De los 82 países con malaria endémica que proporcionaron datos para 2010-2019, 28 han detectado resistencia a las cuatro clases de insecticidas más comúnmente utilizadas en al menos un vector de la malaria y un sitio de recolección, y 73 han detectado resistencia a al menos una clase de insecticida. Hasta el momento, solo ocho países no han detectado resistencia a ninguna clase de insecticida.
- A nivel mundial, la resistencia a los piretroides – la única clase de insecticida que se usa actualmente en los MTI– continúa siendo generalizada. Se detectó en al menos un vector de la malaria en el 69,9% de los sitios para los que se disponía de datos. Se informó de resistencia a los organoclorados en el 63,4% de los sitios. La resistencia a carbamatos y organofosforados fue menos prevalente, detectándose en 31,7% y 24,9% de los sitios que reportaron datos de monitoreo, respectivamente.
- Según los datos de seguimiento de la resistencia a los insecticidas comunicados a la OMS por los Estados Miembros, un total de 330 áreas en 33 países cumplen actualmente los criterios recomendados por la OMS para implementar mosquiteros con piretroide y butóxido de piperonilo.
- Aunque los Estados Miembros de la OMS y sus socios han comenzado a notificar datos de la vigilancia de la resistencia a los insecticidas para neonicotinoides y pirroles, se desaconseja a los Estados Miembros que utilicen datos generados mediante procedimientos no validados para llegar a conclusiones sobre el estado de resistencia de sus poblaciones de vectores locales a estas clases de insecticidas. Está en curso un proceso formal de la OMS para establecer dosis discriminantes y procedimientos de prueba para estas dos clases de insecticidas. Los datos notificados a la OMS se evaluarán de acuerdo con estas dosis y procedimientos a medida que estén disponibles.
- Para guiar el manejo de la resistencia, los países deben desarrollar e implementar un plan nacional para el monitoreo y manejo de la resistencia a los insecticidas, basándose en el *Marco de la OMS para un plan nacional para el monitoreo y manejo de la resistencia a los insecticidas en los vectores de la malaria*. En 2019, el número de países que habían completado esos planes aumentó a 53, y 29 países estaban en proceso de desarrollarlos.
- Los datos estándar de resistencia a los insecticidas notificados a la OMS se incluyen en la base de datos mundial de la OMS sobre la resistencia a los insecticidas en los vectores de la malaria y están disponibles para su exploración a través del Mapa de Amenazas de la Malaria (World Malaria Threats Map). En 2020 se lanzó una nueva versión de esta herramienta con funciones mejoradas y opciones de descarga de datos.



## RESPUESTA A LA MALARIA DURANTE LA PANDEMIA DE COVID-19

---

- Para abril de 2020, el síndrome respiratorio agudo severo coronavirus 2 (SARS-CoV2), que causa la COVID-19, se había extendido a todos los países donde la malaria es endémica, y al final de la segunda semana de noviembre de 2020, alrededor de 22 millones de casos y 600 000 muertes se habían informado en éstos países.
- La pandemia de COVID-19 y las restricciones relacionadas con la respuesta han provocado alteraciones en los servicios esenciales contra la malaria.
- Además, los primeros mensajes dirigidos a reducir la transmisión del coronavirus aconsejaron al público que se quedara en casa si tenían fiebre, lo que podría alterar la búsqueda de tratamiento para enfermedades febriles como la malaria.
- En marzo de 2020, a medida que la pandemia de COVID-19 se propagaba rápidamente por todo el mundo, la OMS convocó un esfuerzo entre socios para mitigar el impacto negativo del coronavirus en los países afectados por la malaria y contribuir a la respuesta a la COVID-19.
- El trabajo se llevó a cabo en estrecha colaboración con la iniciativa Hacer Retroceder la Malaria (Roll Back Malaria) para ponerle fin a la malaria, el Fondo Mundial, la Iniciativa del Presidente de los Estados Unidos contra la Malaria (PMI), varios socios de implementación y promoción e instituciones de investigación.
- El esfuerzo entre socios condujo a una estrecha colaboración que dio lugar a varios resultados:
  - publicación de orientación técnica sobre cómo mantener de manera segura los servicios de control de la malaria en el contexto de la pandemia de COVID-19;
  - publicación de un análisis de modelación para cuantificar el impacto potencial de las alteraciones del servicio debido a la pandemia de COVID-19, para reforzar las consecuencias de las alteraciones del servicio. El análisis sugirió que era probable que la mortalidad por malaria en África subsahariana se duplicara para fines de 2020, en relación con la línea de base de 2018, si se produjera una interrupción extrema en la prevención y el tratamiento;
  - mitigar la presión para cambiar la producción de pruebas de diagnóstico de malaria por pruebas para la detección del virus SARS-CoV2;
  - éxito en la resolución de los principales obstáculos mundiales en la fabricación de medicamentos contra la malaria;
  - mitigar las interrupciones en el envío y entrega de productos para malaria;
  - movilización de recursos para equipos de protección personal (EPP) y otros productos básicos para ayudar con la implementación de campañas de prevención, diagnóstico y tratamiento; y
  - seguimiento de las alteraciones en los países para ayudar a orientar la respuesta.
- El esfuerzo colectivo ha llevado a los países a realizar esfuerzos impresionantes para completar campañas de prevención de la malaria que involucran mosquiteros insecticidas de larga duración (MILD), RRI y quimio-prevención estacional de la malaria (QPE), y para minimizar las interrupciones en el diagnóstico y el tratamiento.
- Todos los países que habían planificado campañas de QPE estaban en camino de completarlas a pesar de retrasos moderados en algunas áreas.
- De los 47 países que tenían campañas del RRI planificadas en 2020, 23 las habían completado, 13 estaban en camino de completarlas y 11 estaban desencaminados o en riesgo de no completarlas.
- Varios países han completado sus campañas de MILD y muchos están en proceso de distribuir MILDs. Sin embargo, a la tercera semana de noviembre, de los 222 millones de MILD planificados para su distribución en 2020, solo se habían distribuido alrededor de 105 millones.
- Muchos países también han informado de niveles moderados de alteraciones, y el análisis de modelos muestra que reducciones en el acceso al tratamiento antimalárico efectivo del 10%, 15%, 25% y 50% en África subsahariana en 2020 podrían conducir a 19 000, 28 000, 46 000 y 100 000 muertes más por malaria, respectivamente, para fines de 2020, incluso si se completan todas las campañas de prevención.





# 1 Introduction

The year 2020 is a milestone for several important health and development goals, including for efforts to reduce the burden of malaria overall and eliminate the disease where possible. It is 20 years since the Abuja Declaration (1) and the launch of the Millennium Development Goals (MDGs) (2); and 5 years since the global agreement on the Sustainable Development Goals (SDGs) framework (3) and the launch of the World Health Organization (WHO) *Global technical strategy for malaria 2016–2030* (GTS) (4) and the RBM Partnership to End Malaria *Action and investment to defeat malaria 2016–2030* (AIM) (5). The WHO *World malaria report 2020* presents both the estimates of disease burden for 2019 and a review of the updated official estimates of global progress in the fight against malaria in the first 2 decades of the 21st century (2000–2019).

To provide the historical context to help interpret the trends, the report also looks back at the key events and milestones that have shaped the global malaria effort over the past 20 years (Section 2). Section 3 presents the global trends in malaria morbidity and mortality, and estimates of the burden of malaria during pregnancy. Progress towards elimination is presented in Section 4. An update of the trends and response in the 11 highest burden countries are presented in Section 5, while Section 6 focuses on the total funding for malaria control and elimination, and for malaria research and development. The supply of key commodities to endemic countries and population-level coverage achieved through these investments is presented in Section 7. Section 8 summarizes globally, by region and country, progress toward the GTS milestones for 2020 and the trajectory towards 2025 and 2030. Section 9 describes the threats posed by *Plasmodium falciparum* parasites that no longer express histidine-rich protein 2 (HRP2), which is detected by the most widely used malaria rapid diagnostic test (RDT), and by drug and insecticide resistance. Section 10 describes the malaria response during the COVID-19 pandemic. Section 11 summarizes the findings of the report, and discusses the findings within the context of the COVID-19 pandemic and the future of the fight against malaria.

The main text is followed by annexes that contain data sources and methods, regional profiles and data tables. Country profiles are presented online (<https://www.who.int/teams/global-malaria-programme>).



# Malaria milestones, 2000–2020

It took almost 30 years from the end of the Global Malaria Eradication Programme (in 1969) for malaria to re-emerge as a public health priority in global health and development discourse (6–8). Although data from 1969 to 2000 are scarce, this period was characterized by a sense of failure and abandonment in the fight against malaria. During these 3 decades, hundreds of millions of people were infected with malaria, tens of millions – mostly in sub-Saharan Africa – died, millions of households failed to emerge out of poverty as they struggled with catastrophic health expenditures, hundreds of thousands of pregnant women died during delivery due to malaria-related complications, and millions of children were born with low birthweight, potentially leading to early death or lifelong disability. Millions of children who survived struggled with learning as they dealt with frequent absenteeism due to multiple episodes of malaria, chronic anaemia, seizures or cognitive impairment – consequences of infection and severe disease. Huge blows were dealt to the growth of already weak post-independence national economies, and their attempts to build viable health systems were hampered by lost productivity and high demand for health care.

Against this background, the first 2 decades of the 21st century represent a golden era in the history of malaria control. The world pulled together to fight malaria, delivering one of the biggest returns on investment in global health. The unprecedented scale-up of malaria interventions over this period has led to considerable reductions in disease incidence and mortality. These efforts coincided with other trends and changes that have had a positive impact on malaria, including a period of considerable economic growth and development, infrastructure and housing improvements, rapid urbanization, and general improvements in health systems and population health. By the end of 2019, about 1.5 billion malaria cases and nearly 7.6 million deaths had been averted since the beginning of the century (Section 3). The indirect effect of these gains on the overall health of populations and economies is poorly documented, but is likely to be substantial. In recent years, however, progress has stalled, at a time when we are still dealing with very high levels of malaria burden, re-emphasizing the need to do a lot more to sustain the gains, accelerate progress and achieve the global ambition of a malaria free world (9).

This section reflects on the key malaria milestones in the past 2 decades and the preceding events that laid the foundation. The aim is not to present a comprehensive review of the malaria journey across this period, but rather to highlight some of the major global and regional events that shaped the direction we have travelled. A summary timeline is presented in Fig. 2.1.



## 2.1 LAYING THE FOUNDATIONS

Following decades of not being a global priority, the 1990s laid both the political and scientific foundations for a renewed response to malaria. There were no

reliable estimates of the global burden, but the situation was considered alarming, particularly in sub-Saharan Africa, because the disease was seen both as



the biggest killer of children and a major obstacle to socioeconomic development (1). Malaria control programmes were weak, little effective vector control was being deployed, and access to treatment was limited. Furthermore, the efficacy of chloroquine (CQ), the most commonly used antimalarial for both treatment and prevention, was rapidly declining, resulting in further increases in malaria mortality (10).

This situation triggered key political events in the 1990s that helped to shape progress in the following 2 decades. The Ministerial Conference on Malaria, held in Amsterdam, the Netherlands, in 1992, endorsed a new WHO Global Strategy Malaria Control to guide the response (11). In 1996, the WHO Member States at the 49th World Health Assembly called for the establishment of a special programme on malaria, considering malaria control as an integral part of primary health care (12). This led to an initial investment of US\$ 20 million<sup>1</sup> from WHO (from unspecified funds of the Director-General), to launch the “Accelerated Implementation of Malaria Control” in Africa (13). In June 1997, at its Assembly of Heads of State and Government, the Organization of African Unity (OAU) released the Harare Declaration on Malaria Prevention and Control (6) – the first formal political commitment in Africa to place malaria within the context of African economic recovery and development. Two months later, the Multilateral Initiative on Malaria (MIM) was launched in Dakar, Senegal, at the first Pan-African Malaria Conference (14). This unprecedented gathering brought together leading researchers and academics working on malaria, heads of African malaria control programmes and key international research institutions. In October 1998, the Director-General of WHO, Dr Gro Harlem Brundtland, launched the Roll Back Malaria (RBM) initiative, established through a partnership between WHO, the World Bank, the United Nations Children’s Fund (UNICEF) and the United Nations Development Programme (UNDP) (15).

During the 1990s, and in the face of limited capacity and financial resources for research and development, the WHO-hosted Special Programme for Research and Training in Tropical Diseases (TDR) (16), and some of the leading international research funders supported the early seminal malaria trials and studies. Large-scale trials in Africa documented the efficacy of pyrethroid-impregnated insecticide-treated mosquito nets (ITNs) in preventing malaria and mortality (17, 18). Early trials of artemisinin-based combination therapies (ACTs) also suggested that these therapies were

efficacious and were expected to reduce the risk of resistance developing and spreading (19, 20). Research on the use of malaria medicines for chemoprevention to reduce severe disease and death among the key target groups (infants, children aged under 5 years and pregnant women) was in progress (21–23). The search for a malaria vaccine intensified, and clinical trials of candidate products began in Africa (24).

In 1998, the INDEPTH Network was established as a network of health and demographic surveillance systems that provide detailed and accurate data on health and population problems in low- and middle-income countries (LMICs) (25). Many of the members of the INDEPTH Network were those that undertook the early ITN and antimalarial trials and studies that helped inform malaria control subsequently. By 1998, WHO had made a recommendation for the use of sulfadoxine-pyrimethamine (SP) for intermittent preventive treatment in pregnancy (IPTp) (26). By 1999, the WHO Pesticides Evaluation Scheme (WHOPES) had recommended the use of pyrethroid-impregnated nets for malaria prevention on the basis of safety, efficacy and quality (27, 28). In 1999, a study confirming the presence of vector resistance to pyrethroids, already extensively used in agriculture and in coils and aerosols for vector control, was published (29). To tackle the rapidly evolving threat of CQ resistance, many countries adopted SP as their first-line treatment of uncomplicated malaria. SP was used widely across Africa through a largely presumptive approach to malaria treatment. However, evidence of failure of SP in the treatment of clinical malaria soon emerged in many malaria endemic countries (30).


Resistance to CQ and SP emerged at a time when there were hardly any antimalarial drugs in the development pipeline, and pharmaceutical companies considered that it was not commercially attractive to invest in such drugs. In recognition of the looming lack of efficacious alternative drugs, the Medicines for Malaria Venture (MMV) was established in November 1999; the aim was to facilitate the discovery, development and delivery of efficacious and affordable antimalarial drugs (31). MMV has since been a leader in the product development partnership for drugs for the prevention and treatment of malaria.

By the end of the 20th century, momentum for a global response to malaria had started, but most malaria endemic countries did not have the resources to mount such a response.

<sup>1</sup> All US\$ figures used in this section have been converted to constant 2019 US\$.

# 2.2

2000–2004



Renewed political commitment, the establishment of RBM as a cabinet project of the WHO Director-General and the growing availability of better tools to fight the disease were all instrumental in the signing of the Abuja Declaration at the African Heads of States and Governments Summit, held in Abuja, Nigeria, on 24–25 April 2000 (1). The overarching aim of the Abuja Declaration was to “Halve the malaria mortality for Africa’s people by 2010, through implementing the strategies and actions for Roll Back Malaria”. This was to be achieved through multiple approaches to ensure that, in malaria endemic Africa, 60% of malaria patients had access to prompt effective treatment, 60% of children aged under 5 years and pregnant women were protected with ITNs, and 60% of pregnant women received presumptive intermittent treatment to alleviate the consequences of malaria infection to the mother and her unborn child. African countries had also committed to achieving expenditure of 15% of gross domestic product (GDP) on health by 2015. They urged donor countries to “fulfil the yet to be met target of 0.7%” of their gross national product (GNP) as official development assistance (ODA) to developing countries (1). The Abuja Declaration was further reinforced by the Group of Eight (G8) countries, during the Okinawa Summit in Japan in July 2000, committing to the target of reducing malaria mortality by 50% by 2010 (32).

In September 2000, the framework of eight MDGs was launched during the Millennium Summit at the United Nations (UN) headquarters in New York (2). Under the MDGs, there was a clear articulation that malaria was a global development issue, with emerging research documenting more clearly the considerable toll of the disease on economic development in endemic countries (33). MDG target 6C required the halting of the malaria epidemic and the reversal of incidence and death rates associated with malaria (34). This strengthened the calls made in the Harare and Abuja

declarations, and by the RBM initiative, for a globally funded partnership to fight malaria, to save lives and to accelerate economic growth in affected countries.

In 2000, in response to reduced efficacy of CQ and SP for the treatment of clinical malaria, WHO published recommendations for the use of ACTs (35). In 2001, the initial evidence of delayed parasite clearance with artesunate was reported in Cambodia (36). The previous year, WHO also recommended the use of RDTs in health facilities, as increasingly accurate and affordable tests became available (37). This led to a major shift away from what had been a predominantly syndromic approach – with the presumptive treatment of all fevers for malaria – to an approach based on pretreatment parasitological confirmation of malaria. This improved the rational use of ACTs and has also subsequently enhanced the value of routinely reported data on malaria burden. However, parasitological diagnosis continues to be used at modest levels, especially in sub-Saharan Africa (**Section 7**).

In 2000, the Bill & Melinda Gates Foundation was established; it is now one of the largest private foundations in the world (38). In its work on malaria, the foundation has focused on development of new vaccines, diagnostics, medicines and vector control products and their delivery and use in public health, while advancing improved surveillance systems and data analytics.

Several new institutions, programmes and initiatives soon followed. In May 2001, the European Union launched the “Programme for accelerated action on HIV/AIDS, malaria and tuberculosis in the context of poverty reduction”, which also led to the creation of the European and Developing Countries Clinical Trials Partnership (EDCTP). Founded as a public–public partnership between countries in Europe and sub-



2005–2010

2011–2015

2016–2019

Saharan Africa, the EDCTP is supported by the European Union (39). The EDCTP aims to accelerate clinical development of vaccines, diagnostics and medicines for infectious diseases of the poor, and has been a major investor in malaria clinical trials (40), several of which have contributed to the development of global normative guidance by WHO. In 2003, the Foundation for Innovative New Diagnostics (FIND) was established as a global non-profit organization, with the aim of accelerating innovation in the development and delivery of diagnostics of infectious diseases of the poor (41). As a WHO Collaborating Centre for Laboratory Strengthening and Diagnostic Technology Evaluation, FIND has supported the generation of evidence for malaria diagnosis policies, producing regular reports on the quality and performance on RDTs.

In 2002, the Global Fund to Fight AIDS, Tuberculosis and Malaria (the Global Fund) was created, marking the beginning of an unprecedented period for malaria funding (42). The Global Fund was originally conceived as a financing mechanism for HIV/AIDS but ministers of health, especially from the WHO African Region, called for it to be extended to malaria (34, 43).


In recognition of the terrible toll of malaria on children and pregnant women, UNICEF stepped up its key role in the malaria response, in addition to being one of the founders of RBM. UNICEF's focus was on strengthening community-based and local action to improve child health and nutrition. By the early 2000s, it was one of the world's largest global procurers of ACTs, ITNs and subsequently long-lasting insecticidal nets (LLINs), supporting the delivery of nets during routine and mass vaccination campaigns (44). UNICEF continues to support the scale-up of diagnosis and treatment of malaria at the community level, through integrated delivery platforms and support for the delivery of seasonal malaria chemoprevention (SMC) (45).

The path from the promising results from the field trials of the efficacy of ITNs and ACTs to scaling these up in malaria endemic countries remained challenging. There were the limited supply of ITNs and ACTs, their high costs and the lack of substantial domestic or external funding for malaria control to scale-up new interventions for prevention and treatment. In 2002, the WHO RBM initiative published a framework for scaling up ITNs in Africa. The framework proposed two key elements: sustained subsidies strictly targeted to vulnerable groups, and a strengthened and expanded commercial market that would provide ITNs at the lowest possible prices for the general population (46). The consensus at the time was not in favour of delivering ITNs to the whole population or providing ITNs at no cost, even to vulnerable groups, mainly because of concerns about the financial sustainability of doing so. Instead, subsidized distribution through social marketing and mother and child clinics became the norm. The overarching aim was to catalyse the growth of commercial markets to meet the demand for ITNs and reduce commodity prices (46).

The *Africa malaria report*, a precursor to the world malaria report, was published in 2003 (47). Despite many important developments, by the end of 2004, most mosquito nets were still conventional ITNs (i.e. they required frequent retreatment), and their use by children aged under 5 years was only 2% (48). Although recommended by WHO since 1998, IPTp scale-up had barely started, only 42% of children with fever sought treatment and received antimalarials, and most malaria treatment was presumptive and predominantly with CQ or SP, which were no longer recommended for treatment by WHO.

# 2.3

2000–2004



In March 2005, the first meeting on the replenishment of the Global Fund took place in Stockholm, Sweden (49). At the end of the replenishment process, US\$ 3.7 billion was pledged to the Global Fund for the period 2006–2007, of which about US\$ 760 million was eventually committed to malaria control (50). In June 2005, the United States President’s Malaria Initiative (PMI) was launched, targeting support to Angola, Uganda and the United Republic of Tanzania (51). By the end of the decade, PMI had extended its support to 12 additional countries in Africa (52).

The injection of funding came at an important time. At the end of 2005, WHO released the first world malaria report, presenting global progress on malaria in the period 2000–2004 (48). The report showed that the malaria burden remained high, with 1 million estimated deaths, mainly in sub-Saharan Africa, and that access to malaria prevention and treatment had barely improved since 2000. In 2005, the WHO RBM initiative published a strategy for improving access to treatment through home management of malaria (53). In the same year, WHO also published a recommendation to use artesunate and artemisinin suppositories for pre-referral treatment of severe malaria (54).

Measuring the burden of malaria and progress in intervention was proving to be a difficult task. Also, as funding increased, a credible measure of the impact of the investment was increasingly seen as critical to make the case for further funding. Surveillance systems in malaria endemic countries remained weak, and most reported malaria case data were not based on parasitological diagnosis. There was limited understanding of the subnational malaria epidemiology to effectively guide investments. In 2004, the RBM Monitoring and Evaluation Reference Group (MERG), with funding support from the United States Agency for International Development (USAID), began the process of developing a malaria indicator survey toolkit (55). The toolkit was intended to support standalone malaria-specific surveys or malaria modules included in standard demographic and health surveys (DHS) (56) or UNICEF-supported multiple indicator cluster surveys (MICS) (57). These surveys have since been the backbone of understanding infection prevalence and malaria intervention coverage in communities in Africa, and in the tracking of global progress annually through the world malaria report. Since 2006, over 100 surveys with malaria-related information have been conducted, mainly in sub-Saharan Africa.

Creation of new partnerships and initiatives continued. To respond to the threat of insecticide resistance, innovation was needed to develop new vector control solutions. In 2005, the Innovative Vector Control Consortium (IVCC) was established as a partnership of industry, the public sector and academia (58). As the main product development partnership for malaria vector control, IVCC has worked with a range of partners to facilitate the development of novel and improved public health insecticides, formulations and products to address these challenges. It has also supported field research and efforts to improve access to these tools through its global access strategy (59). In 2006, Unitaid was established as an agency that is hosted and administered by WHO; Unitaid’s mission is to scale up access to treatment for HIV/AIDS, malaria and tuberculosis in developing countries through price reductions of drugs and diagnostics, and improved availability (60). Unitaid has used an innovative financing approach – the solidarity levy on airline tickets imposed by France and other countries. Since its establishment, Unitaid’s investment in malaria prevention, diagnosis and treatment has developed into a large portfolio (61).

Faced with weak health systems and low domestic funding, approaches to scale-up of interventions remained challenging. Until 2007, the recommendation was still to prioritize coverage of ITNs to key target groups in sub-Saharan Africa; however, it was estimated that by 2007 only 15% of children aged under 5 years and pregnant women were sleeping under an ITN (50). The dominant channels for ITN distributions were social marketing of nets and continuous distributions in health facilities, with the latter moving from being highly subsidized to being free in some countries from around this time (62). General case management practice was also to treat any febrile child as a malaria case, often presumptively, because RDTs had not been widely scaled up and microscopy was limited mainly to large urban health facilities.

In August 2007, supported by evidence from Kenya (63), the WHO Global Malaria Programme (GMP) released a position statement in which it recommended that “insecticidal nets be long-lasting, and distributed either free or highly subsidized and used by all community members” and noted that “... free mass distribution of LLINs is a powerful way to quickly and dramatically increase coverage, particularly among the poorest people” (64). This statement laid the foundation for ITNs becoming by far the largest investment in a single malaria intervention. Free mass campaigns to cover





## 2005–2010

individuals of all ages with LLINs, and continuous distribution channels to sustain coverage, were launched and marked the beginning of a rapid increase in ITN coverage in sub-Saharan Africa (**Section 7**). Although studies showed significant reduction in parasite prevalence following universal coverage, the decision to implement universal coverage was driven primarily by coverage and equity aims rather than comprehensive cost-benefit analysis.

For decades, fuelled by the sense of failure following the first eradication campaign of the 1950s and 1960s, the world had shied away from placing eradication of malaria within its goals (24). However, buoyed by the increasing global commitment to fight malaria, the opportunities to rapidly expand the scale-up of interventions and results from the development of new tools, including vaccines – Bill and Melinda Gates (of the Bill & Melinda Gates Foundation) made a global call for a renewed commitment to eradicate malaria (65) and WHO Director-General Dr Margaret Chan publicly endorsed that vision. This triggered a global discussion on the feasibility of malaria eradication and its critical dependence on the development of new and improved tools. No timeline for that effort was defined.

Shortly after, the RBM Partnership – by now a partnership entity hosted within WHO – released the Global Malaria Action Plan for a malaria free world (GMAP) (66). This plan built on the WHO call for universal coverage and the emerging discussions on malaria eradication. A 2010 target was assigned to achieve universal coverage, reduce malaria morbidity and mortality by half from a 2000 baseline, and eliminate malaria in 8–10 countries. Also explicitly stated in the GMAP was a target of achieving near-zero preventable deaths by 2015, and of malaria eradication through progressive elimination in countries, without a defined date for its achievement. The plan outlined three strategic components with research as a supporting component: scale-up for impact, sustained control and elimination.

Following the first, second and third replenishments (in 2005, 2007 and 2010, respectively), Global Fund resources for malaria increased considerably (67). External investment in malaria was estimated to be US\$ 450 million in 2005, with an estimated US\$ 1 billion spent in the period 2000–2005 (**Section 6**). Increasing access remained the key challenge. Although ITNs were moving from a social marketing scheme towards mass distribution campaigns, new delivery mechanisms were being developed with regard to ACTs (68, 69). In 2008,

## 2011–2015

the Global Fund assumed funding responsibilities, with support from Unitaid, for the Affordable Medicines Facility-malaria (AMFm) as a pilot programme that aimed to take advantage of the relative high use of the private retail sector for treatment of fever, and thus expand access to quality-assured ACTs (68, 69). An evaluation funded by the Global Fund showed that positive achievements included increased availability of ACTs, reduced prices, increased market share and minimal disruption of supplies to the public sector (70). However, given the low levels of parasitological diagnosis and by not subsidizing diagnostic testing in the private sector, the AMFm failed to fully target the subsidized ACTs to those with malaria.

In 2007, confirmation of what was then called partial artemisinin resistance was established in the area of the Thai–Cambodia border, and in 2008 the first clinical cases due to malaria parasites containing gene deletions causing false negative RDTs were described in Peru (36, 71). Since then, monitoring and mitigating ACT resistance has become a major focus of the global malaria community; also, deletions in the *P. falciparum* genes for HRP2 (*pfhrp2*) have emerged in sub-Saharan Africa, and recent evidence suggests worrying levels of prevalence in Horn of Africa countries (**Section 9.1**).

In 2009, the African Leaders Malaria Alliance (ALMA) was established as a forum to provide visibility at high levels of political leadership for the response against malaria in Africa (72).

On the policy front, WHO released a recommendation on the use of intermittent preventive treatment in infants with SP (IPTi-SP) in 2010, following evidence of modest efficacy from pooled analysis of randomized control trials in Gabon, Ghana, Kenya, Mozambique and the United Republic of Tanzania (73).

In 2010, the US National Institute of Allergy and Infectious Diseases established 10 International Centers for Excellence for Malaria Research to support multidisciplinary malaria research across diverse settings in Africa, Asia Pacific and South America (74).

The high-level attention, increased funding and successful implementation of technical strategies were beginning to contribute to a positive impact. By 2010, it was estimated that, globally, substantial reductions in malaria morbidity and mortality had been reported (75). WHO certified United Arab Emirates in 2007 and Morocco and Turkmenistan in 2010 as malaria free (**Section 4**).

## 2016–2019

# 2.4

2000–2004



In 2011, WHO established the Malaria Policy Advisory Committee (MPAC) to provide independent advice to WHO on developing policy recommendations to control and eliminate malaria, and thus improve the quality and independence of the malaria policy-making process. The MPAC is an independent advisory group that aims to bring together the world's foremost experts on malaria to provide strategic technical guidance to the WHO Director-General as part of a transparent, responsive and credible policy-setting process on malaria (76).

In 2011, PMI added the countries of the Greater Mekong subregion (GMS), the Democratic Republic of the Congo, Guinea, Nigeria and Zimbabwe to its list of countries to receive support; this brought its tally of support to 20 high burden African countries (52).

By the end of 2011, global sales of ACTs had exceeded 500 million treatment doses, marking a period of sustained scale-up of effective malaria interventions (77). However, artemisinin resistance was expanding in the GMS and was considered as a potential threat to the global malaria enterprise (36, 78). Learning from the experience of poor mitigation of resistance to previous antimalarials, WHO mobilized the global community by launching the Global Plan for Artemisinin Resistance Containment (GPARC) (79).

In 2012, WHO and partners launched the Mekong Malaria Elimination (MME) programme (78, 80). This

is a multi-country (Cambodia, China, Lao People's Democratic Republic, Myanmar, Thailand and Viet Nam) programme to fight artemisinin resistance, primarily through accelerated progress towards malaria elimination by 2025, focusing especially on *P. falciparum* malaria. To support the MME programme, the Global Fund launched the Mekong Regional Artemisinin-resistance Initiative (RAI) in 2013, and has invested considerable resources (nearly US\$ 600 million) in the subregion since then (81). Dramatic progress has been achieved in the GMS since the launch of the MME programme, and most countries are on target to achieve *P. falciparum* elimination by 2025 (Section 4); also, there is to date no evidence of a spread of artemisinin resistance from the GMS to other parts of the world (Section 9).

In 2012, the Global Fund launched its second strategy for achieving impact through its investments across five strategic objectives: invest more strategically, evolve the funding model, actively support grant implementation success, protect and promote human rights, and sustain the gains and mobilize resources (82). Also in 2012, the Global Fund decided to integrate AMFm into core grant management processes through an orderly transition in 2013, allowing countries to use some of their core grants to implement AMFm as part of a co-payment mechanism (83).

On the global policy front, in 2012 WHO published a recommendation for the use of SMC in children in high



2005–2010

2011–2015

2016–2019


burden and highly seasonal malaria transmission areas, in response to evidence of the strong impact on malaria morbidity (84). In support of the scale-up of SMC, Unitaid launched the ACCESS-SMC project in 2013 (85), the Global Fund mainstreamed the intervention into core grants in 2017 (86), and PMI expanded its support for SMC activities in Benin, Burkina Faso, Cameroon, Ghana, Guinea, Mali, Niger, northern Nigeria and Senegal (87).

In 2012, WHO published the Global Plan for Insecticide Resistance Management in Malaria Vectors (88) as a response to mitigate the spread of insecticide resistance. Adding to the list of biological threats to the global malaria fight, 2014 saw the first evidence of the presence of an *Anopheles stephensi*, an efficient urban malaria species in Asia and Persian Gulf, being reported in sub-Saharan Africa, in Djibouti, where it was implicated in a malaria epidemic (89). Since then, *An. stephensi* has been reported to be established in Ethiopia and is efficient in transmission of both *P. vivax* and *P. falciparum* (90).

Following the call for malaria eradication by Bill and Melinda Gates, several scientific publications – for example, those from the Malaria Eradication Scientific Alliance published in *PLoS Medicine* (91) and the Lancet series on malaria elimination (92) – re-energized the debate on feasibility, approaches and innovation towards malaria elimination and its eventual eradication. At the same time, the application of novel

geospatial methods to the growing number of community parasite prevalence surveys in sub-Saharan Africa began to create a clearer picture of the geographical distribution of *P. falciparum* malaria subnationally (93–96). This increased granularity of malaria risk mapping exposed underlying heterogeneity and the need for strategic planning and resource allocation at subnational levels (Section 5).

Some 15 years after the launch of the MDGs, analysis presented in the *World malaria report 2015* (97) suggested that the target of reversing the malaria trends had been achieved. It was estimated that malaria case incidence had reduced by 37% and mortality rate by 60% between 2000 and 2015. An estimated 438 000 people had died of malaria in 2015; thus, the near-zero death target of the GMAP had not been achieved (66). These major declines in the malaria burden were considered conclusive evidence of achieving, or even surpassing, the MDG target 6C, and were hailed as showing the remarkable strides that could be made with adequate investment and political commitment. Three years earlier, in anticipation of the end of the MDGs, the UN Conference on Sustainable Development was convened in Rio de Janeiro, Brazil, where Member States decided to develop a set of SDGs to build on the MDGs, and to establish the UN High-level Political Forum on Sustainable Development (98). Progress in malaria incidence and mortality rate were recognized as key indicators in SDG Goal 3, target 3.3, which stated “By



2030, end the epidemics of AIDS, tuberculosis, malaria and neglected tropical diseases and combat hepatitis, waterborne diseases and other communicable diseases”; that goal had a target of halving malaria case incidence by 2020, and contributing to the ending of preventable deaths of neonates and children aged under 5 years by 2030, from a baseline of 2015 (3, 99).

Among the global malaria community, there was now consensus on the need to develop a coherent and even more ambitious global strategy, not only to sustain the gains, but also to ensure accelerated progress and align with the SDGs. In 2012, the MPAC discussed the proposal to develop a global technical strategy, and recommended it to the WHO Director-General. The GTS was formally adopted by the Sixty-eighth World Health Assembly in May 2015, in resolution WHA68.2 (4). With a vision of a world free of malaria, and underpinned by five guiding principles, the GTS included three pillars, two supporting elements and four impact goals across three milestone years (2020, 2025 and 2030) using a 2015 baseline (**Table 2.1**). For the first time, transforming surveillance systems was affirmed as a core intervention, recognizing the critical function of reliable information in improving the efficiency and effectiveness of interventions to prevent and treat malaria.

As an investment case for the GTS, the RBM Partnership to End Malaria developed the investment plan AIM (5). Anchored in a strong partnership, with a multisectoral

and coordinated approach, the plan outlines core areas of focus: mobilizing resources; strengthening multisectoral and intercountry collaboration; keeping people at the centre of the response; strengthening the enabling environment; fostering and sharing innovations and solutions; and facilitating change. Both the GTS and the AIM acknowledged that strengthened health systems would be needed, because these would determine the rate of progress towards the bold targets. It was hoped that the adoption by countries of the GTS and AIM would also contribute to the post-2015 SDGs.

By 2015, over 1 billion ITNs had been distributed globally, accounting for the largest proportion of donor investment in malaria. Modelling analysis suggested that, among malaria interventions, use of ITNs was the largest contributor to the reduction in the burden of malaria in sub-Saharan Africa (93). By the end of this period, however, pyrethroid resistance had increased both in terms of geography and intensity (100).

Armenia and Maldives were certified by WHO as free of malaria in 2011 and 2015, respectively. The *Malaria Elimination Strategy in the GMS 2015–2030* was endorsed by the MPAC and adopted by health ministers in GMS countries in 2015; its goals were to eliminate *P. falciparum* malaria in 2025 and all malaria in 2030 in the subregion (101).



2005–2010

2011–2015

2016–2019

**TABLE 2.1.**

**GTS: global targets for 2030 and milestones for 2020 and 2025** *Source: GTS (4).*

**Vision – A world free of malaria**

**Principles**

1. All countries can accelerate efforts towards elimination through combinations of interventions tailored to local contexts
2. Country ownership and leadership, with involvement and participation of communities, are essential to accelerating progress through a multisectoral approach
3. Improved surveillance, monitoring and evaluation, as well as stratification by malaria burden, are required to optimize the implementation of malaria interventions
4. Equity in access to health services, especially for the most vulnerable and hard-to-reach populations, is essential
5. Innovation in tools and implementation approaches will enable countries to maximize their progression along the path to elimination

**Pillars**

Pillar 1	Ensure universal access to malaria prevention, diagnosis and treatment
Pillar 2	Accelerate efforts towards elimination and attainment of malaria free status
Pillar 3	Transform malaria surveillance into a core intervention

**Supporting elements**

Supporting element 1. Harnessing innovation and expanding research

Supporting element 2. Strengthening the enabling environment

Goals	Milestones 2020	2025	Targets 2030
1. Reduce malaria mortality rates globally compared with 2015	At least 40%	At least 75%	At least 90%
2. Reduce malaria case incidence globally compared with 2015	At least 40%	At least 75%	At least 90%
3. Eliminate malaria from countries in which malaria was transmitted in 2015	At least 10 countries	At least 20 countries	At least 35 countries
4. Prevent re-establishment of malaria in all countries that are malaria free	Re-establishment prevented	Re-establishment prevented	Re-establishment prevented

GTS: *Global technical strategy for malaria 2016–2030.*

# 2.5

2000–2004



Following the launch of the GTS and AIM, many WHO regions and national programmes launched their own aligned strategies. In June 2016, the RBM initiative previously hosted by WHO was renamed the RBM Partnership to End Malaria, with new hosting arrangements under the UN Office for Project Services (102). In 2017, the Global Fund launched its new strategy for the period 2017–2022, titled Investing to end epidemics, with four strategic objectives: maximizing impact, promoting and protecting human rights and gender equality, building resilient and sustainable systems for health for all, and mobilizing increased resources (103). Building on the ALMA experience, the Asia Pacific Leaders Malaria Alliance (APLMA) was launched in 2017 (104). In the same year, PMI extended its support to include the GMS and five additional African countries (52).

As part of the commitment to achieving Goal 3 of the GTS (i.e. ensuring at least 10 countries reach malaria elimination by 2020), in April 2017, WHO launched the “eliminating countries for 2020” (E-2020) initiative (105). Twenty-one countries that had made substantial progress over the past decade and were considered close to elimination were selected to take part in the E-2020 (Section 4). By 2018, the Goal 3 milestone for 2020 was already on target, with 10 countries that were malaria endemic in 2015 expected to be malaria free by 2020 (77). Since 2015, Kyrgyzstan (2016), Sri Lanka (2016) and Uzbekistan (2018) have been certified by WHO as malaria free. Paraguay (2018) and Algeria (2019), both E-2020 countries, each became the first country in their respective region to be certified malaria free since 1973. Argentina (2019) followed

Paraguay to become the next country in the WHO Region of the Americas to be certified. The *Ministerial Declaration on Accelerating and Sustaining Malaria Elimination in South-East Asia Region* was signed in November 2017, to accelerate malaria elimination in this region (106).

In contrast to the impressive progress on the GTS elimination goal, estimates published in the *World malaria report 2017* showed that the morbidity and mortality goals were off track, and that gains were beginning to reverse in some countries (107). The main theme of the report was that the malaria world was at a “crossroads”, and an urgent response was required to kickstart the stalling progress (8, 104). The Director-General of WHO, Dr Tedros Ghebreyesus, declared:

***The data showed that less than half of countries with ongoing transmission were on track to reach critical targets for reductions in the death and disease caused by malaria. Progress appeared to have stalled ... The choice before us is clear. If we continue with a ‘business as usual’ approach – employing the same level of resources and the same interventions – we will face near-certain increases in malaria cases and deaths.***



2005–2010

2011–2015

2016–2019


This call led to the formation of the high burden to high impact (HBHI) response coordinated by WHO and the RBM Partnership to End Malaria and led by endemic countries (108). The formal launch of the HBHI approach was held in Maputo, Mozambique, in November 2018, during the 20th anniversary of the RBM Partnership. The approach is based on four response elements: galvanizing political will nationally and globally to reduce malaria deaths; using strategic information to drive impact; implementing best global guidance, policies and strategies suitable for all malaria endemic countries; and applying a coordinated country response (108). This approach has been led by 11 countries that accounted for 70% of the global burden of malaria: Burkina Faso, Cameroon, the Democratic Republic of the Congo, Ghana, India, Mali, Mozambique, Niger, Nigeria, Uganda and the United Republic of Tanzania. Since then, the launch of the HBHI approach has been formally initiated in all countries except in Mali (where HBHI-related activities are underway). In all initiation meetings, there was high-level government and partnership participation, with strong commitment to support the approach.

Since 2018, WHO, the RBM Partnership to End Malaria and collaborating partners have supported the HBHI countries to develop robust national malaria strategic plans (NMSPs), and to prioritize resources using subnational tailoring of interventions, driven by epidemiological, ecological and health system data, and other information (**Section 5**). WHO has embarked on a process to improve the predictability, timeliness and transparency of its policy-making process, and to produce the first set of WHO consolidated malaria guidelines. The aim is to re-position its policy recommendations, moving away from a prescriptive

set of statements to instead providing problem-solving tools for countries to adapt, and inculcating an approach of subnational tailoring of interventions based on local data. As a first step, WHO published a compendium of all policies, clarifying the distinction between actual *recommendations*, which are based on thorough, systematic reviews of the evidence by a guideline development group (109), and *best practice statements*, which are designed to help countries implement policies but should not be considered restrictive. These concepts were further crystallized through a technical brief (110) to countries, to support national malaria programmes (NMPs) making funding requests to the Global Fund and other organizations.

In October 2019, during its Sixth Replenishment Conference in Lyon, France, the Global Fund managed to raise the highest level of funding since its inception, with a commitment of US\$ 14 billion (111). Of this amount, US\$ 4.8 billion was allocated to malaria, an increase of over US\$ 1 billion from the previous allocation period. PMI funding also increased to US\$ 755 billion in 2019 (52).

Incremental improvements to the tools available for malaria control have continued; for example, another ACT (pyronaridine-artesunate) has been developed (112), as have mosquito nets treated with insecticides other than pyrethroids (these are currently undergoing evaluation). In 2016, WHO released a position paper on the world's first malaria vaccine to have received a positive recommendation from the European Medicines Agency (EMA). As part of a collaboration between WHO, PATH, GlaxoSmithKline (GSK), the Global Fund, Gavi and Unitaid, GSK's RTS,S vaccine is undergoing a phased pilot introduction through routine



childhood immunization services in parts of Malawi, Ghana and Kenya, which started in 2019. Some 12 months on, about 500 000 children have been reached with their first dose of the vaccine. An ongoing evaluation is assessing the public health value of the vaccine as a complementary tool that could be added to the existing preventive, diagnostic and treatment measures recommended by WHO.

At about the same time as evidence was emerging that progress towards GTS milestones for burden reduction had stalled and the global community was grappling with ways to support countries to get back on track, active discussions were happening about whether malaria eradication with a defined timeline was feasible (113). In 2016, the then Director-General of WHO, Dr Margaret Chan, established a strategic advisory group tasked with analysing future scenarios for malaria, including the feasibility and expected cost of eradication. The Strategic Advisory Group for Malaria Eradication (SAGme) concluded its work in 2019. Based on SAGme's work, WHO reaffirmed its position on malaria eradication and the importance of investing in universal health coverage (UHC) through a statement by the Director-General, Dr Tedros Ghebreyesus:

This statement was released as part of the WHO push to renew the momentum, to ensure the

**WHO continues to unequivocally support the goal of malaria eradication. To achieve this vision, we must deliver on our promises: to increase domestic and international investments in health; reduce malaria in the highest-burden countries; achieve universal health coverage; ensure no child dies from a preventable disease; and leave no one behind in pursuit of health and development goals because they were born poor. By delivering on these promises and investing in the development of transformative new tools, the world can achieve the health-related Sustainable Development Goals and eradicate malaria.**





2005–2010

2011–2015

2016–2019

establishment of strong primary health care systems through the UHC approach encapsulated in the Astana Declaration of 2018 (114). This declaration was signed by heads of state and government, ministers, and representatives of states and governments during the Global Conference on Primary Health Care held in Astana, Kazakhstan, on 25–26 October 2018. WHO did not define a specific timeline for malaria eradication; instead, it identified a focus on burden reduction and sequential elimination in malaria endemic countries and regions as a logical path forward. To this end, SAGme proposed focused efforts in four areas: research and development of new tools; improved access to affordable, quality, people-centred health services; enhanced surveillance and response; and formulation of subnational, national and regional strategies (113). At around the time that WHO released the SAGme report, the Lancet Commission on Malaria Eradication published a collection of work on the feasibility and affordability of malaria eradication by 2050 (115).


In September 2019, at the UN high-level meeting “Universal Health Coverage: Moving Together to Build a Healthier World”, the political commitment was secured for implementing high-impact health interventions to combat diseases, protect women’s and children’s health, and ensure no one suffers financial hardship. There was commitment for investing in everyone’s health, expanding quality health services

and reaching the most marginalized populations. This would require improved efficiency and equity in the allocation and use of existing resources – based on local context and priorities, and governed by data to identify those in need of interventions (116).

Although major system weaknesses and data quality issues remain, the period 2016–2019 has also been one of considerable progress in the strengthening of health information systems in malaria endemic countries. By 2018, more than 50 malaria endemic countries had installed District Health Information Software 2 (DHIS2) either for direct data entry at health facilities or as the backbone of aggregated data systems (77). Combined with increasing use of RDTs and increased reporting, the volume and quality of data have improved steadily, with DHIS2 offering flexible data analysis and use capabilities. These improvements have been major contributors to the efforts on subnational tailoring of malaria interventions in HBHI countries (**Section 5**).

By the end of 2019, with the emergence of COVID-19 and its subsequent pandemic spread, much of the progress against malaria was under enormous risk, with the potential to wipe out 20 years of malaria gains (117). To mitigate disruptions of essential malaria services, global and national partners joined forces to support countries to mount a response. The nature of this response and the consequences of the pandemic are described in **Section 10**.

**FIG. 2.1. Key milestones in the fight against malaria in the past 2 decades**



ACT: artemisin-based combination therapy; *An.*: *Anopheles*; BMGF: Bill & Melinda Gates Foundation; DG: Director-General; EDCTP: European & Developing Countries Clinical Trials Partnership; G8: Group of Eight; GMAP: Global Malaria Action Plan; GMP: Global Malaria Programme; GMS: Greater Mekong subregion; GPARC: Global Plan for Artemisinin Resistance Containment; GPIRM: Global Plan for Insecticide Resistance Management in Malaria; GTS: Global technical strategy for malaria 2016-2030; HBHI: high burden high impact; IPTi-SP: intermittent preventive treatment in infants using sulfadoxine-pyrimethamine; IPTp: intermittent preventive treatment in pregnancy; IPTp-SP: intermittent preventive treatment in pregnancy using sulfadoxine-pyrimethamine; ITN: insecticide-treated mosquito net; IVCC: Innovative Vector Control Consortium; IVM: integrated vector management; LLIN: long-lasting insecticidal net; MDG: Millennium Development Goal; MIM: Multilateral Initiative on Malaria; MIS: malaria indicator survey; MME: Malaria Mekong Elimination; MMV: Medicines for Malaria Venture; MTG: malaria treatment guidelines; MVI-PATH: Malaria Vaccine Initiative, PATH; PBO: piperonyl butoxide; PMI: President's Malaria Initiative; RAI: Regional Artemisinin-resistance Initiative; RBM: Roll Back Malaria (before 2016); RDT: rapid diagnostic test; SAGme: Strategic Advisory Group for Malaria Eradication; SDG: Sustainable Development Goal; SMC: seasonal malaria chemoprevention; SME: surveillance, monitoring & evaluation; SSA: sub-Saharan Africa; UNOPS: United Nations Office for Project Services; VC: vector control; WHA: World Health Assembly; WHO: World Health Organization; WHOPEs: WHO Pesticides Evaluation Scheme; WMR: world malaria report.



# Global trends in the burden of malaria

The burden estimates presented in this section are the number of cases and deaths estimated to have occurred between 2000 and 2019, as well as case incidence and malaria mortality rates in the same period. These estimates are then used to compute the number of cases and deaths averted, globally and by WHO region, since 2000. An analysis of the prevalence of exposure to malaria and low birthweights is also presented.



Estimation of the burden of malaria cases and deaths relies on several methods, depending on the quality of the national surveillance systems and the availability of data over time (**Annex 1**). Moderate to high transmission countries in sub-Saharan Africa account for most of the global malaria burden, but they generally have weak surveillance systems. For these countries, estimates of cases are derived using an approach that transforms modelled community parasite prevalence into case incidence within a geospatial framework. Malaria deaths for these countries are also estimated from a cause of death fraction for malaria applied to the trends in all-cause mortality in children aged under 5 years, and to which a factor for malaria deaths among those aged over 5 years is applied. For other countries with stronger

surveillance systems, data are used as reported or cases are estimated by adjusting national data for treatment seeking, testing and reporting rates. Where adjustments are applied to national case data, a species-specific case fatality rate is applied to these data to estimate malaria deaths.

Because these estimates are updated each year, computed malaria cases and deaths change across the period of analysis, and estimates over time may vary in the annual world malaria reports from different years. Also, partly because of the separate methods used to compute malaria cases and deaths in sub-Saharan Africa, trends in the two measures of burden may be different for a given country; thus, caution should be applied in their comparison.

## 3.1 GLOBAL ESTIMATES OF MALARIA CASES AND DEATHS, 2000–2019


Globally, there were an estimated 229 million malaria cases in 2019 in 87 malaria endemic countries, declining from 238 million in 2000 (**Table 3.1**) across 108 countries that were malaria endemic in 2000

(**Fig. 3.1**). At the GTS baseline of 2015, there were 218 million estimated malaria cases. The proportion of cases due to *P. vivax* reduced from about 7% in 2000 to 3% in 2019.



**FIG. 3.1.**

**Countries with indigenous cases in 2000 and their status by 2019** Countries with zero indigenous cases over at least the past 3 consecutive years are considered to have eliminated malaria. In 2019, China and El Salvador reported zero indigenous cases for the third consecutive year and have applied for WHO certification of malaria elimination; also, the Islamic Republic of Iran, Malaysia and Timor-Leste reported zero indigenous cases for the second time. *Source: WHO database.*



WHO: World Health Organization.

**TABLE 3.1.**

**Global estimated malaria cases and deaths, 2000–2019** Estimated cases and deaths are shown with 95% upper and lower confidence intervals. *Source: WHO estimates.*

Year	Number of cases (000)				Number of deaths		
	Point	Lower bound	Upper bound	% <i>P. vivax</i>	Point	Lower bound	Upper bound
2000	<b>238 000</b>	222 000	259 000	6.9%	<b>736 000</b>	697 000	782 000
2001	<b>244 000</b>	228 000	265 000	7.4%	<b>739 000</b>	700 000	786 000
2002	<b>239 000</b>	223 000	260 000	7.1%	<b>736 000</b>	698 000	783 000
2003	<b>244 000</b>	226 000	268 000	7.8%	<b>723 000</b>	681 000	775 000
2004	<b>248 000</b>	227 000	277 000	8.0%	<b>759 000</b>	708 000	830 000
2005	<b>247 000</b>	229 000	272 000	8.3%	<b>708 000</b>	662 000	765 000
2006	<b>242 000</b>	223 000	268 000	7.2%	<b>716 000</b>	675 000	771 000
2007	<b>241 000</b>	222 000	265 000	6.8%	<b>685 000</b>	644 000	735 000
2008	<b>240 000</b>	222 000	264 000	6.5%	<b>638 000</b>	599 000	685 000
2009	<b>246 000</b>	226 000	271 000	6.5%	<b>620 000</b>	572 000	681 000
2010	<b>247 000</b>	226 000	273 000	7.0%	<b>594 000</b>	546 000	658 000
2011	<b>239 000</b>	218 000	262 000	7.2%	<b>545 000</b>	505 000	596 000
2012	<b>234 000</b>	213 000	258 000	6.6%	<b>517 000</b>	481 000	568 000
2013	<b>225 000</b>	206 000	248 000	5.3%	<b>487 000</b>	451 000	538 000
2014	<b>217 000</b>	201 000	236 000	4.3%	<b>471 000</b>	440 000	511 000
2015	<b>218 000</b>	203 000	238 000	3.9%	<b>453 000</b>	422 000	496 000
2016	<b>226 000</b>	210 000	247 000	4.0%	<b>433 000</b>	403 000	478 000
2017	<b>231 000</b>	213 000	252 000	3.4%	<b>422 000</b>	396 000	467 000
2018	<b>228 000</b>	211 000	250 000	3.2%	<b>411 000</b>	389 000	458 000
2019	<b>229 000</b>	211 000	252 000	2.8%	<b>409 000</b>	387 000	460 000

*P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.

### 3 Global trends in the burden of malaria


Malaria case incidence (i.e. cases per 1000 population at risk) reduced from 80 in 2000 to 58 in 2015 and 57 in 2019 (Fig. 3.2). Between 2000 and 2015, malaria case incidence declined by 27% and then by less than 2% in the period 2015–2019, indicating a slowing of the rate of decline since 2015 (Fig. 3.2).

Malaria deaths have reduced steadily over the period 2000–2019, from 736 000 in 2000 to 409 000 in 2019

(Table 3.1). The percentage of total malaria deaths among children aged under 5 years was 84% in 2000 and 67% in 2019. The estimate of deaths in 2015, the GTS baseline, was about 453 000. The malaria mortality rate (i.e. deaths per 100 000 population at risk) reduced from about 25 in 2000 to 12 in 2015 and 10 in 2019, with the slowing of the rate of decline in the latter years similar to that seen in number of cases (Fig. 3.2a).

**FIG. 3.2.**



**Global trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019, c) distribution of malaria cases and d) deaths by country, 2019** Source: WHO estimates.





Of the 87 countries that were malaria endemic in 2019, 29 accounted for 95% of malaria cases globally (**Fig. 3.2b**). Nigeria (27%), the Democratic Republic of the Congo (12%), Uganda (5%), Mozambique (4%) and Niger (3%) accounted for about 51% of all cases globally. About 95% of malaria deaths were in

32 countries (**Fig. 3.2c**). Nigeria (23%), the Democratic Republic of the Congo (11%), the United Republic of Tanzania (5%), Burkina Faso (4%), Mozambique (4%) and Niger (4%) accounted for about 51% of all malaria deaths globally in 2019 (**Fig. 3.2c**).



### 3 Global trends in the burden of malaria

#### 3.2 ESTIMATED MALARIA CASES AND DEATHS IN THE WHO AFRICAN REGION, 2000–2019

With an estimated 215 million malaria cases and 384 000 malaria deaths in 2019 (Table 3.2), the WHO African Region accounted for about 94% of cases and deaths globally. Although there were fewer malaria cases in 2000 (204 million) than in 2019, malaria case

incidence reduced from 363 to 225 cases per 1000 population at risk in this period (Fig. 3.3), reflecting the complexity of interpreting changing disease transmission in a rapidly increasing population. The population living in sub-Saharan Africa increased

**TABLE 3.2.**

**Estimated malaria cases and deaths in the WHO African Region, 2000–2019** Estimated cases and deaths are shown with 95% upper and lower confidence intervals. *Source: WHO estimates.*

Year	Number of cases (000)				Number of deaths		
	Point	Lower bound	Upper bound	% <i>P. vivax</i>	Point	Lower bound	Upper bound
2000	<b>204 000</b>	189 000	223 000	0.9%	<b>680 000</b>	657 000	713 000
2001	<b>210 000</b>	194 000	230 000	1.4%	<b>685 000</b>	662 000	720 000
2002	<b>207 000</b>	191 000	227 000	1.3%	<b>685 000</b>	661 000	721 000
2003	<b>211 000</b>	194 000	234 000	1.7%	<b>672 000</b>	644 000	717 000
2004	<b>214 000</b>	194 000	242 000	1.9%	<b>706 000</b>	671 000	771 000
2005	<b>211 000</b>	193 000	234 000	1.3%	<b>653 000</b>	624 000	703 000
2006	<b>211 000</b>	193 000	235 000	1.5%	<b>667 000</b>	637 000	713 000
2007	<b>211 000</b>	193 000	234 000	1.5%	<b>637 000</b>	610 000	678 000
2008	<b>211 000</b>	193 000	232 000	1.2%	<b>590 000</b>	567 000	625 000
2009	<b>215 000</b>	196 000	239 000	1.4%	<b>569 000</b>	538 000	618 000
2010	<b>215 000</b>	195 000	239 000	1.7%	<b>542 000</b>	509 000	597 000
2011	<b>211 000</b>	192 000	234 000	2.2%	<b>501 000</b>	474 000	544 000
2012	<b>209 000</b>	190 000	231 000	2.2%	<b>477 000</b>	449 000	522 000
2013	<b>205 000</b>	186 000	227 000	1.9%	<b>454 000</b>	424 000	500 000
2014	<b>197 000</b>	182 000	215 000	1.1%	<b>435 000</b>	414 000	469 000
2015	<b>199 000</b>	183 000	218 000	0.9%	<b>418 000</b>	397 000	453 000
2016	<b>205 000</b>	189 000	225 000	0.6%	<b>395 000</b>	376 000	430 000
2017	<b>212 000</b>	196 000	234 000	0.5%	<b>388 000</b>	369 000	428 000
2018	<b>212 000</b>	195 000	234 000	0.2%	<b>385 000</b>	367 000	429 000
2019	<b>215 000</b>	197 000	237 000	0.3%	<b>384 000</b>	365 000	433 000

*P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.




from about 665 million in 2000 to 1.1 billion in 2019 (Section 11). Malaria deaths in the WHO African Region reduced by 44%, from 680 000 in 2000 to 384 000 in 2019, and the malaria mortality rate reduced by 67% over the same period, from 121 to 40 per 100 000 population at risk (Fig. 3.3). Since 2014, however, the rate of progress in both cases and deaths

has slowed, attributed mainly to the stalling of progress in several countries with moderate or high transmission (Fig. 3.3). Distributions of malaria cases by country are shown in Fig. 3.3. It can be seen that 27 of the 29 countries that account for 95% of malaria cases globally (Fig. 3.2c) are in the WHO African Region.

**FIG. 3.3.**

**Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO African Region, 2019**  
 Source: WHO estimates.





### 3 Global trends in the burden of malaria

#### 3.3 ESTIMATED MALARIA CASES AND DEATHS IN THE WHO SOUTH-EAST ASIA REGION, 2000–2019

The WHO South-East Asia Region had nine malaria endemic countries in 2019, and contributed to about 3% of the burden of malaria cases globally. Malaria cases reduced by 74%, from 23.0 million in 2000 to about

6.3 million in 2019 (**Table 3.3**). India contributed to the largest absolute reductions, from about 20 million cases in 2000 to about 5.6 million in 2019. Malaria case incidence reduced by 78%, from about 18 to 4 per

**TABLE 3.3.**

**Estimated malaria cases and deaths in the WHO South-East Asia Region, 2000–2019** Estimated cases and deaths are shown with 95% upper and lower confidence intervals. *Source: WHO estimates.*

Year	Number of cases (000)				Number of deaths		
	Point	Lower bound	Upper bound	% <i>P. vivax</i>	Point	Lower bound	Upper bound
2000	<b>23 000</b>	18 700	29 100	47.8%	<b>35 000</b>	8 000	59 000
2001	<b>23 300</b>	19 100	29 200	50.6%	<b>34 000</b>	7 000	57 000
2002	<b>22 200</b>	17 900	28 000	50.0%	<b>33 000</b>	7 000	55 000
2003	<b>23 400</b>	18 900	29 300	52.4%	<b>33 000</b>	7 000	55 000
2004	<b>25 400</b>	20 200	32 400	52.0%	<b>36 000</b>	8 000	62 000
2005	<b>27 800</b>	21 600	36 700	53.8%	<b>39 000</b>	9 000	66 000
2006	<b>22 700</b>	17 500	30 400	51.5%	<b>33 000</b>	7 000	57 000
2007	<b>22 200</b>	17 100	30 300	49.6%	<b>33 000</b>	7 000	58 000
2008	<b>23 600</b>	18 000	32 200	47.5%	<b>36 000</b>	7 000	64 000
2009	<b>24 000</b>	18 100	33 500	45.3%	<b>38 000</b>	7 000	69 000
2010	<b>24 600</b>	19 400	33 100	46.0%	<b>38 000</b>	9 000	66 000
2011	<b>20 700</b>	16 200	27 900	47.7%	<b>31 000</b>	7 000	55 000
2012	<b>18 000</b>	14 200	24 000	47.6%	<b>27 000</b>	7 000	46 000
2013	<b>13 300</b>	10 500	17 400	46.2%	<b>21 000</b>	4 000	36 000
2014	<b>12 900</b>	10 100	17 300	35.2%	<b>23 000</b>	3 000	41 000
2015	<b>13 300</b>	10 400	17 700	34.4%	<b>24 000</b>	3 000	43 000
2016	<b>13 900</b>	10 400	19 500	34.9%	<b>25 000</b>	3 000	47 000
2017	<b>10 400</b>	7 800	14 100	37.3%	<b>18 000</b>	3 000	34 000
2018	<b>7 600</b>	5 500	10 300	50.5%	<b>11 000</b>	2 000	20 000
2019	<b>6 300</b>	4 500	8 600	51.7%	<b>9 000</b>	2 000	16 000

*P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.




1000 population at risk in the period 2000–2019 (Fig. 3.4). Malaria deaths reduced by 74%, from about 35 000 in 2000 to 9 000 in 2019. India accounted for 88% of malaria cases and 86% of malaria deaths in this

region in 2019. Sri Lanka was certified malaria free in 2015, and Timor-Leste reported zero malaria cases in 2018 and 2019.

**FIG. 3.4.**

**Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO South-East Asia Region, 2019** Source: WHO estimates.



### 3 Global trends in the burden of malaria

#### 3.4 ESTIMATED MALARIA CASES AND DEATHS IN THE WHO EASTERN MEDITERRANEAN REGION, 2000–2019

Malaria cases in the WHO Eastern Mediterranean Region reduced by 26%, from 7 million cases in 2000 to about 5 million in 2019 (Table 3.4). About a quarter of the cases in 2019 were due to *P. vivax*, mainly in

Pakistan and Afghanistan. Malaria deaths also reduced by 16%, from about 12 000 in 2000 to 10 100 in 2019. Over the period 2000–2019, malaria case incidence declined from 21 to 10 and mortality

**TABLE 3.4.**

**Estimated malaria cases and deaths in the WHO Eastern Mediterranean Region, 2000–2019**

Estimated cases and deaths are shown with 95% upper and lower confidence intervals. *Source: WHO estimates.*

Year	Number of cases (000)				Number of deaths		
	Point	Lower bound	Upper bound	% <i>P. vivax</i>	Point	Lower bound	Upper bound
2000	<b>7 000</b>	5 500	11 500	27.3%	<b>12 000</b>	4 000	22 000
2001	<b>7 200</b>	5 600	12 000	27.3%	<b>12 700</b>	4 200	22 500
2002	<b>6 800</b>	5 300	12 300	28.2%	<b>11 600</b>	4 400	20 000
2003	<b>6 400</b>	5 000	11 000	29.3%	<b>10 800</b>	3 800	18 600
2004	<b>5 300</b>	4 100	9 000	24.9%	<b>9 400</b>	2 800	16 300
2005	<b>5 500</b>	4 300	9 800	21.9%	<b>10 300</b>	3 200	17 800
2006	<b>5 500</b>	4 100	10 300	20.2%	<b>10 100</b>	3 300	17 400
2007	<b>4 800</b>	3 700	6 600	23.4%	<b>9 800</b>	3 600	17 000
2008	<b>3 700</b>	2 900	5 200	27.9%	<b>7 200</b>	2 500	12 300
2009	<b>3 600</b>	2 700	5 300	29.5%	<b>6 900</b>	2 500	12 200
2010	<b>4 500</b>	3 400	6 500	28.6%	<b>8 700</b>	3 500	14 800
2011	<b>4 600</b>	3 500	6 600	39.0%	<b>7 900</b>	3 200	12 800
2012	<b>4 300</b>	3 300	6 100	33.1%	<b>8 000</b>	3 000	12 900
2013	<b>4 000</b>	3 200	5 500	35.0%	<b>7 300</b>	2 800	11 700
2014	<b>4 200</b>	3 300	5 700	36.1%	<b>7 500</b>	2 800	12 200
2015	<b>4 100</b>	3 200	5 500	29.6%	<b>7 900</b>	2 600	13 100
2016	<b>5 200</b>	4 200	6 700	37.1%	<b>9 100</b>	3 400	15 000
2017	<b>5 000</b>	4 000	6 600	30.5%	<b>9 500</b>	3 200	16 500
2018	<b>5 400</b>	4 200	7 200	30.3%	<b>9 800</b>	3 100	17 600
2019	<b>5 200</b>	3 900	7 300	23.3%	<b>10 100</b>	2 900	19 000

*P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.




incidence rate from 4 to 2 (Fig. 3.5). Sudan is the leading contributor to malaria in this region, accounting for about 46% of cases (Fig. 3.5), followed by Yemen, Somalia, Pakistan, Afghanistan and Djibouti. Saudi Arabia reported only 38 indigenous malaria cases in

2019, and the Islamic Republic of Iran had no indigenous malaria cases in 2018 and 2019. Iraq, Oman and Syrian Arab Republic have last reported indigenous malaria cases in 2009, 2011 and 2004, respectively (Annex 3-F).

**FIG. 3.5.**

**Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO Eastern Mediterranean Region, 2019** Source: WHO estimates.



### 3 Global trends in the burden of malaria

#### 3.5 ESTIMATED MALARIA CASES AND DEATHS IN THE WHO WESTERN PACIFIC REGION, 2000–2019

The WHO Western Pacific Region had an estimated 1.7 million cases in 2019, a decrease of 43% from the 3 million cases in 2000 (Table 3.5). Malaria deaths reduced by 52%, from about 6600 cases in 2000 to 3200

in 2019. Over the same period, malaria case incidence reduced from 5 to 2 cases per 1000 population at risk (Fig. 3.6), and malaria mortality rate reduced from 1 to 0.4 deaths per 100 000 population at risk. Papua New

**TABLE 3.5.**

**Estimated malaria cases and deaths in the WHO Western Pacific Region, 2000–2019** Estimated cases and deaths are shown with 95% upper and lower confidence intervals. *Source: WHO estimates.*

Year	Number of cases (000)				Number of deaths		
	Point	Lower bound	Upper bound	% <i>P. vivax</i>	Point	Lower bound	Upper bound
2000	<b>2 990</b>	1 894	4 289	16.9%	<b>6 600</b>	2 200	11 800
2001	<b>2 631</b>	1 621	3 850	19.7%	<b>5 600</b>	1 800	10 300
2002	<b>2 334</b>	1 411	3 427	20.0%	<b>5 000</b>	1 600	9 300
2003	<b>2 526</b>	1 523	3 674	19.6%	<b>5 400</b>	1 700	10 000
2004	<b>2 936</b>	1 718	4 350	21.9%	<b>6 100</b>	1 800	11 700
2005	<b>2 509</b>	1 455	3 787	28.5%	<b>4 900</b>	1 500	9 500
2006	<b>2 659</b>	1 585	3 987	26.8%	<b>5 300</b>	1 600	9 800
2007	<b>2 018</b>	1 109	3 145	23.7%	<b>4 100</b>	1 100	8 400
2008	<b>1 845</b>	964	2 949	21.5%	<b>3 900</b>	900	7 900
2009	<b>2 436</b>	1 341	3 760	21.6%	<b>5 100</b>	900	10 200
2010	<b>1 839</b>	1 058	2 816	23.6%	<b>3 800</b>	800	7 500
2011	<b>1 576</b>	927	2 343	21.7%	<b>3 300</b>	600	6 700
2012	<b>1 888</b>	969	3 273	23.9%	<b>3 800</b>	700	8 800
2013	<b>1 964</b>	1 269	2 860	14.1%	<b>4 400</b>	600	8 800
2014	<b>2 321</b>	1 603	3 326	31.7%	<b>4 300</b>	700	8 200
2015	<b>1 431</b>	1 122	1 820	28.3%	<b>2 800</b>	500	4 800
2016	<b>1 676</b>	1 291	2 134	25.7%	<b>3 300</b>	500	6 000
2017	<b>1 961</b>	1 503	2 538	29.0%	<b>3 800</b>	600	6 700
2018	<b>1 981</b>	1 495	2 577	34.9%	<b>3 600</b>	500	6 600
2019	<b>1 739</b>	1 394	2 181	33.9%	<b>3 200</b>	500	5 600

*P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.




Guinea accounted for nearly 80% of all cases in this region in 2019. China has had no indigenous malaria cases since 2017. Malaysia had no cases of human malaria in 2018 and 2019, but reported 3212 cases of *P.*

*knowlesi*, considered to be zoonotic malaria, in 2019. Three countries had fewer than 10 000 cases in 2019: Republic of Korea (485), Vanuatu (1047) and Viet Nam (9702).

**FIG. 3.6.**

**Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO Western Pacific Region, 2019** Source: WHO estimates.



### 3.6 ESTIMATED MALARIA CASES AND DEATHS IN THE WHO REGION OF THE AMERICAS, 2000–2019

In the WHO Region of the Americas, malaria cases and case incidence reduced by 40% (from 1.5 million to 0.9 million) and 53% (from 14 to 6), respectively (**Table 3.6, Fig. 3.7**). Over the same period, malaria deaths and mortality rate reduced by 39% (from 909 to 551) and 50% (from 0.8 to 0.4), respectively. The

region's progress in recent years has suffered from the major increase in malaria in Venezuela (Bolivarian Republic of), which had about 35 500 cases in 2000, rising to over 467 000 by 2019. Brazil, Colombia and Venezuela (Bolivarian Republic of) account for 86% of all cases in this region.

**TABLE 3.6.**

**Estimated malaria cases and deaths in the WHO Region of the Americas, 2000–2019** Estimated cases and deaths are shown with 95% upper and lower confidence intervals. *Source: WHO estimates.*

Year	Number of cases (000)				Number of deaths		
	Point	Lower bound	Upper bound	% <i>P. vivax</i>	Point	Lower bound	Upper bound
2000	<b>1 540</b>	1 392	1 701	71.4%	<b>909</b>	666	1 168
2001	<b>1 297</b>	1 171	1 432	67.6%	<b>832</b>	593	1 090
2002	<b>1 183</b>	1 078	1 298	67.9%	<b>764</b>	514	1 030
2003	<b>1 159</b>	1 067	1 262	68.4%	<b>725</b>	480	992
2004	<b>1 146</b>	1 067	1 234	69.4%	<b>710</b>	460	986
2005	<b>1 283</b>	1 211	1 371	70.3%	<b>692</b>	443	968
2006	<b>1 106</b>	1 042	1 181	68.4%	<b>586</b>	348	852
2007	<b>994</b>	912	1 080	70.4%	<b>503</b>	297	744
2008	<b>699</b>	645	762	71.0%	<b>471</b>	224	756
2009	<b>687</b>	634	751	70.8%	<b>463</b>	227	740
2010	<b>821</b>	745	906	70.9%	<b>507</b>	250	791
2011	<b>611</b>	567	667	68.8%	<b>468</b>	206	733
2012	<b>580</b>	542	627	69.4%	<b>416</b>	211	622
2013	<b>562</b>	520	612	66.1%	<b>436</b>	227	642
2014	<b>477</b>	447	512	69.5%	<b>348</b>	196	484
2015	<b>561</b>	525	602	71.3%	<b>398</b>	216	551
2016	<b>677</b>	625	736	67.5%	<b>515</b>	252	731
2017	<b>915</b>	852	998	74.2%	<b>655</b>	287	947
2018	<b>926</b>	861	1 007	75.7%	<b>602</b>	243	880
2019	<b>889</b>	822	970	72.3%	<b>551</b>	220	813

*P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.




### 3.7 ESTIMATED MALARIA CASES AND DEATHS IN THE WHO EUROPEAN REGION, 2000–2019

Since 2015, the WHO European Region has been free of malaria. The last country to report an indigenous malaria case was Tajikistan in 2014. Also, no indigenous malaria deaths have been reported since 2000.

Throughout the period 2000–2019, no indigenous malaria deaths were reported in the WHO European Region.

**FIG. 3.7.**

Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO Region of the Americas, 2019 *Source: WHO estimates.*





### 3 Global trends in the burden of malaria


#### 3.8 CASES AND DEATHS AVERTED SINCE 2000, GLOBALLY AND BY WHO REGION

Cases and deaths averted in the period 2000–2019 were calculated by comparing the current annual estimated burden of malaria to a counterfactual that

was computed by holding the 2000 malaria case incidence and mortality rates constant throughout the period 2000–2019. The analysis shows that 1.5 billion

**FIG. 3.8.**

**Cumulative number of cases and deaths averted globally and by WHO region, 2000–2019** *Source: WHO estimates.*






malaria cases and 7.6 million malaria deaths have been averted globally in the period 2000–2019. Most of the cases (82%) and deaths (94%) averted were in the WHO African Region, followed by the South-East Asia Region (cases 10% and deaths 3%) (Fig. 3.8,

Fig. 3.9). In addition to malaria interventions, cases and deaths averted could also be due to other factors that modify malaria transmission or disease, such as improvements in socioeconomic status, malnutrition, infrastructure, housing and urbanization.

**FIG. 3.9.**

Percentage of a) cases and b) deaths averted by WHO region, 2000–2019 *Source: WHO estimates.*



AFR: WHO African Region; AMR: WHO Region of the Americas; EMR: WHO Eastern Mediterranean Region; EUR: WHO European Region; SEAR: WHO South-East Asia Region; WHO: World Health Organization; WPR: WHO Western Pacific Region.

### 3 Global trends in the burden of malaria

#### 3.9 BURDEN OF MALARIA IN PREGNANCY

Malaria infection during pregnancy has substantial risks for the pregnant woman, her fetus and the newborn child. For the pregnant woman, malaria infection can lead to severe disease and death, and placental sequestration of the parasite which can lead to maternal anaemia; it also puts the mother at increased risk of death before and after childbirth, and is an important contributor to stillbirth and preterm birth. Placental infection can also lead to poor fetal growth and low birthweight, which in turn can lead to child growth retardation and poor cognitive outcomes, as well as being a major risk factor for perinatal, neonatal and infant mortality (118–120).

To avert the consequences to women and children of malaria infections, WHO recommends – in combination with vector control, and prompt diagnosis and effective treatment of malaria – the use of IPTp with SP as part of


antenatal care (ANC) (Section 7.4) in areas of moderate to high transmission in sub-Saharan Africa.

The *World malaria report 2019* presented, for the first time, an analysis of the prevalence of malaria in pregnancy and the resulting burden of low birthweight (77). This section presents a follow-up analysis of the exposure to malaria infections during pregnancy and the prevalence of low birthweight. It also presents an analysis of the number of low birthweights averted if coverage of the first dose of IPTp was optimized, by ensuring that all women attending ANC clinics for the first time received the first dose.

The analysis in this section is restricted to moderate to high transmission countries in the WHO African Region (Annex 1), where the burden of malaria in pregnancy is most pronounced.

FIG. 3.10.

Estimated prevalence of exposure to malaria infection during pregnancy, overall and by subregion in 2019, in moderate to high transmission countries in the WHO African Region Source: Imperial College and WHO estimates.






### 3.9.1 Prevalence of exposure to malaria infections during pregnancy and contribution to low birthweight newborn

Malaria infection exposure during pregnancy (measured as cumulative prevalence over 40 weeks) was estimated from mathematical models (121) that relate estimates of the geographical distribution of *P. falciparum* exposure by age across Africa in 2019 with patterns of infections in placental histology by age and parity (122) (Annex 1). Country-specific age- and gravidity-specific fertility rates, stratified by urban or rural status, were obtained from DHS and malaria indicator surveys (MIS) (55), where such surveys had been carried out since 2014 and were available from the DHS programme website (56). For countries where

surveys were not available, fertility patterns were allocated based on survey data from a different country, matched on the basis of total fertility rate (123) and proximity. The exposure prevalence and the expected number of pregnant women who would have been exposed to infection were computed by country and subregion.

In 2019, in 33 moderate to high transmission countries<sup>1</sup> in the WHO African Region, there were an estimated 33.2 million pregnancies, of which 35% (11.6 million) were exposed to malaria infection (Fig. 3.10). By WHO subregion, Central Africa had the highest prevalence of exposure to malaria during pregnancy (40%) closely followed by West Africa (39%), while prevalence was 24% in East and Southern Africa.

<sup>1</sup> Angola, Benin, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad, Congo, Côte d'Ivoire, Democratic Republic of the Congo, Gabon, Gambia, Ghana, Guinea, Guinea-Bissau, Equatorial Guinea, Kenya, Liberia, Madagascar, Malawi, Mali, Mauritania, Mozambique, Niger, Nigeria, Senegal, Sierra Leone, South Sudan, Togo, Uganda, United Republic of Tanzania, Zambia and Zimbabwe.




### 3 Global trends in the burden of malaria

It is estimated that malaria infection during pregnancy in these 33 countries resulted in 822 000 children with low birthweight (**Table 3.8**) with almost half of these children (49%) being in the subregion of West Africa (**Table 3.8, Fig. 3.11**).

In the 33 countries, on average, 80% of all pregnant women visited ANC clinics at least once during their pregnancy, 62% received at least one dose of IPTp, 49% received at least two doses of IPTp and 34% received at least three doses of IPTp (**Section 7.4**). At current levels of IPTp coverage across all doses, an estimated 426 000 low birthweights were averted in 2019. If the

**FIG. 3.11.**

**Estimated number of low birthweights due to exposure to malaria infection during pregnancy, overall and by subregion in 2019, in moderate to high transmission countries in sub-Saharan Africa** Sources: Imperial College and WHO estimates.



WHO: World Health Organization.




80% of pregnant women visiting ANC clinics at least once during pregnancy received a single dose of IPTp, assuming they were all eligible, an additional 56 000 low birthweights would be averted, representing a significant missed opportunity under current levels of ANC use (Fig. 3.12). Urgent attention is clearly needed to optimize these missed opportunities while at the same

time ensuring high coverage of subsequent doses of IPTp. It is hoped that the recent call from the RBM Partnership to End Malaria to leaders and health policy-makers to increase protection of mothers and newborn children will result in an accelerated increase in IPTp coverage (124).

**FIG. 3.12.**

**Estimated number of low birthweights averted if current levels of IPTp coverage are maintained and the additional number averted if coverage of first dose of IPTp was optimized to match levels of coverage of first ANC visit in 2019, in moderate to high transmission countries in the WHO African Region** Sources: Imperial College and WHO estimates.



ANC: antenatal care; ANC1: first ANC visit; IPTp: intermittent preventive treatment in pregnancy; IPTp1: first dose of IPTp; WHO: World Health Organization.

# 4 Elimination

Globally, the number of countries that were malaria endemic in 2000 and that reported fewer than 10 000 malaria cases increased from 26 in 2000 to 46 in 2019. In the same period, the number of countries with fewer than 100 indigenous cases increased from 6 to 27. Between 2015 and 2019, the number of countries with fewer than 10 indigenous cases increased from 20 to 24 (Fig. 4.1).



## 4.1 MALARIA ELIMINATION CERTIFICATION

Between 2000 and 2019, 21 countries had achieved 3 consecutive years of zero indigenous malaria cases; 10 of these countries were certified malaria free by WHO (Table 4.1). Algeria is the first country in the WHO African

Region to be certified malaria free since 1973. The process to certify El Salvador is underway and would probably have been completed had the COVID-19 pandemic not disrupted the evaluation process.


## 4.2 E-2020 INITIATIVE

Progress in the reduction of malaria cases since 2010 in the 21 countries that are part of the E-2020 initiative is shown in Table 4.2. In the period 2010–2019, total malaria cases in the 21 countries reduced by 80%.

No indigenous cases were reported in Paraguay and Algeria, which were certified malaria free by WHO in 2018 and 2019, respectively.

**FIG. 4.1.**

**Number of countries that were malaria endemic in 2000, with fewer than 10, 100, 1000 and 10 000 indigenous malaria cases between 2000 and 2019** Sources: NMP reports and WHO estimates.





**TABLE 4.1.**

**Countries eliminating malaria since 2000** Countries are shown by the year that they attained 3 consecutive years of zero indigenous cases; countries that have been certified as malaria free are shown in green (with the year of certification in parentheses). *Sources: Country reports and WHO.*

2000	Egypt	United Arab Emirates (2007)		
2001				
2002				
2003				
2004	Kazakhstan			
2005				
2006				
2007	Morocco (2010)	Syrian Arab Republic	Turkmenistan (2010)	
2008	Armenia (2011)			
2009				
2010				
2011	Iraq			
2012	Georgia	Turkey		
2013	Argentina (2019)	Kyrgyzstan (2016)	Oman	Uzbekistan (2018)
2014	Paraguay (2018)			
2015	Azerbaijan	Sri Lanka (2016)		
2016	Algeria (2019)			
2017	Tajikistan			
2018				
2019	China	El Salvador		

WHO: World Health Organization.

Note: Although Maldives was certified in 2015, it was already malaria free before 2000.

**TABLE 4.2.**

**Number of indigenous malaria cases in E-2020 countries, 2010–2019** *Source: NMP reports.*

Country	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Algeria	1	1	55	8	0	0	0	0	0	0
Belize	150	72	33	20	19	9	4	7	3	0
Bhutan	436	194	82	15	19	34	15	11	6	2
Botswana	1 046	432	193	456	1 346	326	716	1 900	585	169
Cabo Verde	47	7	1	22	26	7	48	423	2	0
China	4 990	3 367	244	86	56	39	1	0	0	0
Comoros	36 538	24 856	49 840	53 156	2 203	1 300	1 066	2 274	15 613	17 599
Costa Rica	110	10	6	0	0	0	4	12	70	95
Ecuador	1 871	1 219	544	368	242	618	1 191	1 275	1 653	1 803
El Salvador	19	9	13	6	6	2	12	0	0	0
Eswatini	268	549	562	962	711	157	350	724	308	239
Iran (Islamic Republic of)	1 847	1 632	756	479	358	167	81	60	0	0
Malaysia	5 194	3 954	3 662	2 921	3 147	242	266	85	0	0
Mexico	1 226	1 124	833	495	656	517	551	736	803	618
Nepal	3 894	3 414	3 230	1 974	832	591	507	623	619	127
Paraguay	18	1	0	0	0	0	0	0	0	0
Republic of Korea	1 267	505	394	383	557	627	602	436	501	485
Saudi Arabia	29	69	82	34	30	83	272	177	61	38
South Africa	8 060	9 866	5 629	8 645	11 705	4 357	4 323	28 295	9 540	3 096
Suriname	1 771	771	356	729	401	81	76	40	29	95
Timor-Leste	48 137	19 739	5 208	1 025	347	80	81	16	0	0
<b>Total</b>	<b>116 859</b>	<b>71 790</b>	<b>71 668</b>	<b>71 776</b>	<b>22 661</b>	<b>9 237</b>	<b>10 166</b>	<b>37 094</b>	<b>29 793</b>	<b>24 366</b>

E-2020: eliminating countries for 2020; NMP: national malaria programme.



## 4 Elimination

China and El Salvador had no indigenous malaria cases for a third consecutive year and have made a formal request for certification. Iran (Islamic Republic of), Malaysia and Timor-Leste reported zero indigenous malaria cases in 2018 and 2019. In 2019, Belize and Cabo Verde reported zero indigenous malaria cases for the

first time since 2000. There were more cases in 2019 than in 2018 in Comoros, Costa Rica, Ecuador and Suriname, which reported 1986, 25, 150 and 66 additional cases, respectively. A malaria outbreak in Timor-Leste in 2020 is under investigation to determine whether any cases should be classified as indigenous.

### 4.3 THE GREATER MEKONG SUBREGION


The six countries of the GMS – Cambodia, China (Yunnan Province), Lao People’s Democratic Republic, Myanmar, Thailand and Viet Nam – have made huge gains against malaria as they aim for *P. falciparum* malaria elimination by 2025 (Fig. 4.2, Fig. 4.3) and elimination of all malaria by 2030. Between 2000 and 2019, the reported number of *P. falciparum* malaria cases fell by 97%, while all malaria cases fell by 90%. Of the 239 000 malaria cases

reported in 2019, 65 000 were *P. falciparum* cases. Overall, Cambodia (58%) and Myanmar (31%) accounted for most cases of malaria in the GMS.

The rate of decline has been fastest since 2012, when the MME programme was launched. During this period, malaria cases reduced sixfold, while *P. falciparum* cases reduced by a factor of nearly 14.

**FIG. 4.2.**

**Total malaria and *P. falciparum* cases in the GMS, 2000–2019** Sources: MME programme database and NMP reports.



GMS: Greater Mekong subregion; MME: Mekong Malaria Elimination; NMP: national malaria programme; *P. falciparum*: *Plasmodium falciparum*.



This accelerated decrease in *P. falciparum* is especially critical because of the increasing drug resistance; in the GMS, *P. falciparum* parasites have developed partial

resistance to artemisinin, the core compound of the best available antimalarial drugs.


#### 4.4 PREVENTION OF RE-ESTABLISHMENT

Once countries have eliminated malaria, re-establishment of transmission must be prevented through continued preventive measures in areas with malariogenic potential (risk of importation in areas receptive to transmission), vigilance to identify suspected malaria cases in the health system, quality-assured diagnosis and treatment, and follow-up to ensure complete cure and no onward transmission. After elimination, imported cases of malaria are expected, while any introduced or indigenous cases

signify local transmission and warn of deficiencies with prevention and surveillance strategies that must be addressed. Transmission of malaria may be considered re-established when at least three indigenous cases of malaria of the same species are found in the same transmission focus for 3 consecutive years. Between 2000 and 2019, no country that was certified malaria free has been found to have malaria transmission re-established.

**FIG. 4.3.**

**Regional map of malaria incidence in the GMS by area, 2012–2019** *Source: NMP reports.*



GMS: Greater Mekong subregion; NMP: national malaria programme.



# High burden to high impact approach

In November 2018, WHO and the RBM Partnership to End Malaria launched the high burden to high impact (HBHI) country-led approach (108), as a mechanism to support the 11 highest burden countries to get back on track to achieve the GTS 2025 milestones (4). The approach includes the four key response elements shown in Fig. 5.1. These 11 countries (Burkina Faso, Cameroon, the Democratic Republic of the Congo, Ghana, India, Mali, Mozambique, Niger, Nigeria, Uganda and the United Republic of Tanzania) account for 70% of the global estimated case burden and 71% of global estimated deaths from malaria. Several countries with a smaller population but with high malaria incidence have also adopted the HBHI approach.

Since November 2018, the HBHI response has been launched in 10 of the 11 countries (it has not yet been launched in Mali owing to disruptions due to the COVID-19 pandemic). However, all 11 countries have implemented HBHI-related activities across the four response elements. This section presents a summary of key activities and case studies for each of the first three response elements: political will, strategic information and better guidance.



## 5.1 GALVANIZING POLITICAL WILL, MOBILIZING RESOURCES AND MOBILIZING COMMUNITY RESPONSE

In each HBHI country initiation, there has been high-level political engagement and support. Several countries have begun adapting the RBM Partnership to End Malaria campaign, 'Zero Malaria Starts with Me' (125), through high-level national committees and councils, community mobilization and engagement activities, including the private sector.


Following the sixth replenishment of the Global Fund in October 2019, the global malaria allocation for the period 2020–2022 was US\$ 4.8 billion, an increase of about US\$ 1 billion from the previous allocation period. Of this, US\$ 2.1 billion was allocated to the 11 HBHI countries, an increase of about US\$ 500 million from the previous allocation in the period 2017–2019 (126).

In 2020, all HBHI countries except Mali submitted funding requests to the Global Fund, based on the analysis of subnational tailoring of interventions described in **Section 5.2**. At the same time, PMI increased its overall allocation to malaria in 2020 to about US\$ 770 million (from about US\$ 755 million in 2019), with most of the funds allocated to HBHI countries (52).

This section presents the Mass Action Against Malaria (MAAM) initiative in Uganda as an example of a country-led process of political engagement at all levels, and of multisectoral and community mobilization (**Box 5.1** on next page).

**FIG. 5.1.**

**HBHI: a targeted malaria response to get countries back on track to achieve the GTS 2025 milestones**  
 Sources: WHO GMP and RBM Partnership to End Malaria.



GMP: Global Malaria Programme; GTS: *Global technical strategy for malaria 2016–2030*; HBHI: high burden to high impact; WHO: World Health Organization.

**BOX 5.1.**

**Uganda's MAAM** Source: *Uganda NMP.*



**Background**

With a slogan of “Am I free of malaria today?” MAAM was launched in April 2018 in Kampala by the President of Uganda, His Excellency President Yoweri Museveni. To address the high malaria burden in the country, and its impact on individual and community development, MAAM was targeted at high-level state leadership, parliamentarians, government civil servants, religious and cultural leaders, media personnel, private sector, district health teams, health facilities, schoolteachers, community leaders, and households and the public at large. A handbook to guide MAAM has been developed, detailing the roles and responsibilities of all key stakeholders (127).



**Key stakeholders**

- The cabinet
- Parliamentarians
- Government ministries, parastatals and departments
- National and regional leaders (religious and cultural)
- Private sector
- Media
- Regional health directors and administrators
- District leaders
- Health care facility service providers
- Community leaders
- School administrators, teachers and other staff
- Households



**Expectations from stakeholders at all levels**

- To have a re-orientation of one's own values, to think about malaria prevention as a public health action to save lives
- To acknowledge that one's actions or inaction affects others
- To have full commitment to and accountability for the fight against malaria
- To support the scaling up of interventions against malaria
- To have a sense of urgency, acknowledging that each minute delayed or wasted costs lives, with negative consequences for the individual, the community and the economy



**Achievements**

- High-level launch and widespread media dissemination
- Development of MAAM handbook
- Incorporation of malaria agenda into the 2021–2025 National Development Plan III, Health Sector Development Plan III
- Establishment of Uganda Parliamentary Forum for Malaria (UPFM), supported by government
- Establishment of the UPFM scorecard for periodic rating of performance at constituency level
- Malaria agenda included in the political party manifesto for the 2021 national election
- Establishment of district task forces, and support for malaria operational interventions and local dissemination through music, dance and drama
- Increase in domestic malaria financing, through institutions such as the Ministry of Finance, Planning and Economic Development, with a budget call circular to all sectors to prioritize the malaria agenda
- Establishment of Malaria Free Uganda Initiative – a private mechanism to drive the malaria agenda
- Establishment of Rotary Malaria Partnership



**Challenges**

- Sustained funding for MAAM is required to ensure high impact
- Domestic financing is increasing but is not yet optimal
- Accountability at subnational level requires capacity-building
- Operationalization of initiatives is often delayed and slow paced



## 5.2 USING STRATEGIC INFORMATION TO DRIVE IMPACT

The HBHI Response Element 2 set out to implement work under five main areas in two phases (Table 5.1), with Phase 1 to be achieved by the end of 2020 and Phase 2 by the end of 2021. The HBHI countries have

successfully implemented all the Phase 1 activities, with support from a collaborative partnership coordinated by WHO.

**TABLE 5.1.**

**HBHI Response Element 2: work areas and status update** *Source: WHO.*

	Work area	Status
Phase 1 (by end of 2020)	<b>Progress review:</b> Country-level malaria situation analysis, and review of malaria programmes to understand progress and bottlenecks	Malaria programme reviews have been completed in all countries except Mali, where a review is in progress
	<b>Analysis of stratification, intervention mixes and prioritization:</b> Data analysis for stratification, optimal intervention mixes and prioritization for NMSP development and implementation	<p>Subnational tailoring of interventions has been completed in all countries except Mali, where tailoring is in progress. The example of Nigeria is shown in Fig. 5.3</p> <p>New NMSPs have been developed in all countries using analysis for subnational tailoring of interventions, and is in progress in Mali</p> <p>New NMSP have been used to develop funding requests to the Global Fund and other funders; these requests have been submitted for review and are in the grant-making process</p>
Phase 2 (by end of 2021)	<b>National malaria data repositories:</b> Functioning national malaria data repositories, with programme tracking dashboards	<p>WHO has developed a master indicators list for national integrated malaria databases</p> <p>WHO has developed a generic DHIS2 national repository platform that can be linked directly with HMIS DHIS2 instances</p> <p>An integrated malaria database repository has been launched in Nigeria, and repositories are under development in Ghana, Mozambique, Uganda and the United Republic of Tanzania</p> <p>Other countries have not yet started repositories, but discussions among countries and partners are ongoing</p>
	<b>Subnational operational plans:</b> Subnational operational plans linked to subnational health plans	<p>New NMSP have 5-year workplans</p> <p>Specific workplans will be developed once discussions with the Global Fund and other donors are complete</p> <p>WHO and partners will work with countries to develop annual workplans</p>
	<b>Monitoring and evaluation:</b> Ongoing national and subnational monitoring and evaluation of programmatic activities (including data systems) and impact	<p>Discussions are ongoing between WHO and each country and partners on enhanced monitoring and evaluation processes</p> <p>Learning from experience in Benin and Nigeria, countries will be encouraged to digitalize their ITN, IRS and SMC campaigns, to ensure efficient micro planning and distribution, with real-time data availability</p> <p>Comprehensive surveillance assessments are planned in Burkina Faso, the Democratic Republic of the Congo and Ghana; rapid surveillance system assessments are planned in other countries</p>

DHIS2: District Health Information Software 2; Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; HBHI: high burden to high impact; HMIS: health management information system; IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; NMSP: national malaria strategic plan; SMC: seasonal malaria chemoprevention; WHO: World Health Organization.

## 5 High burden to high impact approach

The process for analysing subnational tailoring of malaria interventions in the HBHI countries starts with the identification and mapping of operational units in each HBHI country. Demographic, epidemiological, entomological, climatic, health system and other contextual information is assembled for the operational units. Using flexible, context-specific criteria for the targeting of each WHO-approved intervention (4), countries then identify operational units that meet the criteria for each intervention. At the end of the process, each unit will have a mix of interventions tailored to its context. At various stages of the process, mathematical models are used to help countries understand the impact on malaria of the scenarios with different combinations of interventions. This information is then used to review and refine the goals of the NMSP, and to help with costing and prioritization of resources during funding requests to the government, the Global Fund, PMI and other donors. WHO coordinates this process; WHO also led the analysis support to countries and collaborated with several mapping and modelling groups to support the HBHI countries.<sup>1</sup>

The example of intervention mixes for each local government authority in Nigeria is presented in **Fig. 5.2**. This intervention-mix map was used to inform Nigeria's new NMSP, and funding requests to the Global Fund and PMI. It also helped with anticipating interventions that would be implemented if a joint malaria loan from the World Bank and Islamic Development Bank is approved, to target states that do not receive support from the Global Fund, PMI or other donors.

The main highlights of the analysis of subnational tailoring of intervention mixes in Nigeria were an increase in SMC-targeted states, from 114 to 395 local government authorities (LGAs), with actual planned implementation increasing from 114 to 305 LGAs based on available funding, increasing the number of children targeted for SMC from about 4 million to 16 million;

funding for new-generation piperonyl butoxide (PBO) nets to cover more than 160 million people; and a recognition that, before the next ITN campaign, a comprehensive exercise of urban microstratification to better target interventions and improve efficiencies will be implemented by the National Malaria Elimination Programme (NMEP), with support from WHO and partners, given that just over half of the 215 million people in Nigeria live in urban areas.




A modelling analysis of the impact of four intervention scenarios was implemented: business as usual (BAU), which is the pre-HBHI approach; a fully funded NMSP updated using the HBHI approach, where 80% or more of coverage of core interventions is achieved in areas where they are targeted; a funding request based on updated NMSPs that limits SMC to five states; and one that increases SMC to an additional five states (**Fig. 5.2**). The analysis shows that the BAU approach will lead to very small reductions in malaria prevalence in Nigeria, whereas full implementation of the subnationally tailored NMSP will lead to substantial reductions in malaria prevalence – by 2023, infection prevalence in children aged under 5 years will be about 16%, a reduction from the estimated prevalence of 28% in 2020. For the period 2020–2023, preliminary analysis by the NMEP of Nigeria shows that US\$ 2.75 billion is needed to achieve high coverage of interventions in targeted areas, and full availability of diagnosis and treatment in public health facilities. Additional funding is required to cover all SMC eligible populations as well as major improvements in treatment seeking behaviour, access to care, compliance with SMC and use of LLINs. Currently, available funding for LLINs, RDTs and ACTs for the period 2020–2023 is about US\$ 1.75 billion. If the current gap in funding is filled through to 2023, it is projected that, compared with BAU, about 73 million malaria cases and 66 000 deaths will be averted.

<sup>1</sup> The Swiss Tropical and Public Health Institute supported five countries (Cameroon, Ghana, Mozambique, Uganda and the United Republic of Tanzania), PATH supported three countries (the Democratic Republic of the Congo, Mali and Niger), and Northwestern University and the Institute for Disease Modeling supported two countries (Burkina Faso and Nigeria). Subnational maps of parasite prevalence and all-cause mortality in children aged under 5 years were received from the Malaria Atlas Project (MAP) and the Institute for Health Metrics and Evaluations, respectively.



**FIG. 5.2.**

**Example of subnational tailoring of malaria intervention mixes and their projected impacts implemented as part of the HBHI response (in Nigeria)** Sources: NMEP, WHO, Northwestern University, IDM.



Scenario	Cases and deaths averted compared to a business as usual scenario, 2020–2023			
	Cases: all ages	Cases: U5	Deaths: all ages	Deaths: U5
Full implementation of NMSP	103 000 000	32 000 000	90 000	75 000
Funded scenario with prioritized above allocation request (PAAR) SMC LGAs	73 000 000	24 000 000	66 000	54 000
Funded scenario without PAAR SMC LGAs	71 000 000	23 000 000	64 000	53 000

BAU: business as usual; HBHI: high burden to high impact; IDM: Institute for Disease Modeling; IPTi: intermittent preventive treatment in infants; IPTp: intermittent preventive treatment in pregnancy; LGA: local government authority; LLIN: long-lasting insecticidal net; NMEP: National Malaria Elimination Programme; NMSP: national malaria strategic plan; PAAR: prioritized above allocation request; PBO: piperonyl butoxide; PPR: *Plasmodium falciparum* parasite rate; SMC: seasonal malaria chemoprevention; U5: aged under 5 years; WHO: World Health Organization.



## 5 High burden to high impact approach

### 5.3 IMPROVING WHO'S MALARIA POLICY-MAKING AND DISSEMINATION PROCESSES

Before the launch of HBHI, the GMP had already begun an extensive review of the WHO process for developing and disseminating policy guidance on malaria (128). The overall aim was to improve the transparency, consistency, efficiency and predictability of the policy-making process, to make policies timely and more readily adaptable by countries. The resulting pathway was structured as a three-tier process: better anticipate, develop policy and optimize uptake. The HBHI response has added further urgency to this process (128).

In 2019, WHO created a compendium of its malaria guidance (109), to list all WHO recommendations and associated guidance on malaria in a single resource, and to inform programme managers, and national and international stakeholders. The compendium also references relevant WHO handbooks, manuals and

other resources, to guide readers on how these global recommendations can best be implemented. In the same year, WHO updated its technical brief to countries, to support them in the development of robust funding proposals that are tailored to their context (110). The document provided information on the process of stratification, which guides tailoring of interventions to the local context and the prioritization of resources, while adhering to the evidence-based recommendations that have been developed through WHO's standard, stringent processes.

Based on the new WHO policy pathway, in 2020, the GMP established several guideline development groups focusing on vector control, case management, chemoprevention and elimination strategies. The results from the deliberations of these groups are expected in early 2021.

### 5.4 COORDINATED RESPONSE

Several areas of focus were identified for the HBHI fourth response element: stakeholder landscaping; identification of in-country processes requiring coordination; strengthening coordination structures; and aligning partner support around the national strategic and implementation plans.

Although countries have undertaken some assessment of the status of coordination during the initiation phase, most have not formally evaluated their needs. Early feedback from some countries shows that, although they are grateful for the support they receive from partners and WHO, gaps remain; for example:

- weak NMP organizational and staff capacities;
- weak cross-partner coordination structures;
- weak subnational coordination structures;
- potential risks of partner misalignment with NMSPs and operational research priorities;
- issues around the sustainability of project-driven interventions and activities;
- challenges of complex emergencies, including the COVID-19 pandemic.

### 5.5 MALARIA IN HBHI COUNTRIES SINCE 2018

Comparisons of malaria cases, case incidence, deaths and mortality rates in 2018 (the year HBHI was launched) and 2019 are presented in **Fig. 5.3**. Overall, there have been no major changes in the burden of malaria in these countries since 2018. Although cases in India reduced by 1.2 million and Mali by 0.8 million, increases were estimated in Nigeria (2.4 million) and

the Democratic Republic of the Congo (1.2 million). Overall, cases increased slightly from 155 million to 156 million between the two years, and deaths reduced from 263 000 to 226 000. In 2015, at GTS baseline, there were 148 million estimated malaria cases in the 11 HBHI countries.

### 5.6 REPORTED MALARIA CASES IN HBHI COUNTRIES SINCE 2018 AND COMPARISONS WITH ESTIMATED CASES


The methods used in this report to estimate the burden of malaria cases and deaths have several limitations. These methods are elaborated in **Annex 1**. The implications of the limitations become stark in the HBHI countries because they account for 70% or more of the burden of malaria morbidity and mortality. In moderate to high malaria transmission countries in Africa, including the HBHI countries in this region,

WHO uses a parasite rate-to-incidence model to estimate malaria cases (**Annex 1, Section 3**). The process of estimation relies on community parasite surveys, interventions and climatic data to quantify parasite prevalence, which is then transformed to incidence using epidemiological methods (93). The estimates are often different from cases reported by countries, and this difference has been an important



**FIG. 5.3.**

**Estimated malaria a) cases, b) cases per 1000 population at risk, c) deaths and d) deaths per 100 000 population at risk, 2018 and 2019, in HBHI countries** *Source: WHO estimates.*



HBHI: high burden to high impact; WHO: World Health Organization.

## 5 High burden to high impact approach

source of unease among NMPs. Another method used for Southern African countries and those outside Africa where malaria transmission is low is the adjustment of reported data, mainly from the public health sector, for reporting, testing and treatment seeking rates (**Annex 1**).

**Table 5.2** compares the results of two methods used to estimate burden: the parasite-rate-to-incidence method (107) used by WHO and the approach based on adjustment of routine data. The WHO method estimates about 150 million cases in 2019 but the

method based on adjustment of routine data estimates about 265 million cases. Previous analysis showed that similar differences (i.e. with the routine data method generally resulting in more cases) are seen in most of the 20 sub-Saharan countries that use the parasite rate-to-incidence method. These discrepancies could be explained by data quality, epidemiological and methodological issues (129). However, improving national data systems (e.g. in terms of granularity, frequency and quality) is the clear path towards a better understanding of the malaria burden.

**TABLE 5.2.**

**Comparisons of estimated malaria cases (millions) using the parasite rate-to-incidence model (Annex 1) and the reported data from the routine public health sector in high burden countries of the WHO African Region, 2019** Sources: WHO estimates and NMP reports.

Country	Estimated cases using parasite rate-to-incidence model (population-wide estimate)	Reported cases from the routine system (public health sector)	Reported cases adjusted for reporting and testing rates (public health sector)	Reported cases adjusted for reporting and testing rates and treatment seeking (population-wide estimate)	Population at risk 2019
Burkina Faso <sup>a</sup>	7.9	10.3	12.2	14.9	20.3
Cameroon	6.3	1.2	1.5	6.3	25.9
Democratic Republic of the Congo	28.3	20.5	21.6	80.9	86.8
Ghana	4.9	5.0	5.6	14.0	30.4
Mali	6.6	2.7	3.1	9.0	19.7
Mozambique	9.4	11.7	12.5	15.7	30.4
Niger	8.0	3.7	3.9	6.3	23.3
Nigeria	61.0	17.8	22.8	72.0	201.0
Uganda	11.6	12.3	14.0	33.8	44.3
United Republic of Tanzania	6.5	5.9	6.3	11.8	58.0
<b>Total</b>	<b>150.3</b>	<b>91.0</b>	<b>103.5</b>	<b>264.8</b>	<b>540.0</b>

<sup>a</sup> For Burkina Faso, monthly data from 2018 was used due to major disruptions of the surveillance system due to the 2019 health workers' strikes in 2019.





# Investments in malaria programmes and research

The GTS sets out estimates of the funding required to achieve milestones for 2020, 2025 and 2030. Total annual resources needed were estimated at US\$ 4.1 billion in 2016, rising to US\$ 6.8 billion in 2020. An additional US\$ 0.72 billion is estimated to be required annually for global malaria research and development (R&D) (4). **Section 6.1** presents the most up-to-date funding trends for malaria control and elimination (by source and channel of funding) for the period 2000–2019, where permitted through available data, both globally and for major country groupings. **Section 6.2** presents investments in malaria-related R&D for the period 2007–2018.



## 6.1 FUNDING TRENDS FOR MALARIA CONTROL AND ELIMINATION

Malaria-related annual funding from donors through multilateral agencies was estimated from donors' contributions to the Global Fund from 2010 through 2019. Organisation for Economic Co-operation and Development (OECD) contributions were available from 2011 through 2018, with 2010 estimates using 2011 data and 2019 estimates using 2018 data. In addition, contributions from malaria endemic countries to multilateral agencies were allocated to governments of endemic countries for the years 2010 through 2019.

For the 91 countries analysed in this section, total funding for malaria control and elimination in 2019 was estimated at US\$ 3.0 billion, compared with US\$ 2.7 billion in 2018 and US\$ 3.2 billion in 2017. The amount invested in 2019 falls short of the US\$ 5.6 billion estimated to be required globally to stay on track towards the GTS milestones (4). Moreover, the funding gap between the amount invested and the resources needed has continued to widen significantly over recent years, increasing from US\$ 1.3 billion in 2017 to US\$ 2.3 billion in 2018, and to US\$ 2.6 billion in 2019. Over the period 2010–2019, international sources provided 70% of the total funding for malaria control and elimination, led by the US, the United Kingdom of Great Britain and Northern Ireland (United Kingdom) and France over this period (**Fig. 6.1**). Of the US\$ 3.0 billion invested in 2019, US\$ 2.1 billion came from international funders. The highest contributions in 2019 were from the government of the United States of


America (USA), which provided a total of US\$ 1.1 billion through planned bilateral funding and contributions to multilateral funding agencies. This was followed by bilateral and multilateral disbursements from the United Kingdom of US\$ 0.2 billion; contributions of over US\$ 0.1 billion from each of France, Germany and Japan totalling US\$ 0.4 billion; and a combined US\$ 0.4 billion from other countries that are members of the Development Assistance Committee and from private sector contributors (**Fig. 6.2**). Governments of malaria endemic countries contributed 31% of the total funding (**Fig. 6.1**), with investments nearing US\$ 0.9 billion in 2019 (**Fig. 6.2**). Of this amount, an estimated US\$ 0.2 billion was spent on malaria case management in the public sector and US\$ 0.7 billion on other malaria control activities.

To analyse malaria investment since 2000, international bilateral funding data were obtained from several sources, with the historical availability varying across donors. From the USA, data on total annual planned funding from the Centers for Disease Control and Prevention (CDC), Department of Defense and USAID are available from 2001 through 2019. Total annual planned funding for USAID was utilized from 2001 through 2005, until the introduction of country-specific funding in 2006. The country recipient for funding has been labelled as “unspecified” for all years where country-specific data are not available.



**FIG. 6.1.**


**Funding for malaria control and elimination, 2010–2019 (% of total funding), by source of funds (constant 2019 US\$)** Sources: ForeignAssistance.gov, Global Fund, NMP reports, OECD CRS database, United Kingdom Department for International Development, WHO estimates and World Bank DataBank.



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; NMP: national malaria programme; OECD: Organisation for Economic Co-operation and Development; United Kingdom: United Kingdom of Great Britain and Northern Ireland; WHO: World Health Organization.

**FIG. 6.2.**

**Funding for malaria control and elimination, 2010–2019, by source of funds (constant 2019 US\$)** Sources: ForeignAssistance.gov, United Kingdom Department for International Development, Global Fund, NMP reports, OECD CRS database, the World Bank Data Bank and WHO estimates.



CRS: creditor reporting system; Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; NMP: national malaria programme; OECD: Organisation for Economic Co-operation and Development; United Kingdom: United Kingdom of Great Britain and Northern Ireland; WHO: World Health Organization.

## 6 Investments in malaria programmes and research

Data on annual disbursements by the Global Fund to malaria endemic countries are available from 2003 through 2019. For the government of the United Kingdom, funding data towards malaria control are available from 2007 through 2019: for the years 2007 through 2016, disbursement data were obtained through the OECD creditor reporting system (CRS) on aid activity; for 2017 through 2019, disbursement data were sourced from *Statistics on International Development: final UK aid spend 2019 (130)*. For all other donors, disbursement data were also obtained from the OECD CRS database for the period 2002–2018. For years with no data available for a particular funder, no imputation was conducted; hence, the trends presented throughout **Figs 6.3–6.5** should be interpreted carefully.

Contributions from governments of endemic countries were estimated as the sum of contributions reported by NMPs for the world malaria report of the relevant year plus the estimated costs of patient care delivery services at public health facilities. From 2000 to 2019, where available, government expenditures were used for their contributions (if unavailable, then government budgets or estimates were used), whereas patient care delivery costs were estimated using unit cost estimates from WHO-CHOosing Interventions that are Cost-Effective (WHO-CHOICE) 2010, with values included for the years 2010 through 2019.

Of the US\$ 3.0 billion invested in 2019, nearly US\$ 1.2 billion (39%) was channelled through the Global Fund (**Fig. 6.4**). Compared with 2018, the Global Fund's disbursements to malaria endemic countries increased by about US\$ 0.2 billion in 2019. This difference reflects the cyclical distribution of ITNs supported by the Global Fund combined with an increase in disbursements in 2019, which corresponded to the end of most malaria grants in that year (**Fig. 6.4**).


Planned bilateral funding from the government of the USA amounted to US\$ 0.8 billion in 2019, which matched the levels of funding in 2017 and 2018, but is higher than the levels of all other annual planned contributions from 2001, when data first became available, to 2016 (**Fig. 6.3**). The United Kingdom remains the second largest bilateral funder, with less than US\$ 0.1 billion in 2019, followed by the World Bank and other Development Assistance Committee members (**Fig. 6.3**). The total contribution from governments of malaria endemic countries remained constant, at US\$ 0.9 billion invested, in both 2018 and 2019.

**Fig. 6.3** shows the substantial variation across country income groups in the share of funding received from domestic and international sources. The 27 low-income countries accounted for 41% of total malaria funding in 2019, down from 47% in 2018 (corresponding to >90% of global malaria cases and deaths), with 84% of their funding coming from international sources. International funding also dominated in the group of 37 lower-middle-income countries (48% of total funding in 2019), accounting for 69% of the amount invested in these countries. In contrast, in the group of 20 upper-middle-income countries (10% of the total funding in 2019), 13% of their malaria funding came from international sources, and 87% from domestic public funding. Lastly, the three high-income countries accounted for 1% of total malaria funding, with 100% of their funding coming from domestic sources.

Of the US\$ 3.0 billion invested in 2019, 73% benefited the WHO African Region, 9% went to the WHO South-East Asia Region, 5% each to the WHO Region of the Americas and the WHO Western Pacific Region, and 4% to the WHO Eastern Mediterranean Region (**Fig. 6.5**). Funding flows for which no geographical information on recipients was available represented 4% of the total funding in 2019 (**Fig. 6.5**).

**FIG. 6.3.**


**Funding for malaria control and elimination, 2000–2019, by World Bank 2019 income group and source of funding (constant 2019 US\$)<sup>a</sup>** Sources: *ForeignAssistance.gov, Global Fund, NMP reports, OECD creditor reporting system database, United Kingdom Department for International Development, WHO estimates and World Bank DataBank.*





**FIG. 6.4.**


**Funding for malaria control and elimination, 2000–2019, by channel (constant 2019 US\$)** Sources: ForeignAssistance.gov, Global Fund, NMP reports, OECD creditor reporting system database, United Kingdom Department for International Development, WHO estimates and World Bank DataBank.



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; NMP: national malaria programme; OECD: Organisation for Economic Co-operation and Development; United Kingdom: United Kingdom of Great Britain and Northern Ireland; USA: United States of America; WHO: World Health Organization.

**FIG. 6.5.**

**Funding for malaria control and elimination, 2000–2019, by WHO region (constant 2019 US\$)<sup>a</sup>** Sources: ForeignAssistance.gov, United Kingdom Department for International Development, Global Fund, NMP reports, OECD creditor reporting system database, World Bank Data Bank and WHO estimates.



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; NMP: national malaria programme; OECD: Organisation for Economic Co-operation and Development; United Kingdom: United Kingdom of Great Britain and Northern Ireland; WHO: World Health Organization.

<sup>a</sup> "Unspecified" category refers to funding flows, with no information on the geographical localization of their recipients.



# 6 Investments in malaria programmes and research

## 6.2 INVESTMENTS IN MALARIA-RELATED R&D

### 6.2.1 Overarching trends

Between 2007 and 2018, almost US\$ 7.3 billion was invested in basic research and product development for malaria. The malaria R&D funding landscape has been led by investment in drugs (US\$ 2.6 billion, 36% of malaria funding between 2007 and 2018), followed by relatively similar shares for basic research (US\$ 1.9 billion, 26%) and vaccines R&D (US\$ 1.8 billion, 25%). Investments in vector control products and diagnostics were notably lower, reaching overall totals of US\$ 453 million (6.2%) and US\$ 185 million (2.5%), respectively (Fig. 6.6).

Changes in total malaria funding have largely reflected the progression of the overall pipeline. For example, a spike in vaccine funding in 2008–2009 – related to a surge of funding for Phase III trials of the RTS,S malaria vaccine candidate – was followed by a sharp drop and some subsequent stagnation in malaria R&D funding between 2010 and 2015. Driven in part by increased public sector investments in discovery and preclinical


R&D for drugs and vaccines, as well as increased industry investment in several Phase II trials of new chemical entities with potential for single-exposure radical cure, overall funding has climbed again since 2016, steadily returning to near-peak levels in 2018.

Between 2007 and 2018, the public sector held a leading role in malaria R&D funding, growing from US\$ 246 million in 2007 to a peak of US\$ 365 million in 2017. Within the public sector and among all malaria R&D funders, the US National Institutes of Health was the largest contributor, focusing just over half of its US\$ 1.9 billion investment into basic research (US\$ 1.02 billion, 54% of their overall malaria investment between 2007 and 2018).

The Bill & Melinda Gates Foundation has been another instrumental player, investing US\$ 1.8 billion (25% of all malaria R&D funding) between 2007 and 2018, and supporting the clinical development of key innovations such as the RTS,S vaccine. The Bill & Melinda Gates Foundation has given more funding to malaria than

**FIG. 6.6.**

**Funding for malaria-related R&D, 2007–2018, by product type (constant 2019 US\$)<sup>a</sup>** Sources: Policy Cures Research G-FINDER data portal (104).



R&D: research and development.

<sup>a</sup> "Unspecified" category refers to funding flows, with no information on the geographical localization of their recipients.



any other disease-specific investment reported by G-FINDER.

The industry sector has played a prominent role in advancing malaria drug development. From an overall investment of US\$ 1.4 billion between 2007 and 2018, most of the funding (US\$ 932 million, 68%) went towards drug R&D. Overall industry investment has increased in recent years, related mainly to an expanded focus on clinical development as drug candidates advanced through clinical trials from 2015 onwards. This change in focus, combined with declines in philanthropic funding during the same period, led to funding from industry surpassing philanthropic funding in 2017 for the first time in the past decade.

### 6.2.2 Funding flows


Two thirds (US\$ 4.9 billion, 67%) of all funding for malaria basic research and product development between 2007 and 2018 was given externally in the form of grants or contracts, with internal investments (US\$ 2.4 billion, 33%) making up the remainder (Fig. 6.7). Academic and nongovernment research

institutes received the largest share of direct, external funding (US\$ 2.4 billion, 49%), 54% (US\$ 1.3 billion) of which went to basic research between 2007 and 2018. Most internal investment, on the other hand, was accounted for by industry (US\$ 1.4 billion, 58%), followed by the public sector (US\$ 972 million, 40%). About 74% (US\$ 722 million) of the public sector funds came from intramural funding by the US Department of Defense and US National Institutes of Health.

Product development partnerships and other intermediaries received US\$ 1.7 billion (23%) of overall external malaria R&D funding, which was used primarily for investment in drugs (US\$ 867 million, 51% of their overall funding) and vaccines (US\$ 522 million, 31%). During this period, multiple product development partnerships – including PATH’s Malaria Vaccine Initiative (MVI), MMV, FIND and IVCC – have worked to advance development of key malaria product innovations, including numerous drugs, next-generation vector control tools, and, of course, the world’s first malaria vaccine to provide partial protection against malaria in young children.

**FIG. 6.7.**

**Malaria R&D funding from 2007 to 2018, by sector (constant 2019 US\$)** Source: Policy Cures Research, G-FINDER data portal (104).



R&D: research and development.



# Distribution and coverage of malaria prevention, diagnosis and treatment

WHO recommends several interventions for the prevention, diagnosis and treatment of malaria (106). The prevention interventions tracked in this report are ITNs, indoor residual spraying (IRS), SMC and IPTp, discussed here in **Sections 7.1–7.4**. To measure progress in access to prompt case management, **Section 7.5** presents the latest results on distribution of RDTs and ACTs, and population-level coverage of malaria diagnosis and treatment.



## 7.1 DISTRIBUTION AND COVERAGE OF ITNs

Manufacturers delivered about 253 million ITNs to malaria endemic countries in 2019, an increase of 56 million ITNs compared with 2018 (**Fig. 7.1**). About 84% of these ITNs were delivered to countries in sub-Saharan Africa. About 46% of the ITNs delivered by manufacturers were received in Nigeria (33.4 million), the Democratic Republic of the Congo (28.0 million), Ethiopia (15.1 million), Mali (10.4 million), Mozambique (10.2 million), Sudan (10.1 million) and Benin (9.7 million). Data from 2010–2019 are presented here; however, manufacturers' delivery data for 2004–2019 show that nearly 2.2 billion ITNs were supplied globally in that period, of which 1.9 billion (86%) were supplied to sub-Saharan Africa.

In 2019, 154 million ITNs were distributed globally by NMPs in malaria endemic countries. Of these ITNs, 140 million were distributed in sub-Saharan Africa, with a combined total of about 103 million ITNs being distributed in seven countries: Nigeria (31 million), the Democratic Republic of the Congo (21 million), Ethiopia (11 million), Guinea (9 million), Senegal (9 million), Burundi (8 million) and Cameroon (8 million). Outside of sub-Saharan Africa, the largest distribution was in Myanmar (11 million).

Indicators of population-level coverage of ITNs were estimated for sub-Saharan African countries in which ITNs are the main method of vector control. Household




surveys were used, together with manufacturer deliveries and NMP distributions, to estimate the following main indicators:

- ITN use (i.e. percentage of a given population group that slept under an ITN the night before the survey);
- ITN ownership (i.e. percentage of households that owned at least one ITN);
- percentage of households with at least one ITN for every two people; and
- percentage of the population with access to an ITN within their household (i.e. percentage of the population that could be protected by an ITN, if each ITN in a household could be used by two people).

**FIG. 7.1.**

**Number of ITNs delivered by manufacturers and distributed<sup>a</sup> by NMPs, 2010–2019** Sources: Milliner Global Associates and NMP reports.



ITN: insecticide-treated mosquito net; NMP: national malaria programme.

<sup>a</sup> A lag between manufacturer deliveries to countries and NMP distributions of about 6–12 months is expected; thus, deliveries by manufacturers in a given year are often not reflected in distributions by NMPs in that year. Also, distributions of ITNs reported by NMPs do not always reflect all the nets that have been distributed to communities, depending on completeness of reporting. These issues should be considered when interpreting the relationship between manufacturer deliveries, NMP distributions and likely population coverage. Additional considerations include nets that are in storage in country but have not yet been distributed by NMPs, and those sold through the private sector that are not reported by programmes.

# 7 Distribution and coverage of malaria prevention, diagnosis and treatment


By 2019, 68% of households in sub-Saharan Africa had at least one ITN, increasing from about 5% in 2000. The percentage of households owning at least one ITN for every two people increased from 1% in 2000 to 36% in 2019. In the same period, the percentage of the population with access to an ITN within their household increased from 3% to 52%. The percentage of the population sleeping under an ITN also increased considerably between 2000 and 2019, for the whole

population (from 2% to 46%), for children aged under 5 years (from 3% to 52%) and for pregnant women (from 3% to 52%). These indicators represent impressive progress since 2000, although coverage peaked in 2017 (**Fig. 7.2**).

Using concentration indices, socioeconomic equity of ITN use by the children aged under 5 years at the subnational level was analysed. The most recent

**FIG. 7.2.**

**Indicators of population-level coverage of ITNs, sub-Saharan Africa, 2000–2019: a) percentage of households with at least one ITN, b) percentage of households with one ITN for every two people, c) percentage of population with access to an ITN, d) percentage of population using an ITN, e) percentage of children aged under 5 years using an ITN and f) percentage of pregnant women sleeping under an ITN**  
*Sources: ITN coverage model from MAP (131).*



CI: confidence interval; ITN: insecticide-treated mosquito net; MAP: Malaria Atlas Project.




household survey data from DHS and MIS from 24 countries<sup>1</sup> for 2015–2019 were used (Fig. 7.3). In most West African countries, ITN use was generally pro-poor (i.e. concentration index <0) (Fig. 7.3). The concentration index varies from -1 to +1, with a value of zero indicating perfect equality. In this analysis, negative and positive

values suggest that ITN use is concentrated in the poorest and richest households. In contrast, ITN use was higher in wealthier households (i.e. concentration index >0) in many parts of the Democratic Republic of the Congo, Kenya, Mozambique, Uganda and the United Republic of Tanzania.

<sup>1</sup> Angola (DHS 2018), Benin (DHS 2017–2018), Burkina Faso (MIS 2017–2018), Burundi (DHS 2016–2017), Cameroon (DHS 2018), Ethiopia (DHS 2016), Ghana (MIS 2019), Guinea (DHS 2018), Kenya (MIS 2015), Liberia (MIS 2016), Madagascar (MIS 2016), Malawi (MIS 2017), Mali (DHS 2018), Mozambique (MIS 2018), Nigeria (DHS 2018), Rwanda (MIS 2017), Senegal (DHS 2018), Sierra Leone (MIS 2016), Togo (MIS 2017), Uganda (MIS 2018–2019), United Republic of Tanzania (MIS 2017), Zambia (DHS 2018) and Zimbabwe (DHS 2015).

**FIG. 7.3.**

**Concentration index of ITN use by children aged under 5 years, sub-Saharan Africa at administrative level 1** Source: Most recent household surveys from the period 2015–2019. Kenya Medical Research Institute – Wellcome Trust Research Programme.




### 7.2 POPULATION PROTECTED WITH IRS

Globally, the percentage of the populations at risk protected by IRS in malaria endemic countries declined from 5% in 2010 to 2% in 2019. The percentage of the population protected by IRS decreased in all WHO regions (Fig. 7.4). The number of people protected

globally fell from 180 million in 2010 to 115 million in 2015, but declined to 97 million in 2019. By country, Burundi, Ethiopia, India and Somalia each had the number of people protected with IRS reducing by a million or more when 2019 was compared with 2018.

**FIG. 7.4.**


**Percentage of the population at risk protected by IRS, by WHO region, 2010–2019** Source: IVCC data and NMP reports.



AFR: WHO African Region; AMR: WHO Region of the Americas; EMR: WHO Eastern Mediterranean Region; IRS: indoor residual spraying; IVCC: Innovative Vector Control Consortium; NMP: national malaria programme; SEAR: WHO South-East Asia Region; WHO: World Health Organization; WPR: Western Pacific Region.

**FIG. 7.5.**

**Subnational areas where SMC was delivered in implementing countries in sub-Saharan Africa, 2019** Source: LSHTM.





## 7.3 SCALE-UP OF SMC

In Benin, SMC was scaled up for the first time, taking the number of countries in the Sahel that implement SMC to 13. The number of children reached with at least one dose of SMC steadily increased, from about 0.2 million in 2012 to about 21.5 million in 2019 (Table 7.1). Subnational areas in each country where

SMC was targeted in 2019 are shown in Fig. 7.5. In the 13 countries, a total of about 21.7 million children were targeted in 2019. On average, 21.5 million children received treatment each month (Table 7.2), but household surveys are needed to establish coverage gaps.

**TABLE 7.1.**

**Average number of children treated with at least one dose of SMC by year in countries implementing SMC, 2012–2019** Sources: NMPs, LSHTM and MMV.

Country	2012	2013	2014	2015	2016	2017	2018	2019
Benin	0	0	0	0	0	0	0	114 165
Burkina Faso	0	0	307 770	860 058	2 648 083	2 949 901	3 298 397	3 298 397
Cameroon	0	0	0	0	1 070 865	1 581 183	1 636 658	1 681 737
Chad	10 000	263 972	27 307	322 493	824 806	899 320	1 184 706	1 491 905
Gambia	0	0	48 953	76 450	73 710	76 726	101 511	110 870
Ghana	0	0	0	115 309	151 510	327 446	329 953	964 956
Guinea	0	0	0	201 283	442 177	575 927	840 120	750 903
Guinea-Bissau	0	0	0	1 999 987	36 681	166 162	90 998	82 918
Mali	160 000	537 294	699 880	646 173	3 849 672	3 990 096	4 278 401	3 767 205
Niger	0	225 970	518 110	787 399	1 994 345	2 545 885	3 810 884	4 151 103
Nigeria	0	209 451	370 280	471 803	1 579 229	2 284 915	3 460 733	4 110 152
Senegal	0	55 709	446 809	0	477 614	485 717	0	671 132
Togo	0	119 222	127 624	5 480 954	308 858	382 319	325 621	296 332
<b>Total</b>	<b>170 000</b>	<b>1 411 618</b>	<b>2 546 733</b>	<b>10 961 909</b>	<b>13 457 550</b>	<b>16 265 597</b>	<b>19 357 982</b>	<b>21 491 775</b>

**TABLE 7.2.**

**Average number of children targeted and treated, and total treatment doses targeted and delivered, in countries implementing SMC, 2019** Sources: NMPs, LSHTM and MMV.

Country	Average number of children targeted	Average number of children treated	Total treatments targeted	Total treatments delivered
Benin	117 470	114 165	469 881	456 660
Burkina Faso	3 588 271	3 298 397	14 353 085	13 193 588
Cameroon	1 687 880	1 681 737	6 751 520	6 726 948
Chad	1 424 920	1 491 905	5 699 681	5 967 620
Gambia	142 695	110 870	570 780	443 480
Ghana	1 074 214	964 956	4 296 856	3 859 824
Guinea	726 402	750 903	2 905 606	3 003 612
Guinea-Bissau	93 364	82 918	373 456	331 672
Mali	3 548 968	3 767 205	14 195 872	15 068 820
Niger	4 188 304	4 151 103	16 753 217	16 604 412
Nigeria	3 989 073	4 110 152	15 956 290	16 440 608
Senegal	821 473	671 132	3 285 893	2 684 528
Togo	346 259	296 332	1 385 035	1 185 328
<b>Total</b>	<b>21 749 293</b>	<b>21 491 774</b>	<b>86 997 172</b>	<b>85 967 096</b>

LSHTM: London School of Hygiene & Tropical Medicine; MMV: Medicines for Malaria Venture; NMP: national malaria programme; SMC: seasonal malaria chemoprevention.




### 7.4 COVERAGE OF IPTp USE BY DOSE

To date, 33 African countries have adopted IPTp to reduce the burden of malaria during pregnancy. These countries reported routine data from health facilities in the public sector on the number of women visiting ANC clinics, and the number receiving the first, second, third and fourth doses of IPTp (i.e. IPTp1, IPTp2, IPTp3 and IPTp4). Using annual expected pregnancies as the

denominator (adjusted for fetal loss and stillbirths), the percentage of IPTp use by dose was computed. Despite a slight increase in IPTp3 coverage from 31% in 2018 to 34% in 2019, coverage remains well below the target of at least 80% and underscores the substantial number of missed opportunities, given that 62% of women receive IPTp1 (Fig. 7.6).

**FIG. 7.6.** Percentage of pregnant women attending an ANC clinic at least once and receiving IPTp, by dose, sub-Saharan Africa, 2010–2019 Source: NMP reports, US CDC and Prevention estimates and WHO estimates.



ANC: antenatal care; CDC: Centers for Disease Control and Prevention; IPTp: intermittent preventive treatment in pregnancy; IPTp1: first dose of IPTp; IPTp2: second dose of IPTp; IPTp3: third dose of IPTp; NMP: national malaria programme; US: United States; WHO: World Health Organization.




## 7.5 MALARIA DIAGNOSIS AND TREATMENT

This section presents information on manufacturer sales and deliveries and national distribution of RDTs and ACTs, treatment seeking for fever in children aged under 5 years, and population-level coverage of malaria diagnosis and treatment with ACTs. RDT data shown in this section reflect sales by manufacturers eligible for procurement (i.e. under the Malaria RDT Product Testing Programme) from 2010 to 2017, and since 2018 for WHO Prequalification, and NMP distributions of RDTs. The types of ACTs tracked are those recommended by WHO for use in the treatment of clinical malaria.

Globally, 2.7 billion RDTs for malaria were sold by manufacturers in 2010–2019, with nearly 80% of these sales being to sub-Saharan African countries. In the same period, NMPs distributed 1.9 billion RDTs – 84% in sub-Saharan Africa (**Fig. 7.7**). In 2019, 348 million RDTs were sold by manufacturers and 267 million distributed by NMPs. RDT sales and distributions in 2019 were lower than those reported in 2018, by 63 million and 24 million, respectively, with most decreases being in sub-Saharan Africa.

**FIG. 7.7.**

**Number of RDTs sold by manufacturers and distributed by NMPs for use in testing suspected malaria cases, 2010–2019<sup>a</sup>** Sources: NMP reports and sales data from manufacturers eligible for the WHO Malaria RDT Product Testing Programme.



NMP: national malaria programme; *P. falciparum*: *Plasmodium falciparum*; RDT: rapid diagnostic test; WHO: World Health Organization.  
<sup>a</sup> NMP distributions do not reflect those RDTs still in storage that have yet to be delivered to health facilities and community health workers.

## 7 Distribution and coverage of malaria prevention, diagnosis and treatment

More than 3.1 billion treatment courses of ACT were sold globally by manufacturers in 2010–2019 (Fig. 7.8). About 2.1 billion of these sales were to the public sector in malaria endemic countries, and the rest were sold through either public or private sector co-payments (or both), or exclusively through the private retail sector. National data reported by NMPs show that, in the same period, 1.9 billion ACTs were delivered to health service providers to treat malaria patients in the public health


sector. In 2019, some 190 million ACTs were sold by manufacturers to the public health sector; in that same year, 183 million ACTs were distributed to this sector by NMPs, of which 90% were in sub-Saharan Africa.

Aggregated data from household surveys conducted in sub-Saharan Africa between 2005 and 2019 in 21 countries<sup>1</sup> with at least two surveys (baseline – 2005–2011 and most recent – 2015–2019) in this period

<sup>1</sup> Angola (MIS 2011; DHS 2018), Benin (DHS 2006; DHS 2017–2018), Burkina Faso (DHS 2010; MIS 2017–2018), Burundi (DHS 2010; DHS 2016–2017), Cameroon (DHS 2011, DHS 2018), Ghana (DHS 2008; MIS 2019), Guinea (DHS 2005; DHS 2018), Kenya (DHS 2008–2009; MIS 2015), Liberia (MIS 2011; MIS 2016), Madagascar (MIS 2011; MIS 2016), Malawi (DHS 2010; MIS 2017), Mali (DHS 2006; DHS 2018), Mozambique (DHS 2011; MIS 2018), Nigeria (MIS 2010; DHS 2018), Rwanda (DHS 2010; MIS 2017), Senegal (DHS 2010–2011; DHS 2018), Sierra Leone (DHS 2008; MIS 2016), Uganda (DHS 2011; MIS 2018–2019), United Republic of Tanzania (DHS 2010; MIS 2017), Zambia (DHS 2007; DHS 2018) and Zimbabwe (DHS 2010–2011; DHS 2015).

**FIG. 7.8.**

**Number of ACT treatment courses delivered by manufacturers and distributed by NMPs to patients, 2010–2019<sup>a,b</sup>** Sources: Companies eligible for procurement by WHO/UNICEF and NMP reports.



ACT: artemisinin-based combination therapy; AMFm: Affordable Medicines Facility–malaria; GF: Global Fund to Fight AIDS, Tuberculosis and Malaria; NMP: national malaria programme; UNICEF: United Nations Children’s Fund; WHO: World Health Organization.

<sup>a</sup> NMP deliveries to patients reflect consumption reported in the public health sector.

<sup>b</sup> AMFm/GF indicates that the AMFm operated from 2010 to 2013, with the GF co-payment mechanism operating from 2014.



were used to analyse coverage of treatment seeking, diagnosis and use of ACTs by children aged under 5 years (**Table 7.3**). Comparing baseline and latest surveys, there was little change in prevalence of fever within the 2 weeks preceding surveys (median 24% versus 21%) and treatment seeking for fever (median 64% versus 69%). Comparisons of the source of treatment between baseline and more recent surveys shows that a median 63% versus 71% received care from public health facilities, and a median 39% versus 30% from the private sector. Use of community health workers was low in both periods, at a median of less than 2%.

The rate of diagnosis among children aged under 5 years for whom care was sought increased considerably, from a median of 15% at baseline to 38% in the latest household surveys. Use of ACTs also increased more than twofold, from 39% at baseline to 81% in the latest surveys when all children with fever for whom care was sought were considered. Among those who received a finger or heel prick, use of ACTs was 42% in the most recent survey, suggesting that many children received ACTs without parasitological diagnosis.

**TABLE 7.3.**

**Summary of coverage of treatment seeking for fever, diagnosis and use of ACTs for children aged under 5 years from household surveys in sub-Saharan Africa, at baseline (2005–2011) and most recent (2015–2019)** *Source: household surveys.*

Children aged under 5 years	Baseline (2005–2011)			Most recent survey (2015–2019)		
	Median estimate	Lower bound	Upper bound	Median estimate	Lower bound	Upper bound
<b>Prevalence of fever</b>						
With fever in past 2 weeks	24.1%	18.3%	34.3%	20.6%	16.1%	30.9%
<b>Treatment seeking for fever</b>						
With fever in past 2 weeks for whom treatment was sought	63.5%	57.7%	71.6%	69.1%	56.3%	73.8%
<b>Source of treatment for fever among those who were treated</b>						
Public sector (health facility)	62.9%	52.8%	80.3%	71.0%	49.0%	85.0%
Public sector (community health worker)	2.0%	0.2%	3.4%	1.3%	0.4%	4.9%
Private sector (formal and informal)	39.1%	21.6%	50.3%	30.2%	16.3%	51.9%
<b>Diagnosis among those with fever and for whom care was sought</b>						
Received a finger or heel prick	15.4%	6.5%	26.9%	37.7%	17.8%	49.1%
<b>Use of ACTs among those for whom care was sought</b>						
Received treatment with ACTs	38.9%	23.6%	68.2%	80.5%	30.6%	93.4%
<b>Use of ACTs among those for whom care was sought and received a finger or heel prick</b>						
Received ACTs	18.9%	14.3%	37.7%	42.4%	17.1%	58.7%

ACT: artemisinin-based combination therapy.

## 7 Distribution and coverage of malaria prevention, diagnosis and treatment


Analysis of equity of fever prevalence and treatment seeking at subnational level was conducted using the

most recent household survey data for 2015–2019, from 23 countries<sup>1</sup> (Fig. 7.9). In most countries, children in

<sup>1</sup> Angola (DHS 2018), Benin (DHS 2017–2018), Burkina Faso (MIS 2017–2018), Burundi (DHS 2016–2017), Cameroon (DHS 2018), Ethiopia (DHS 2016), Ghana (MIS 2019), Guinea (DHS 2018), Kenya (MIS 2015), Liberia (MIS 2016), Madagascar (MIS 2016), Malawi (MIS 2017), Mali (DHS 2018), Mozambique (MIS 2018), Nigeria (DHS 2018), Rwanda (MIS 2017), Senegal (DHS 2018), Sierra Leone (MIS 2016), Togo (MIS 2017), Uganda (MIS 2018–2019), United Republic of Tanzania (MIS 2017), Zambia (DHS 2018) and Zimbabwe (DHS 2015).

**FIG. 7.9.**

**Concentration index of a) prevalence of fever in, and b) care seeking for children aged under 5 years at administrative level 1, sub-Saharan Africa** Source: most recent household surveys from the period 2015–2019, Kenya Medical Research Institute – Wellcome Trust Research Programme.






poorer households had a higher prevalence of having a fever in the 2 weeks preceding the household surveys (i.e. concentration index  $<0$ ). In contrast, treatment

seeking was higher in febrile children from wealthier households in all subnational units, although in some units the difference was small.

b)



# 8 Global progress towards the GTS milestones

The GTS aims for a reduction in malaria case incidence and mortality rate of at least 40% by 2020, 75% by 2025 and 90% by 2030 from a 2015 baseline (4). Trends in malaria cases and deaths were used to make annual projections from 2020 to 2030, to track progress towards the targets and milestones of the GTS as mandated to WHO by the World Health Assembly (4). The projections presented here do not account for potential disruptions due to the COVID-19 pandemic, which – despite commendable global and national efforts to maintain essential malaria services – is likely to lead to higher than expected malaria morbidity and mortality (Section 10).



## 8.1 GLOBAL PROGRESS

Despite the considerable progress made since 2000, the GTS 2020 milestones for morbidity and mortality will not be achieved globally (Fig. 8.1). Without actions to reverse this trend, the 2030 GTS and SDG targets for malaria morbidity and mortality will also not be met (Fig. 8.1). The malaria case incidence of 56 per 1000 population at risk in 2020 instead of the expected 35 cases per 1000 means that, globally, we are off track by 37%; at the current trajectory, we could be off track by 87% in 2030 (Fig. 8.1a). Although relative progress in the mortality rate is greater than that in case incidence (see Section 3 for potential methodological reasons), globally projected malaria deaths per 100 000 population at risk in 2020 was projected to be 9.8, reducing from 11.9 in 2015. This implied that globally we were off track by 22% (Fig. 8.1b).

Fig. 8.2 and Fig. 8.3 on the next page present progress in all countries considered to be malaria endemic in 2015. Countries were ranked into eight categories of reduction of case incidence and mortality rates in 2020 from a 2015 baseline:

- achieved zero malaria by 2020;
- reduced by 40% or more;
- reduced by between 25% and less than 40%;
- reduced by less than 25%;
- no change since 2015 (less than 5% increase or decrease in case incidence or mortality rate);

- increased by less than 25%;
- increased by between 25% and less than 40%; and
- increased by 40% or more.


Of the 92 countries that were malaria endemic globally in 2015, 31 (34%) were estimated to be on track for the GTS morbidity milestone for 2020, having achieved 40% or more reduction in case incidence or reported zero malaria cases. Another 21 (23%) had made progress in reducing malaria case incidence but were not on track for the GTS milestone. Thirty-one countries (34%) are estimated to have experienced increased incidence, with 15 countries (16%) estimated to have experienced an increase of 40% or more in malaria case incidence in 2020 compared with 2015. Malaria case incidence in nine (10%) countries in 2020 was estimated to be at levels similar to those of 2015.

Thirty-nine (42%) countries that were malaria endemic in 2015 were on track for the GTS mortality milestone for 2020, with 28 of them reporting zero malaria cases. An additional 34 countries (37%) were estimated to have achieved reductions in mortality rate but progress was below the 40% target. Malaria mortality rates remained at the same level in 2020 as in 2015 in seven countries (8%), while another 12 countries (13%) had estimated increases, with six of these countries having increases of 40% or more.



**FIG. 8.1.**

**Comparison of global progress in malaria: a) case incidence and b) mortality rate, considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)** *Source: WHO estimates.*




GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.



**FIG. 8.2.**

**Map of malaria endemic countries showing progress towards the GTS 2020 malaria case incidence milestone of at least 40% reduction from a 2015 baseline** *Source: WHO estimates.*




GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization.



**FIG. 8.3.**

Map of malaria endemic countries showing progress towards the GTS 2020 malaria mortality rate milestone of at least 40% reduction from a 2015 baseline *Source: WHO estimates.*



GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization.


### 8.2 WHO AFRICAN REGION

Analysis of the trends by region shows that the WHO African Region is off track for both the malaria morbidity and mortality 2020 GTS milestones, by 37% and 25%, respectively (Fig. 8.4). Only Botswana, Cabo Verde, Ethiopia, the Gambia, Ghana, Namibia and South Africa are on track to achieve the GTS 2020 target of a 40% reduction in malaria case incidence, and Algeria has already been certified malaria free.

Although not on track, 17 countries (Equatorial Guinea, Gabon, Guinea, Guinea-Bissau, Kenya, Malawi, Mali,

Mauritania, Mozambique, Niger, Senegal, Sierra Leone, South Africa, Togo, United Republic of Tanzania, Zambia and Zimbabwe) were estimated to have achieved reductions in malaria case incidence by 2020 compared with 2015 (Fig. 8.2). There was no difference (<5% increase or decrease) in case incidence in 2020 compared with 2015 in Benin, Burkina Faso, Cameroon, Central African Republic, Liberia, Madagascar, Nigeria, South Sudan and Uganda. Case incidence was higher in 2020 than in 2015 by less than 25% in Angola, Chad, Congo, Côte d'Ivoire, Democratic Republic of the

**FIG. 8.4.** Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO African Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)  
Source: WHO estimates.




GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.



Congo, Rwanda, and Sao Tome and Principe, and increased by 40% or more in Burundi, Comoros, Eritrea and Eswatini.

Botswana, Cabo Verde, Eswatini, and Sao Tome and Principe reported zero malaria deaths in 2019 and were projected to maintain this in 2020 (Fig. 8.3). Ethiopia and Namibia were estimated to have achieved a reduction in mortality rate of more than 40%. Although not on track for the GTS 2020 mortality milestones, 30 countries (Angola, Benin, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad,

Congo, Côte d'Ivoire, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Gambia, Ghana, Guinea, Kenya, Malawi, Mali, Mauritania, Mozambique, Niger, Nigeria, Senegal, Sierra Leone, South Africa, Togo, Uganda, United Republic of Tanzania, Zambia and Zimbabwe) had achieved mortality rate reductions of less than 40%. Guinea-Bissau, Liberia, Madagascar, Rwanda and South Sudan showed no change in levels of mortality rate (<5% decrease or increase) in 2020 compared with 2015, whereas increases in mortality rate of more than 40% were reported in Comoros, Eritrea and Sudan.



GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.


### 8.3 WHO REGION OF THE AMERICAS

In the WHO Region of the Americas, both Belize and El Salvador had zero malaria cases in 2019 and are projected to remain unchanged in 2020. Belize, French Guiana, Guatemala, Haiti, Honduras and Peru were all on target for the 2020 malaria morbidity GTS milestone of a reduction of at least 40% in case incidence (Fig. 8.5). Bolivia (Plurinational State of), Brazil, Mexico and Suriname are estimated to have reduced malaria case incidence by less than 25% in 2020 compared with 2015. Colombia, Costa Rica, Dominican Republic, Ecuador, Guyana, Nicaragua, Panama and Venezuela

(Bolivarian Republic of) are estimated to have increases in case incidence of more than 40% in 2020 compared with 2015.

At regional level, most of the worsening of the trend is attributable to the epidemic in Venezuela (Bolivarian Republic of). Progress analysis in the WHO Region of the Americas shows that the region would be about 43% off the GTS 2020 malaria case incidence milestones *with* the estimated cases in Venezuela (Bolivarian Republic of) and 15% off *without* those

**FIG. 8.5.** Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO Region of the Americas considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green) Source: WHO estimates.




GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.



estimated cases (**Fig. 8.5**). Urgent control of the epidemic in Venezuela (Bolivarian Republic of) is required to get the region back on track.

There are few malaria deaths in the WHO Region of the Americas, and changes in 2020 relative to the 2015 GTS baseline should be interpreted with caution. For example, although the mortality rate in Bolivia

(Plurinational State of), Dominican Republic and Nicaragua has increased by more than 40% (**Fig. 8.3**), it is estimated that the actual number of deaths would be fewer than 15 in all these countries. Malaria deaths in Venezuela (Bolivarian Republic of), however, are estimated to have doubled and there have been more than 400 cases in 2020.




GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.

### 8.4 WHO EASTERN MEDITERRANEAN REGION

Overall, the WHO Eastern Mediterranean Region is off track for both the 2020 GTS milestone for malaria morbidity and mortality, by twice the expected levels (Fig. 8.6). However, the Islamic Republic of Iran has reported no indigenous malaria cases in 2018 and 2019, and Saudi Arabia has reduced case incidence by more than 40%. Although not on track for the GTS 2020 case incidence milestones, Pakistan and Somalia have reduced case incidence, but by less than 40% in 2020

compared with 2015. Djibouti and Sudan were both off track, with malaria case incidence higher by more than 40% in 2020 compared with 2015. Afghanistan and Yemen’s case incidence was higher in 2020 than in 2015, but by less than 25% in Afghanistan and by 25% to less than 40% in Yemen (Fig. 8.3). Malaria mortality rate had decreased by less than 25% in Afghanistan and Somalia, and by between 25% and 40% in Pakistan in 2020 compared with 2015.

**FIG. 8.6.** Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO Eastern Mediterranean Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green) Source: WHO estimates.






## 8.5 WHO SOUTH-EAST ASIA REGION

The WHO South-East Asia Region is on track for both the mortality and morbidity milestones (Fig. 8.2, Fig. 8.3, Fig. 8.7). Sri Lanka was certified malaria free in 2015 and remains malaria free. Timor-Leste reported zero malaria cases and deaths in 2019. All other

countries reduced malaria case incidence by 40% or more, and mortality rate by more than 40%, except Indonesia where the rate reduced by between 25% and less than 40% in 2020 compared with 2015 (Fig. 8.2, Fig. 8.3).

**FIG. 8.7.**

**Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO South-East Asia Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)** Source: WHO estimates.



GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.




### 8.6 WHO WESTERN PACIFIC REGION

Overall, the WHO Western Pacific Region was off track for both the malaria morbidity and mortality 2020 GTS milestones by 50%, and at the current trajectory the burden could increase through to 2030 (Fig. 8.8). However, most of this increase in burden is attributable

to Papua New Guinea, which accounts for about 80% of the burden of malaria in the region. Malaria case incidence was higher by 25% or less in Vanuatu, by between 25% and 40% in Papua New Guinea and the Philippines, and by 40% or more in the Solomon

**FIG. 8.8.** Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO Western Pacific Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green) Source: WHO estimates.




GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.



Islands (Fig. 8.2). However, China and Malaysia both reported zero malaria cases in 2019 and were expected to maintain this into 2020. Case incidence reduced by 40% or more from the 2015 baseline in Cambodia and Lao People's Democratic Republic, and

by between 5% and 25% in the Republic of Korea and Viet Nam. When Papua New Guinea is excluded from analysis, the projections suggest that the region is almost on track for the 2020 GTS incidence milestones (Fig. 8.8).



GTS: Global technical strategy for malaria 2016–2030; WHO: World Health Organization; WMR: world malaria report.



# Biological threats

## 9.1 DELETIONS IN *P. FALCIPARUM* HISTIDINE-RICH PROTEIN 2 AND PROTEIN 3 GENES

Histidine-rich protein 2 (HRP2) is the predominant target of the 345 million *P. falciparum*-detecting malaria RDTs sold annually. Parasites that no longer express HRP2 may not be detectable by RDTs based on HRP2, and those that no longer express HRP2 and histidine-rich protein 3 (HRP3) are completely undetectable by these RDTs. Deletions in the *P. falciparum* genes for HRP2 (*pfhrp2*) and HRP3 (*pfhrp3*) in clinical isolates were first identified in 2010 in the Peruvian Amazon basin, by researchers characterizing blood samples that were negative by HRP2-based RDTs but positive by microscopy (71). In recent years, *pfhrp2/3*-deleted parasites have been documented outside of South America, including in Asia, the Middle East, and Central, East, Southern and West Africa. Prevalence estimates vary widely both within and between countries. The examples of Eritrea and Peru – where the prevalence of dual *pfhrp2* and *pfhrp3* deleted parasites among symptomatic patients reached as high as 80% – demonstrate that these parasites can become dominant in the population, posing a serious global threat to patients and to the efficacy of HRP2-based RDTs.

WHO has published guidance on investigating suspected *pfhrp2/3* deletions (132), and recommends that countries that have reports of *pfhrp2/3* deletions, and their neighbouring countries, should conduct representative baseline surveys among suspected malaria cases, to determine whether the prevalence of *pfhrp2/3* deletions causing false negative RDT results has reached a threshold for RDT change (>5% *pfhrp2* deletions causing false negative RDT results). Alternative RDT options (e.g. based on detection of the

*Plasmodium* lactate dehydrogenase [pLDH]) are limited; in particular, there are currently no WHO-prequalified non-HRP2 combination tests that can detect and distinguish between *P. falciparum* and *P. vivax*.

WHO is tracking published reports of *pfhrp2/3* deletions using the Malaria Threats Map mapping tool (100, 133), and is encouraging a harmonized approach to mapping and reporting of *pfhrp2/3* deletions through publicly available survey protocols. Among the 39 reports published by 39 countries, 32 (82%) reported *pfhrp2* deletions, but variable methods in sample selection and laboratory analysis mean that the scale and scope of clinically significant *pfhrp2/3* deletions is still unclear. Between 2019 and September 2020, investigations for *pfhrp2/3* deletions were reported in 16 publications from 15 countries. *Pfhrp2/3* deletions were confirmed in 12 reports from 11 countries: China, Equatorial Guinea, Ethiopia, Ghana, Myanmar, Nigeria, Sudan, Uganda, United Kingdom (imported from various malaria endemic countries), the United Republic of Tanzania and Zambia. No deletions were identified in France (among returning travellers), Haiti, Kenya and Mozambique.

The WHO Global Response Plan for *pfhrp2/3* deletions outlines several areas for action beyond scaling up surveillance. The plan includes identifying new biomarkers, improving the performance of non-HRP2-based RDTs, market forecasting and strengthening laboratory networks to support the demands of molecular characterization to determine the presence or absence of these gene deletions.



## 9.2 THERAPEUTIC EFFICACY OF ACTs

Effective treatment for malaria is a critical component of malaria control and elimination. The emergence of multidrug resistance, including resistance to artemisinin and partner drugs, threatens the global effort to reduce the burden of malaria. The GTS calls on countries and global malaria partners to monitor the efficacy of antimalarial medicines, to ensure that the most appropriate and effective treatments are selected for national treatment policies (4).

Therapeutic efficacy studies (TES) track clinical and parasitological outcomes in patients after they have received antimalarial treatment. When conducted according to the WHO protocol, TES offer a consistent measure of treatment efficacy over time. These studies provide NMPs with the data required to evaluate their treatment policies and make changes where necessary. In areas of malaria elimination, the routine surveillance system incorporates the treatment and follow-up of all malaria cases. In this context, the data generated on patient outcomes become part of integrated drug efficacy surveillance (iDES) (135).

This section summarizes TES findings from studies conducted on patients infected with *P. falciparum* and *P. vivax* for each WHO region between 2010 and 2019.

Given that ACTs are currently the recommended first-line treatment in all malaria endemic countries, and artesunate (injectable) is the main treatment for severe malaria, **Section 9.3** summarizes the prevalence of *PfKelch13* molecular mutations associated with artemisinin partial resistance. The latest available information and references can be found online in the Malaria Threats Map, which provides a geographical representation of drug efficacy and resistance data.<sup>1</sup> The data from the most recent TES are also summarized in reports available online.<sup>2</sup>

### 9.2.1 WHO African Region

In the WHO African Region, the first-line treatments for *P. falciparum* include artemether-lumefantrine (AL), artesunate-amodiaquine (AS-AQ) and dihydroartemisinin-piperazine (DHA-PPQ). The overall average efficacy rates for *P. falciparum* – 98.0% for AL, 98.4% for AS-AQ and 99.4% for DHA-PPQ – remained consistent over time (**Fig. 9.1**). Treatment failure rates of more than 10% were observed in four studies of AL but can be considered statistical outliers. There is no evidence of confirmed lumefantrine resistance in Africa. For all other medicines, treatment failure rates remain below 10%.


<sup>1</sup> See <https://www.who.int/malaria/maps/threats-about/en/>.

<sup>2</sup> See [https://www.who.int/malaria/areas/drug\\_resistance/drug\\_efficiency\\_database/en/](https://www.who.int/malaria/areas/drug_resistance/drug_efficiency_database/en/).

**FIG. 9.1.**

#### Treatment failure rates among patients with *P. falciparum* malaria, WHO African Region, 2010–2019

Source: WHO Global database on antimalarial drug efficacy and resistance.



AL: artemether-lumefantrine; AS-AQ: artesunate-amodiaquine; AS-MQ: artesunate-mefloquine; AS-PY: artesunate-pyronaridine; AS+SP: artesunate-sulfadoxine-pyrimethamine; DHA-PPQ: dihydroartemisinin-piperazine; *P. falciparum*: *Plasmodium falciparum*; WHO: World Health Organization.

## 9 Biological threats

*P. vivax* is only endemic in a few countries in the WHO African Region. In Ethiopia, the AL efficacy rate was low in one study, probably due to lumefantrine's short half-life, which does not protect against early relapse. In most studies of CQ in Ethiopia, treatment failure rates were consistently below 10% except in one study that had a treatment failure of 22.0%. No treatment failures were observed in TES of AS-AQ in Madagascar and CQ in Mauritania.

### 9.2.2 WHO Region of the Americas

The first-line treatments for *P. falciparum* in the WHO Region of the Americas include AL (in Bolivia [Plurinational State of], Brazil, Colombia, Ecuador, French Guiana, Guyana, Panama, Paraguay, Suriname and Venezuela [Bolivarian Republic of]), AS-MQ (in Brazil, Peru and Venezuela [Bolivarian Republic of]) and CQ (in Dominican Republic, Guatemala, Haiti, Honduras and Nicaragua). Efficacy of AL and AS-MQ remains high in Brazil, Colombia and Suriname.


In all malaria endemic countries in the WHO Region of the Americas, the first-line treatment policy for *P. vivax* is CQ but some ACTs were tested. Countries conducted studies of CQ alone or of CQ combined with primaquine (PQ) (Fig. 9.2). One study of CQ from Bolivia (Plurinational State of) in 2011 detected a treatment failure rate of 10.4%.

### 9.2.3 WHO South-East Asia Region

The first-line treatments for *P. falciparum* in the WHO South-East Asia Region include AL (in Bangladesh, Bhutan, India, Myanmar, Nepal and Timor-Leste), AS-MQ (in Myanmar), AS+SP (in India) and DHA-PPQ (in Bangladesh, Indonesia, Myanmar and Thailand). TES of AL demonstrated high treatment efficacy in Bhutan, India, Myanmar, Nepal and Timor-Leste, with treatment failure of less than 10% in all studies (Fig. 9.3). AL treatment failure rates exceeded 10% in three studies: one in Thailand (11.3% in 2012) and two in Bangladesh (11.1% in 2013 and 14.3% in 2017). Both of the

FIG. 9.2.

Treatment failure rates among patients with *P. vivax* malaria, WHO Region of the Americas, 2010–2019  
Source: WHO Global database on antimalarial drug efficacy and resistance.



CQ: chloroquine; CQ+PQ: chloroquine plus primaquine; *P. vivax*: *Plasmodium vivax*; WHO: World Health Organization.



studies in Bangladesh had small sample sizes ( $n < 10$ ). All TES of AS+SP were conducted in India. Following high rates of treatment failure in the north-eastern provinces, in 2013, India changed its treatment policy in those provinces to AL; AS+SP remains effective elsewhere in the country. TES of AS-AQ were conducted in Indonesia in 2011 and 2012, with a treatment failure rate of 16.7% observed in the 2012 study of 24 patients. TES of AS-MQ were conducted in Myanmar, where the treatment remains effective, and in Thailand, where high rates of treatment failure were observed. TES findings in Thailand led to the adoption of DHA-PPQ as the first-line treatment in 2015. Among the four TES of AS-PY in Myanmar, no treatment failures were observed. Studies of DHA-PPQ were conducted in Indonesia, Myanmar and Thailand. All results from Indonesia and Myanmar demonstrated high rates of treatment efficacy, with treatment failure rates of less than 5%. In Thailand, high rates of treatment failure were observed with DHA-PPQ in two of five studies: 86.7% in a study of 15 patients and 100% in a study of


two patients. Both studies were completed in 2018 in the eastern part of the country; Thailand is currently recommending treatment with AS-PY in this area.

The first-line treatments for *P. vivax* are CQ (in Bangladesh, Bhutan, Democratic People's Republic of Korea, India, Myanmar, Nepal, Sri Lanka and Thailand), AL (in Timor-Leste) and DHA-PPQ (in Indonesia). High treatment efficacy was found in studies of CQ conducted in Bangladesh, Bhutan, the Democratic People's Republic of Korea, India, Myanmar and Nepal except in two studies from Myanmar (11.9% in 2010 and 21.7% in 2012) and one from Timor-Leste (17.5% in 2011). There was high efficacy of AL in the Democratic People's Republic of Korea and Timor-Leste, AS-PY in Myanmar and DHA-PPQ in Indonesia.

**FIG. 9.3.**

**Treatment failure rates among patients with *P. falciparum* malaria, WHO South-East Asia Region, 2010–2019**

Source: WHO Global database on antimalarial drug efficacy and resistance.



AL: artemether-lumefantrine; AS-AQ: artesunate-amodiaquine; AS-MQ: artesunate-mefloquine; AS+SP: artesunate+sulfadoxine-pyrimethamine; AS-PY: artesunate-pyronaridine; DHA-PPQ: dihydroartemisinin-piperaquine; *P. falciparum*: *Plasmodium falciparum*; WHO: World Health Organization.

**9.2.4 WHO Eastern Mediterranean Region**

The first-line treatments for *P. falciparum* in the WHO Eastern Mediterranean Region are AL (in Afghanistan, Pakistan, Somalia and Sudan) and AS+SP (in Iran [Islamic Republic of], Saudi Arabia and Yemen). The TES of AL from Afghanistan, Pakistan, Somalia, Sudan and Yemen all demonstrated good treatment efficacy, with treatment failure rates below 10% (Fig. 9.4). The TES of AS+SP from Somalia and Sudan, conducted from 2011 to 2016, found low efficacy, with treatment failure rates as high as 22.2% in Somalia in 2011 and 18.1% in Sudan in 2014 (Fig. 9.4). This prompted a subsequent change in treatment policy to the use of AL in both countries. Elsewhere, TES of AS+SP from Afghanistan, Iran (Islamic Republic of), Pakistan and Yemen all demonstrated high treatment efficacy, with fewer than 5% of patients failing treatment.

The first-line treatments for *P. vivax* are AL (in Somalia and Sudan) and CQ in all other countries. TES of CQ were conducted in Afghanistan (n=1), Iran (Islamic

Republic of) (n=1) and Pakistan (n=1), all of which showed high treatment efficacy. In addition, TES of AL in Afghanistan (n=4), Somalia (n=1) and Sudan (n=1) demonstrated high treatment efficacy.


**9.2.5 WHO Western Pacific Region**

The first-line treatments for *P. falciparum* in the WHO Western Pacific Region are AL (in Lao People’s Democratic Republic, Malaysia, Papua New Guinea, Philippines, Solomon Islands and Vanuatu), AS-MQ (in Cambodia), DHA-PPQ (in China and Viet Nam) and AS-AQ (in China) (Fig. 9.5).

TES of AL were conducted in Cambodia, Lao People’s Democratic Republic, Malaysia, Papua New Guinea, Philippines, Solomon Islands and Viet Nam. Treatment failure rates were 10% or less in four studies in Lao People’s Democratic Republic but those studies did not have the recommended sample sizes. A study with an adequate number of patients is currently underway to further investigate these high rates of treatment failure.

**FIG. 9.4.**

**Treatment failure rates among patients with *P. falciparum* malaria, WHO Eastern Mediterranean Region, 2010–2019** Source: WHO Global database on antimalarial drug efficacy and resistance.



AL: artemether-lumefantrine; AS+SP: artesunate+sulfadoxine-pyrimethamine; DHA-PPQ: dihydroartemisinin-piperaquine; *P. falciparum*: *Plasmodium falciparum*; WHO: World Health Organization.



All other studies of AL in the region demonstrated high treatment efficacy. TES of AS-MQ conducted in Cambodia, Lao People’s Democratic Republic, Malaysia and Viet Nam showed that the treatment efficacy of AS-MQ has remained high over the past 10 years, except in one 2019 study from Cambodia, where treatment failed in two of 16 patients.

TES of AS-PY were conducted in Cambodia, Lao People’s Democratic Republic and Viet Nam. High rates of treatment failure were observed in two studies from Cambodia in 2014, of 10.2% and 18.0%, but subsequent studies have found treatment failure rates below 5.0%. In one study in Viet Nam from 2017, treatment failed in three of 19 patients; all other studies in Viet Nam and Lao People’s Democratic Republic found treatment failure rates of 5.0% or less.

Studies of DHA-PPQ were conducted in Cambodia, China, Lao People’s Democratic Republic, Papua New Guinea and Viet Nam. Following high rates of treatment failure, Cambodia removed DHA-PPQ from its


treatment policy. High rates of failure for treatment with DHA-PPQ were also observed in Lao People’s Democratic Republic and Viet Nam. AS-PY has become the first-line treatment in the Viet Nam provinces where treatment failures with DHA-PPQ were observed.

The first-line treatments for *P. vivax* in the WHO Western Pacific Region are AL (in Lao People’s Democratic Republic, Malaysia, Papua New Guinea, Solomon Islands and Vanuatu), AS-MQ (in Cambodia) and CQ (in China, Philippines, Republic of Korea and Viet Nam). TES of AL were conducted in Papua New Guinea, Solomon Islands and Vanuatu between 2011 and 2014. High failure rates for treatment with AL were observed in each country: 35.0% in Papua New Guinea, 31.6% in Solomon Islands and 12.1% in Vanuatu. These high failure rates are probably due to the short half-life of lumefantrine, which does not protect against early relapse. TES of AS-MQ conducted in Cambodia, Lao People’s Democratic Republic and Malaysia demonstrated 100% efficacy. TES of AS-PY in Cambodia and Lao People’s Democratic Republic found treatment failure rates below 5%.

**FIG. 9.5.**

**Treatment failure rates among patients with *P. falciparum* malaria, WHO Western Pacific Region, 2010–2019**

Source: WHO Global database on antimalarial drug efficacy and resistance.



AL: artemether-lumefantrine; AS-AQ: artesunate-amodiaquine; AS-MQ: artesunate-mefloquine; AS-PY: artesunate-pyronaridine; DHA-PPQ: dihydroartemisinin-piperazine; *P. falciparum*: *Plasmodium falciparum*; WHO: World Health Organization.



### 9.3 THE GLOBAL PREVALENCE OF *PFKELCH13* MOLECULAR MUTATIONS

Molecular marker studies help to identify and track the prevalence of molecular mutations associated with drug resistance. WHO has established the following list of validated *PfKelch13* markers of partial resistance to artemisinin: F446I, N458Y, M476I, Y493H, R539T, I543T, P553L, R561H, P574L and C580Y. The candidate markers are P441L, G449A, C469F/Y, A481V, R515K, P527H, N537I/D, G538V, V568G, R622I and A675V. In some areas, there is evidence of a clonal expansion of *PfKelch13* mutations associated with artemisinin partial resistance, as discussed below.

Artemisinin partial resistance emerged independently in several foci in the GMS. WHO continues to monitor the situation, which has evolved rapidly since the first detections of *PfKelch13* mutations in the GMS. Some mutations have disappeared, whereas the prevalence of others has increased. Currently, the most prevalent markers west of Bangkok (western Thailand and Myanmar) are F446I, M476I and R561H. The most prevalent markers east of Bangkok (eastern Thailand, Cambodia, Lao People's Democratic Republic and Viet Nam) are Y493H and P553L. Two markers, R539T and C580Y, are also highly prevalent in both areas. The change in treatment policy in Cambodia from

DHA-PPQ to AS-MQ resulted in a reduction in the prevalence of strains carrying both C580Y and PPQ resistance.

Rwanda has detected an increasing prevalence of the R561H mutation, a validated marker that emerged independently in the GMS between 2012 and 2015. The presence of this mutation was confirmed in Rwanda in 2018, but so far it seems that delayed clearance associated with this mutation has not affected the efficacy of the ACTs that are currently among those tested and used in Rwanda. The R622I mutation seems to be appearing independently in Africa, having been found in Eritrea, Ethiopia, Somalia and Sudan, and with increasing frequency in the Horn of Africa. The ACTs used in these four countries remain effective, despite the presence of the mutation. Further investigation of delayed parasite clearance is needed in this region.

In Guyana, the C580Y mutation also emerged independently between 2010 and 2017. However, in recent studies (including surveys and TES), 100% of samples were found to be wild type, indicating that the mutation may be disappearing in Guyana.

### 9.4 VECTOR RESISTANCE TO INSECTICIDES

Resistance of malaria vectors to pyrethroid insecticides that are commonly used for malaria vector control – namely, pyrethroids, organophosphates, carbamates and the rarely used organochlorine dichlorodiphenyl-trichloroethane (DDT) – threatens malaria control and elimination efforts.

#### 9.4.1 Update on the status of data reporting

From 2010 through 2019, a cumulative total of 82 countries reported data. The extent and frequency of insecticide resistance monitoring continues to vary considerably between countries, despite continued increases in the number of sites from which standard resistance monitoring data were reported, from 3143 in 2010–2018 to 3559 in 2010–2019. The number of sites per country for which resistance monitoring data were reported between 2010 and 2019 varied widely, from 1 to 287. Pyrethroids continue to be the most frequently monitored insecticide class.

A total of 66 countries reported insecticide resistance monitoring data at least once over the past 3 years and 16 did not report such data. Among 82 countries, only 28 have reported on their insecticide resistance status consistently every year for the past 3 years. Low reporting in 2019 was probably due to competing priorities arising from the COVID-19 pandemic.

Although 29% of the countries that used IRS reported the status of insecticide resistance for every insecticide class used in the year of implementation or the preceding year, concerningly, 57% of countries did not report the status for at least one of the insecticide classes used and 14% did not report the status for any insecticide class used. Although this may reflect a gap in data reporting to WHO, malaria endemic countries are highly encouraged to ensure adequate monitoring of resistance to insecticide classes that are either in use or under consideration for use in malaria vector control interventions, and to prioritize monitoring of these classes.


#### 9.4.2 Update on the status of insecticide resistance

Of the 82 countries that reported resistance monitoring data to WHO, 73 confirmed resistance to at least one insecticide in one malaria vector species from one mosquito collection site in 2010–2019. The number of countries that reported insecticide resistance to all four main insecticide classes in at least one malaria vector species increased from 26 in 2010–2018 to 28 in 2010–2019, and among those 28, 17 reported resistance to three of the four classes between 2010 and 2019 (Fig. 9.6). Of those countries that reported insecticide resistance monitoring data to WHO, the proportion of



**FIG. 9.6.**

**Number of classes to which resistance was confirmed in at least one malaria vector in at least one monitoring site, 2010–2019** Source: national health institutes, national implementation partners, NMPs, research institutions and scientific publications.



NMP: national malaria programme.

## 9 Biological threats

countries that confirmed resistance to these insecticide classes was as follows: 86.4% for pyrethroids, 80.6% for organochlorines, 68.8% for carbamates and 58.8% for organophosphates. Only nine countries that reported data did not confirm resistance to any insecticide class.

Resistance to the four main insecticide classes was detected in all WHO regions except the WHO European Region. Globally, resistance to pyrethroids was detected in at least one malaria vector in 69.9% of the sites for which data were available, while resistance to organochlorines was reported in 63.4% of the sites. Resistance to carbamates and to organophosphates was less prevalent, being detected in 31.7% and 24.9%, respectively, of the sites that reported monitoring data. However, the geographical extent of confirmed resistance to each insecticide class differed considerably across regions (**Fig. 9.7**). Maps showing the status of insecticide resistance to different insecticides at each site are available on the Malaria Threats Map website (100).

There is continued improvement in the collection and reporting of data to guide deployment of recently prequalified vector control tools covered by WHO policy recommendations. The number of countries that monitored the involvement of metabolic resistance mechanisms by means of PBO pre-exposure bioassays increased from 23 in 2010–2018 to 30 in 2010–2019. All 30 countries detected partial or full involvement of metabolic resistance mechanism in phenotypic resistance to pyrethroids in at least one monitoring site for at least one vector species and one pyrethroid insecticide. The number of sites where the involvement of metabolic resistance mechanisms in pyrethroid resistance was monitored by means of PBO pre-exposure bioassays increased by more than twofold, reaching 438 by 2019. Full or partial involvement of metabolic resistance mechanisms for at least one vector species and one pyrethroid insecticide was reported in 392 sites.

Results of biochemical and molecular assays conducted to detect metabolic resistance mechanisms were available for 35 countries and 308 sites for 2010–2019. Of the sites for which reports were available, mono-oxygenases were detected in 66.9%, glutathione-S-transferases in 74.6%, esterases in 74.8% and acetylcholinesterases in 73.2%. Results of assays conducted to detect target-site resistance mechanisms were available for 39 countries and 539 sites. *Kdr L1014F* was detected in 76.3% of the sites and *Kdr L1014S* in 48.9% of the sites.

Recently, WHO Member States and their implementing partners have started to explore insecticide dosages to monitor resistance to neonicotinoid and pyrrole insecticides using two assays: the WHO tube test and the US CDC bottle bioassay. To date, WHO has received

a total of 1326 test results from 323 sites in 23 countries from the WHO regions of Africa and the Western Pacific. A formal WHO process to establish discriminating dosages and test procedures for these two insecticide classes is ongoing. The data reported so far to WHO on mosquito mortality after exposure to neonicotinoid and pyrrole insecticides will be assessed against these discriminating dosages once they have been fully defined. Also, WHO test procedures for insecticide resistance monitoring will be updated to incorporate the new discriminating dosages and potential changes to the methodology. Until that time, Member States are discouraged from using data generated by means of non-validated procedures to arrive at conclusions regarding the resistance status of their local vector populations to these insecticide classes.

All the standard insecticide resistance data reported to WHO are included in the WHO global insecticide resistance database (136) and are available for exploration via the online interactive data visualization tool Malaria Threats Map (100). The latest version of this tool, launched in 2020, provides a summary table with the status of phenotypic resistance and resistance mechanisms for each country; presents maps to inform discussions on the deployment of pyrethroid-PBO nets; allows for download of selected datasets; and includes an animation of the evolution of insecticide resistance, as per reports received by WHO.

### 9.4.3 Mitigating and managing insecticide resistance

The selection of effective vector control interventions needs to be based in part on representative data on the susceptibility of local vectors to insecticides that are covered by a policy recommendation and prequalified by WHO. In addition, insecticide resistance data are crucial for assessing the potential impact that resistance may have on the effectiveness of malaria vector control, an area that continues to be poorly understood. To meet these data needs, countries and their implementing partners are advised to conduct regular insecticide resistance monitoring following the WHO-recommended *Test procedures for insecticide resistance monitoring in malaria vector mosquitoes* (137), and to report and share results in a timely manner. To facilitate reporting, WHO has developed data reporting templates (138) and DHIS2 modules (139) for use by its Member States and their implementing partners, and is supporting the rollout of these tools.

The impact of insecticide resistance on the effectiveness of malaria vector control interventions continues to be poorly understood; however, it is highly likely that such resistance reduces the efficacy of currently available interventions. Countries should therefore not delay in



implementing effective policies and practices for the prevention, mitigation and management of resistance. Two relatively new vector control options that should be considered as part of an insecticide resistance management strategy – pyrethroid-PBO nets and neonicotinoid insecticides for IRS – have been recommended by WHO, and a number of prequalified products that fall into these classes are available. Based on insecticide resistance monitoring data reported to WHO by Member States, and considering recent data from each site, a total of 330 areas in 33 countries currently meet WHO-recommended criteria for pyrethroid-PBO net deployment. Maps showing these sites, along with higher level maps highlighting areas and countries where these sites are present, have been incorporated into the Malaria Threats Map to inform discussions on the deployment of pyrethroid-PBO nets.

To guide resistance management, WHO recommends that countries develop and implement national insecticide resistance monitoring and management plans, drawing on the WHO *Framework for a national plan for monitoring and management of insecticide resistance in malaria vectors* (140). Up to the end of 2019, countries have made considerable progress in developing such plans, with 53 countries having completed plans for resistance monitoring and management, and 28 currently in the process of developing such plans. Further effort and support will be required to ensure that every malaria endemic country has such a plan in place, updates it regularly and has the necessary resources to implement it.

**FIG. 9.7.**

**Reported insecticide resistance status as a proportion of sites for which monitoring was conducted, by WHO region, 2010–2019: pyrethroids, organochlorines, carbamates and organophosphates** Status was based on mosquito mortality, where <90% = confirmed resistance, 90–97% = possible resistance and ≥98% = susceptibility. Where multiple insecticide classes or types, mosquito species or time points were tested at an individual site, the highest resistance status was considered. Numbers above bars indicate the total number of sites (n) for which data were reported. Sources: national health institutes, national implementation partners, NMPs, research institutions and scientific publications.



AFR: WHO African Region; AMR: WHO Region of the Americas; EMR: WHO Eastern Mediterranean Region; EUR: WHO European Region; n: number; NMP: national malaria programme; SEAR: WHO South-East Asia Region; WHO: World Health Organization; WPR: WHO Western Pacific Region.



# Malaria response during the COVID-19 pandemic

## 10.1 THE 2020 COVID-19 PANDEMIC

On 31 December 2019, Chinese authorities alerted WHO to an outbreak of pneumonia cases of unknown cause in Wuhan City, Hubei Province, China. These cases were later confirmed as cases of COVID-19; by the end of January 2020, China had more than 7700 confirmed cases, 12 000 suspected cases and 170 deaths (141). On 30 January 2020, the Director-General of WHO declared the novel coronavirus outbreak a public health emergency of international concern (PHEIC), WHO's highest level of alarm under the International Health Regulations (IHR) (2005) (142). By the first quarter of 2020, COVID-19 – caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV2) – had started spreading aggressively outside China. It became clear that the pandemic would be a major test of the resilience of health systems, even those considered strong and well resourced. Unfortunately, the pandemic continued to spread rapidly, with all countries soon affected. By the second week of November 2020, the COVID-19 pandemic had resulted in more than 54 million cases and more than 1.3 million deaths (143). Older patients and those with certain pre-

existing morbidities had a higher risk of severe disease and death (144). Outside of China, several malaria endemic countries in the WHO South-East Asia Region had reported COVID-19 cases by January 2020. By April 2020, the virus had spread globally to all malaria endemic countries, and by the third week of November 2020, 24 million cases and about 636 000 deaths had been reported (Fig. 10.1).

Brazil and India accounted for more than 64% of all cases reported from malaria endemic countries. In sub-Saharan Africa, a region that accounts for over 90% of malaria infections, the spread of the disease was much slower and case fatality rates were lower than had first been feared. Factors that are being considered as possible contributors to the slower spread in this region include early adoption of aggressive control strategies, prior experience in the control of disease outbreaks, a youthful population, a relatively high proportion of rural population with limited mobility and higher ambient temperatures (145, 146).




In several high-income countries, health systems have become overwhelmed with the efforts required to stop the transmission of the coronavirus, and hospitals have struggled to cope with increasing numbers of severe COVID-19 cases. This led to global concerns about the

potential consequences of the pandemic, including disruptions of essential health services, especially in LMICs, where the population was already dealing with a considerable burden of other infectious diseases.

**FIG. 10.1.**

**Trends in COVID-19 cases and deaths in malaria endemic countries globally and by WHO region (as of 23 November 2020)** Source: WHO Coronavirus disease (COVID-19) dashboard (143).



AFR: WHO African Region; AMR: WHO Region of the Americas; EMR: WHO Eastern Mediterranean Region; SEAR: WHO South-East Asia Region; WHO: World Health Organization; WPR: WHO Western Pacific Region.


# 10 Malaria response during the COVID-19 pandemic

Indeed, the COVID-19 pandemic and restrictions related to the response have caused major disruptions in essential malaria services. Furthermore, early messaging targeted at reducing SARS-CoV2 transmission advised the public to stay at home if they had fever, potentially disrupting treatment seeking for febrile diseases such as malaria. At the same time,

many high malaria burden countries had plans to implement large prevention campaigns before the peak malaria transmission season (which was likely to coincide with peak COVID-19 cases). These plans needed to be adapted to conform with COVID-19 restrictions (**Fig. 10.2**).

**FIG. 10.2.**

**Malaria seasonality and trends of COVID-19 cases in malaria endemic countries and areas, 2020 (as of 23 November 2020)** Source: WHO Coronavirus disease (COVID-19) dashboard (143) and PATH.





## 10.2 GLOBAL WORKSTREAMS ON SUSTAINING THE MALARIA RESPONSE DURING THE COVID-19 PANDEMIC

In March 2020, as the COVID-19 pandemic spread rapidly around the globe, WHO convened a cross-partner effort to mitigate the negative impact of the coronavirus in malaria-affected countries and contribute to the COVID-19 response. The work was carried out in close collaboration with the RBM Partnership to End Malaria, the Global Fund, PMI,

several implementation and advocacy partners, and research institutions.

This collaborative work was implemented across seven cross-partner workstreams set up to address various thematic areas (**Table 10.1**).

TABLE 10.1. The global workstreams on the malaria response during the COVID-19 pandemic	
Workstream	Areas of work
 Clinical trials of COVID-19 treatment with antimalarials and product development	<ul style="list-style-type: none"> <li>■ Develop a generic protocol to evaluate anti-COVID-19 prophylaxis in malaria endemic settings</li> <li>■ Coordinate with researchers</li> <li>■ Disseminate information</li> </ul>
 Modelling, surveillance and clinical epidemiology	<ul style="list-style-type: none"> <li>■ Establish a network of sites involved in clinical epidemiology in countries with malaria transmission</li> <li>■ Consider potential scenarios for COVID-19–malaria interactions and feed these into other workstreams</li> <li>■ Model the impact of service disruptions</li> <li>■ Track country-level service disruptions using routine health information systems</li> </ul>
 Supplies and commodities	<ul style="list-style-type: none"> <li>■ Assess and monitor commodity stocks and supply-chain bottlenecks</li> <li>■ Estimate potential demand for key malaria commodities</li> <li>■ Work with international partners to consider how to use global purchasing power to stimulate ongoing production – and potential stockpiling – of key commodities</li> <li>■ Coordinate and collaborate to optimize global stocks and distribution through careful prioritization</li> <li>■ Work with international financiers to ensure that the necessary resources for the global COVID-19 response do not divert resources away from malaria or other public health priorities</li> </ul>
 Malaria response and guidance	<ul style="list-style-type: none"> <li>■ Develop integrated guidance to support maintenance of essential malaria services</li> <li>■ Ensure the continuation of the effective delivery of malaria interventions within a COVID-19 transmission setting</li> <li>■ Anticipate that the demand for health services may outstrip the ability to deliver routine care</li> <li>■ Consider resource requirements (e.g. commodities and workforce) for extraordinary measures</li> </ul>
 Communications	<ul style="list-style-type: none"> <li>■ Communicate to avoid conflicting advice and misinformation</li> <li>■ Ensure that current advice and public messaging intended to curb coronavirus transmission is appropriate in malaria endemic settings</li> </ul>
 Coordination	<ul style="list-style-type: none"> <li>■ Identify early warning signs of increased costs for implementing malaria programmes or decreased funding for the global malaria response, as both donor and malaria endemic countries respond to COVID-19</li> <li>■ Protect and ensure follow-through on existing commitments (e.g. to the Global Fund) as resources are allocated to the COVID-19 response</li> <li>■ Develop proposals and conduct donor outreach during the COVID-19 pandemic, to fill near-term health system gaps, including critical commodities for malaria and other communicable diseases</li> </ul>
 Resource mobilization	<p>Under the leadership of the RBM Partnership to End Malaria, support countries to mobilize resources, through channels such as the Global Fund and others, to:</p> <ul style="list-style-type: none"> <li>■ purchase PPE to help protect health workers in the provision of services at clinics and during campaigns</li> <li>■ provide emergency resources to adapt the response during COVID-19</li> <li>■ ensure gains are sustained despite the pandemic</li> </ul>



## 10.3 GLOBAL HIGHLIGHTS IN THE MALARIA RESPONSE DURING THE COVID-19 PANDEMIC

### 10.3.1 Partnership alignment and technical guidance

The cross-partner global response achieved several important milestones, starting with an initial urgent call to countries to maintain core malaria control services while protecting health workers and communities against COVID-19 transmission. A WHO statement, shared widely in March 2020, was issued in response to reports that some countries in sub-Saharan Africa had suspended mass ITN campaigns (147). This statement encouraged countries to move forward with ITN, IRS and SMC campaigns, and to advise the public to avoid delays in seeking treatment for illnesses.

To support malaria-affected countries to maintain essential services, in April 2020, WHO issued technical guidance on how to safely maintain malaria control services in the context of the COVID-19 pandemic. This document was developed in close collaboration with partners, and was consistent with broader guidance on maintaining essential services in COVID-19 settings and on facilitating the role of community-based health care during the pandemic. It provided specific malaria guidance on the prevention of infection through vector control and chemoprevention, testing, treatment of cases, clinical services, supply chains and laboratory activities (148).

### 10.3.2 Modelling the potential impact of service disruptions on the burden of malaria

To reinforce the urgent call to maintain essential malaria control services during the pandemic, the WHO GMP, in collaboration with the Malaria Atlas Project (MAP), conducted modelling to quantify the potential impact of service disruptions due to the COVID-19 pandemic (117). This analysis showed that, under the worst-case scenario – in which all ITN campaigns are suspended and there is a 75% reduction in access to effective antimalarial medicines – a staggering 769 000 people in sub-Saharan Africa could die from malaria by the end of 2020. This figure represents a doubling in the number of malaria deaths compared with 2018 and a return to mortality levels last seen 20 years ago. These dire projections were extensively communicated through the media, and directly to the governments of malaria endemic countries and their partners, catalysing an impressive response, with countries tailoring the delivery of essential malaria services to the COVID-19 response, as described below.

### 10.3.3 Responding to the pressure to shift diagnostic production away from malaria

As early as February or March 2020, during the initial acceleration wave of the pandemic, international demand for the development and large-scale production of SARS-CoV2 antigen-detecting rapid immunoassays increased dramatically, driven by the need to diagnose and track the pandemic. By April, some of the world's leading RDT suppliers announced plans to reallocate manufacturing capacity away from malaria RDTs and towards the production of COVID-19 tests. To avoid a potentially devastating shortfall of more than 100 million RDTs, the malaria RDT task force, which involves 15 organizations,<sup>1</sup> began immediate discussion with suppliers that led to the convening of a suppliers' summit in June 2020, attended by 12 companies,<sup>2</sup> including all major manufacturers. In response, the Global Fund and PMI announced tenders to secure unallocated volumes for the remainder of 2020, allowing some flexibility in price offers. The floating of these tenders in July and August secured the malaria RDT requirements for the remainder of 2020, minimizing the risk of stockouts. Since then, PMI and the Global Fund have been expanding their collaborative focus "downstream", tracking RDT supply levels in countries they support and, together with UNICEF and UNDP, coordinating orders and deliveries to minimize disruptions at the country level (Fig. 10.3).

### 10.3.4 Resolving global manufacturing bottlenecks for malaria medicines

In February 2020, preliminary results from small trials employing CQ and hydroxychloroquine (HCQ) for COVID-19 treatment created high expectations for the therapeutic and prophylactic properties of these medicines. These early expectations led to CQ/HCQ treatment of hospitalized COVID-19 patients, and multiple stockpiling initiatives nationally and globally, fed in part by interest from the media, the general public and heads of governments. Unregulated demand by consumers led to instances of cardiotoxicity and death through self-administration of these medicines in several countries. The massive spike in demand for these medicines – normally used for the treatment of *P. vivax* malaria, and conditions such as rheumatoid arthritis and lupus – generated high demand for their active pharmaceutical ingredients. Sales of a key starting material (4,7-dichloroquinoline) increased up to sixfold from April to June 2020. This key starting material is essential for producing other

<sup>1</sup> Bill & Melinda Gates Foundation; Clinton Health Access Initiative (CHAI); FIND; Global Fund; Global Health Supply Chain Program – Procurement and Supply Management (GHSC-PSM); Médecins Sans Frontières (MSF); PATH; PMI; RBM Partnership to End Malaria; UNDP; UNICEF; Unitaid; US CDC; USAID; and WHO GMP and WHO Prequalification Programme.

<sup>2</sup> Abbott, Access Bio Inc., Advy Chemicals, Arkray, Hangzhou Biotest, J. Mitra, Meril, Mologic, Premier Medical Corp, Rapigen, SD Biosensor and Tulip Group.




antimalarial drugs, such as piperazine and amodiaquine; thus, the supply of other critical artemisinin-combination treatments was also constrained. At the time of spike in demand, a major donor sought to ensure that over 120 million tablets of CQ would remain available for deployment for COVID-19 treatment in LMICs, after WHO validation of

properly conducted solidarity trials. Following the release of data showing no benefits of CQ/HCQ for COVID-19, these medicines have been donated to countries in need of CQ for treating their high burden of *P. vivax* malaria (e.g. in Ethiopia, India and certain countries in Latin America).

**FIG. 10.3.**

**Potential RDT stockouts forecast in June 2020, if country orders were not delivered** The July tenders address all but the immediate stockouts through early 2021. Sources: PMI and Global Fund.



### 10.3.5 Mitigating the disruptions in the shipment and delivery of malaria commodities

The COVID-19 pandemic also impacted ITNs and insecticides for IRS, affecting the availability of raw starting materials and production, and the shipment or movement of product between and within countries. Increasing costs of raw materials and freight for many manufacturers, especially in India, could no longer be absorbed in the price of final products. Lockdown measures in countries led to increasing restrictions that limited movement of people and goods; these in turn affected the timely production, packaging, shipment, customs clearance and in-country delivery of goods from countries of manufacture to customer countries. Requirements for COVID-19 testing of drivers who transport goods across borders led to backlogs at ports and borders, and delayed import of goods. Similar factors delayed pre-shipment inspection by limiting movement of personnel. Quality assurance and quality control for ITNs and insecticides were also delayed due to closed laboratories. The availability of and prices for procuring personal protective equipment (PPE) were also affected by the COVID-19 pandemic, because there was high national and international demand for these supplies, especially for N95 masks, which are essential for sprayers engaged in IRS campaigns in 2020 and early 2021. The collaboration of over 20 organizations in tracking progress in ITN and IRS campaigns led to early resolution of bottlenecks, coordinated procurement and delivery, and mobilization of resources for PPE.

### 10.3.6 Supplementing funding for countries

The Global Fund has established an overall response fund of US\$ 1 billion, and has allowed countries to access an amount equivalent to up to 10% of their allocations to help with the response (149). This support includes providing funding to countries to purchase personal protective equipment such as masks, gloves and gowns that are critical for the continuation of non-COVID-19 health care services including malaria. PMI, the second largest donor to the fight against malaria,

has also made significant investments, particularly across its 24 focus countries in sub-Saharan Africa (including in all the HBHI African countries). The investments are for both for enhanced routine programming and flexibilities within existing allocation, to help countries support and adapt their malaria programmes while responding to their COVID-19 situation. Additional specific resource mobilization has also been supported by several other partners.<sup>1</sup>

### 10.3.7 Tracking malaria service disruptions during the COVID-19 pandemic

COVID-19 overwhelmed health delivery systems across the world, requiring adaptation or, in some cases, suspension of routine and elective services. However, many countries are compromised by the lack of accurate and timely data for tracking and monitoring the extent of disruptions to essential health services. This is limiting the understanding of the scale of the problem and hampering the development of locally appropriate mitigation strategies.

A range of global trackers, implemented at different intervals, have been developed by various agencies to monitor disruptions in broader essential health services during the COVID-19 pandemic, including some developed specifically for malaria. Information from these trackers was assembled to inform the level of malaria service disruption by country.<sup>2</sup> Trackers, other than those for campaign-type interventions, had important limitations related to periodicity, scope and reliability. In particular, information on disruptions of clinical management of malaria (diagnosis and treatment) was not adequately captured by all the trackers. Where attempts were made to capture such information, the responses were qualitative and difficult to validate. This exercise highlights the need to ensure that countries' health information systems can capture critical data elements related to service disruptions and mechanisms, and complement these with low-cost sentinel surveillance and rapid community surveys.

## 10.4 COUNTRY RESPONSES TO MITIGATE GLOBAL SERVICE DISRUPTIONS

Several malaria endemic countries with moderate or high transmission had plans to implement campaigns to distribute LLINs, IRS and SMC in 2020. The COVID-19 pandemic threatened the safe and effective delivery of these interventions. Faced with the possibility that most of the gains over the past 20 years could be reversed in a single year if major malaria intervention programmes

were disrupted (**Section 10.3.2**), many malaria endemic countries mounted an impressive response by adapting service delivery approaches while still adhering to the restrictions imposed by national attempts to curb the spread of SARS-CoV2 infections. The guidance provided by the WHO GMP (with support from partners) (148) coupled with documents

<sup>1</sup> RBM Partnership to End Malaria. *Best practices in mitigating the effect of COVID-19 on malaria at country and sub-regional level*. October 2020, report in preparation.

<sup>2</sup> RBM Country/Regional Support Partner Committee (CRSPC) tracker (150), Workstream 3 trackers (ITN, IRS and SMC), RBM MERG routine data tracker (151) and WHO essential health services survey (152).



developed by partners to support implementation, were critical in helping countries tailor their responses to the COVID-19 pandemic.

In their mitigation response, countries faced several challenges: lack of funds and delays in procurement of PPE; delays in procurement and delivery to country of adequate nets, insecticides, diagnostics and drugs because of global supply chain disruptions

(Sections 10.3.3–10.3.5); delays in shipping due to mobility restrictions; and the need to acquire high-level political support in an environment where most of the focus was on direct efforts to fight COVID-19.

A case study of Benin, as an example of a country adapting and maintaining malaria services during the COVID-19 pandemic, is presented in **Box 10.1**.

### **BOX 10.1.**

#### **Benin: Country example for sustaining malaria programming during COVID-19**

In March 2020, the first cases of COVID-19 were recorded in Benin, just as the country was planning its LLIN campaign. Following the WHO recommendation to continue with the implementation of malaria control interventions in the face of COVID-19 (148), and with strong support from the RBM Partnership, the Ministry of Health was authorized to continue with the implementation of the planned LLIN campaign. Working closely with the RBM Partnership through the Alliance for Malaria Prevention (AMP), Benin's National Malaria Control Programme reviewed and revised their distribution strategy to mitigate the risks of COVID-19 transmission<sup>a</sup> during the campaign. The AMP guidance for distribution of ITNs during COVID-19 transmission facilitated adaptation of the distribution strategy, with the adoption of a door-to-door distribution approach rather than distribution from a fixed point. The change in approach meant an increase in the number of days needed for community mobilization, modifications to briefings, training and supervision, plus the purchase of PPE. The Global Fund rapidly approved the release of funds from Benin's existing grant to cover any increased costs.

The strong leadership from the Government of Benin, the Ministry of Health and the NMP, and effective collaboration with international and implementing partners facilitated the door-to-door distribution of 7 638 192 nets in just 20 days, ensuring that Benin's population of 14 million were protected from malaria. Benin was the first country to proceed with its planned LLIN campaign in the face of COVID-19, providing a valuable "proof of concept" for other countries to follow. Other countries across Africa subsequently adopted the approach pioneered by Benin to ensure that life-saving mosquito nets were distributed.

Benin also successfully conducted IRS during the COVID-19 pandemic, spraying a total of 350 349 structures. With support from partners, the NMP updated the IRS strategy and training to include COVID-19 prevention measures. Additional protection measures were established, including increasing the number of handwashing stations for frontline workers and provision of additional vehicles to transport spray personnel in accordance with national travel recommendations. Measures were put in place for COVID-19 testing of spray personnel and for managing any suspected cases among the spray teams.

Benin also successfully completed four rounds of SMC in four health zones. With support from partners, the NMP adapted the SMC strategy to include COVID-19 prevention measures. Sensitization of communities and compliance with the government's protective measures (wearing a mask, using sanitizing gels and physical distancing), as well as limiting the number of participants in meetings and trainings, helped to build confidence in the community.

Finally, the country has worked to sustain case management of malaria during the COVID-19 pandemic. This has included ensuring sufficient supplies of essential malaria commodities (e.g. diagnostics and treatment) at health facility level.

Through strong leadership, and coordinated partnership, Benin has successfully implemented the LLIN, IRS and SMC campaigns planned for 2020, while working to sustain access to case management. All this has been achieved under the very difficult circumstances of the COVID-19 pandemic.

<sup>a</sup> <https://allianceformalariaprevention.com/wp-content/uploads/2020/10/Key-guidance-EN.pdf>

### 10.5 LEVELS OF SERVICE DISRUPTION BY COUNTRY AND IMPLICATIONS FOR DELIVERY OF INTERVENTIONS


According to available information,<sup>1</sup> all 31 countries (25 in sub-Saharan Africa) that had ITN campaigns planned in 2020 aimed to complete them by the end of the year. As of 23 November 2020, five countries had completed on time (within the planned period before the pandemic), seven had completed with moderate delays (within the second quarter of the original planned period), 12 had ongoing campaigns with moderate delays, and another seven had campaigns in progress but with major delays (beyond the second quarter of the original planned period). Of the 222 million ITNs

expected to be distributed in 2020, 105 million had been distributed by 23 November 2020. Of the 47 countries that planned IRS campaigns in 2020, 23 had completed them, with eight of those countries doing so with delays. Thirteen countries are on track to complete their IRS campaigns, six of them with delays. Eleven countries, eight of them in sub-Saharan Africa, were either off track or at risk of not completing their IRS campaigns. By the third week of November 2020, all countries that had planned SMC campaigns were on track to complete them, despite moderate delays in some areas.

<sup>1</sup> RBM Country/Regional Support Partner Committee (CRSPC) tracker (150) and Workstream 3 trackers (ITN, IRS and SMC).

**FIG. 10.4.**

**Results from WHO surveys on disruptions of malaria-related services during the COVID-19 pandemic: a) ANC services and b) diagnosis and treatment** No disruption (<5%); partial disruption (< 50%); severe disruption (>50%). Surveys were conducted in May–September 2020 Sources: WHO Integrated Health Services.




AFR: WHO African Region; AMR: WHO Region of the Americas; ANC: antenatal care; EMR: WHO Eastern Mediterranean Region; SEAR: WHO South-East Asia Region; WHO: World Health Organization; WPR: WHO Western Pacific Region.



Understanding the disruptions in malaria case management is difficult because it requires data from multiple household surveys of disruptions to treatment seeking for fevers, combined with information at health facility level about changes in patient caseloads. In addition, disruptions varied greatly within countries by geography and over time, making it difficult to draw conclusions from point-in-time data. These data should be combined with detailed country information on supply chains, and stockouts of diagnosis and treatment commodities in order to identify not only disruptions but also their potential causes and solutions. In the absence of such data, several proxies have been explored.

**Figure 10.4** shows responses from countries on the extent of disruptions of malaria diagnosis and treatment, collected through the WHO Essential Health Service pulse survey from mid-May to September 2020. The findings suggest that among the 64 malaria endemic countries that responded, 39 experienced partial disruption (of between 5% and 50%) of ANC services (**Fig. 10.4a**), and 37 experienced similar disruptions of malaria diagnosis and treatment (**Fig. 10.4b**). Djibouti reported severe disruptions of ANC services. This information is similar to that shown on other more recent surveys implemented by the Global Fund (153), suggesting that most malaria endemic countries surveyed have experienced at least moderate levels of

### b) Diagnosis and treatment



AFR: WHO African Region; AMR: WHO Region of the Americas; ANC: antenatal care; EMR: WHO Eastern Mediterranean Region; SEAR: WHO South-East Asia Region; WHO: World Health Organization; WPR: WHO Western Pacific Region.

# 10 Malaria response during the COVID-19 pandemic

disruption of malaria case management, of up to 50% based on the knowledge of the respondents.


Analysis of routine aggregate data, while potentially biased by many factors related to the quality of the surveillance system, may add value to our understanding of disruptions to clinical services.

**Fig. 10.5** shows monthly trends in all-cause outpatients in 2019, and up to June or September 2020 in the public health sector, for 23 countries in sub-Saharan Africa. Most of the countries show reductions in outpatient attendances from March 2020 onwards, compared with a similar period in 2019, suggesting a general decline in use of health services.

**FIG. 10.5.**

**Monthly trends in all-cause outpatients attendances in 23 countries in sub-Saharan Africa in 2019 and 2020**

Source: NMP reports.



NMP: national malaria programme.

Note: For Burkina Faso, monthly data from 2018 were used due to major disruptions of the surveillance system due to the 2019 health workers' strikes in 2019.




A similar analysis of malaria outpatient data shows that, despite decreasing overall attendance at public health facilities, malaria cases were generally higher in 2020 than in 2019 in 10 countries, and were lower in the remaining 14 countries (Fig. 10.6). There are several potential reasons for discordance in the trends in all-cause and malaria outpatient data, such as changes in

diagnostic practice or reporting of presumptively treated cases as parasitologically confirmed. However, a potential concern would be that there is increasing malaria transmission, whereby there is more malaria among those patients using services at a time when use of services has generally reduced due to COVID-19 disruptions.

**FIG. 10.6.**

**Monthly trends in malaria outpatients attendances in 24 countries in sub-Saharan Africa in 2019 and 2020**

Source: NMP reports.



NMP: national malaria programme.

Note: For Burkina Faso, monthly data from 2018 were used due to major disruptions of the surveillance system due to the 2019 health workers' strikes in 2019.




### 10.6 THE CONSEQUENCES OF SERVICE DISRUPTIONS DURING THE COVID-19 PANDEMIC

The analysis in this report of the consequences of disruption of services focuses on sub-Saharan Africa, a region that accounts for more than 90% of the burden of malaria morbidity and mortality. Within this region, the analysis further focuses on mortality because it is assumed that most of the prevention campaigns will be completed by the end of 2020, averting major increases in cases. Delays in the campaigns in 2020

have been included in the analysis of the effect of vector control coverage on infection and malaria cases. Different scenarios of disruptions of access to effective antimalarial treatment were applied to each country, to estimate the number of untreated cases. A uniform *P. falciparum* case fatality rate was then applied to the untreated cases, to estimate mortality by country (**Annex 1**).

**FIG. 10.7.** Estimated potential increase in malaria deaths in sub-Saharan Africa (excluding Botswana, Eswatini, Namibia and South Africa) corresponding to varying levels of disruptions of access to effective antimalarial treatment *Source: WHO estimates.*



WHO: World Health Organization.



The analysis shows that, even with completion of the prevention campaigns, relatively small disruptions in access to effective antimalarial treatments (similar to those suggested by the various trackers) can lead to considerable loss of life (**Fig. 10.7**). Thus, a disruption in access to treatment of 10% in sub-Saharan Africa is likely to lead to an estimated 19 000 additional deaths

among people of all ages. This is likely to increase to 28 000, 46 000 and 100 000 deaths if access is reduced by 15%, 25% and 50%, respectively.

Had the ITN, IRS and LLIN campaigns not happened in 2020 as planned, mortality would have increased several times more than currently projected.

# 11 Key results, context and conclusion

This concluding section of the *World malaria report 2020* highlights some of the progress made against malaria in the past 2 decades, calls out the major current challenges and threats (including the COVID-19 pandemic), and draws attention to opportunities for the global malaria community to work together to ensure even greater achievements in the next decade of the GTS.



## 11.1 KEY RESULTS

Following years of neglect, remarkable progress was made in malaria during the MDG era and that progress should be considered one of the first great public health success stories of the millennium. Despite modest levels of investment in research and development (R&D), new tools became available in the form of ITNs, ACTs and RDTs. New strategies to deploy existing tools were developed, including various forms of chemoprevention (e.g. IPTp, IPTi and SMC), the use of community health workers and greater engagement with the private sector.

A range of financing mechanisms were developed to augment the national investments of endemic countries: between 2000 and 2019, about US\$ 39 billion was invested in the fight against malaria, of which US\$ 26 billion represented funds from external donors (**Section 6**). These developments led to an unprecedented scale-up of effective malaria interventions (**Section 7**). Over 2.2 billion ITNs, 3.1 billion ACTs and 2.7 billion RDTs have been delivered to malaria endemic countries. In sub-Saharan Africa, between 2000 and 2019, the percentage of children aged under 5 years and of pregnant women sleeping under an ITN both increased from below 3% to over 50%. More than 21 million children aged under 5 years have received SMC, and about 23 million (62%) pregnant women received at least one dose of IPTp in 2019 alone. The percentage of children being diagnosed using a parasitological test increased from 14% before the large rollout of RDTs to, on average,

40% in the most recent household surveys conducted in sub-Saharan Africa.

By 2019, there were 229 million malaria cases and 409 000 deaths globally, reducing from 238 million and 736 000 since 2000, respectively. It is estimated that 1.5 billion malaria cases and 7.6 million deaths had been averted since 2000 (**Section 3**). Since 2000, 21 countries had achieved malaria free status or were certified by WHO as having interrupted malaria transmission (**Section 4**). Thirty-one and 35 countries were on target for the 2020 GTS morbidity and mortality reduction targets, respectively (**Section 8**). Each WHO region had shown reductions in malaria case incidence and mortality rates since 2000, and the entire WHO European Region had been free of malaria since 2015 (**Section 3**). Under the HBHI approach, the 11 highest burden countries globally had concluded an intensive initial exercise to use their local data to develop and implement evidence-based subnationally tailored malaria interventions plans (**Section 5**). Through support from the Global Fund and PMI, these countries are expected to receive more funding in the period 2020–2022 than in the preceding 3 years.

Despite the overall progress made in the first 15 years of this century, global trends in malaria case and mortality rates have been plateauing since 2015 (**Section 3**), particularly in the highest burden countries that account for most of the cases and deaths globally (**Section 5**).



## 11.2 THE ENABLING ENVIRONMENT AND THREATS TO THE MALARIA PROGRESS

The unprecedented investment in malaria and the scale-up of interventions coincided with a period of considerable demographic and socioeconomic change in malaria endemic countries. In sub-Saharan Africa, where over 90% of the malaria burden occurs, the population increased from 665 million in 2000 to 1.1 billion in 2019, and it is projected to rise to 1.5 billion by 2030 (154). The proportion of this population that resides in urban areas increased from 31% in 2000 to 41% in 2019, and is projected to increase to 47% by 2030. GDP growth has averaged 4% since 2000, with several countries exceeding an average of 5% in this period (155), and the percentage of the population considered poor (i.e. living on <US\$ 1.90 a day at 2011 international prices) reducing from 60% in 2000 to 40% in 2018 (156). The level of rural electrification rose from 11% to 32% of households, giving those households better economic opportunities, connectivity and access to information (157). The 11 million mobile cellular subscriptions in 2000 increased dramatically to 537 million subscriptions in 2019 (158). Major improvements in socioeconomic growth and development have also occurred in many malaria endemic countries outside sub-Saharan Africa (159). These factors have no doubt contributed to general improvements in health and – both directly and in combination with the massive scale-up of malaria interventions – to the progress made against malaria since 2000.

The plateauing of the burden of malaria at what is still a very high level is a wake-up call, drawing attention not only to the need to innovate against the vector and the parasite – by developing new tools, strategies and problem-solving approaches at the frontline of malaria control – but also to ensure that the global response evolves. Sustained, strengthened and coordinated investments and actions are needed to build on earlier successes.

The efficacy of most of the current malaria prevention tools remains modest. High levels of coverage and user

compliance remain challenging, and the different approaches are threatened by emerging resistance (Section 9). The spread of resistance to insecticides used in ITNs and IRS is extensive and, although the epidemiological impact of such resistance remains inconclusive, reinforces the need for vigilance and development of new insecticides (Section 9). The emerging spread of *pfhrp2* deletions means that the most widely used malaria diagnostic test is no longer reliable in most countries in the Horn of Africa, and this situation could spread rapidly to other countries. ACT resistance; it has not spread from the GMS to the rest of the world as was previously feared; nevertheless, it remains a threat to which WHO continues to pay attention.

Funding for malaria has plateaued since 2010 (Section 6) and, despite the welcome increase in Global Fund replenishment in 2019, per capita investments for populations at risk are unlikely to change greatly in the period 2020–2022. The 2019 malaria funding of about US\$ 3 billion is considerably below the US\$ 5.6 billion estimated as being needed to achieve the GTS targets. Despite impressive economic growth in malaria endemic countries, domestic funding for malaria has also stagnated over the past decade.

Inadequate funding and inefficiencies in service delivery systems have resulted in some people failing to access and use malaria interventions. In sub-Saharan Africa, the population sleeping under ITNs has remained similar to 2015 levels (and actually declined slightly between 2018 and 2019), with important inequities in several countries (Section 7.1). Nearly 30% of children with fever are still not receiving care and less than half of those who seek care are not diagnosed using a parasitological test (Section 7.5). A third of these children use private health facilities (Section 7), with households incurring expenses they can barely afford. This draws further attention to the importance of UHC and of ensuring that mechanisms exist to deliver interventions without creating financial hardship.


# 11 Key results, context and conclusion

The link between improving human development and reducing the burden of infectious diseases is strong (160). It is anticipated that as the world strives for a future without malaria, human development, in all its facets, will be one of the biggest drivers for this change (113). At the same time, reducing the burden of malaria through prevention and treatment is likely to contribute

to accelerated development. Currently, however, more than 80% of the burden of malaria is concentrated in countries with low human development indices (Fig. 11.1), assessed using dimensions of health, education and standard of living indicators (159), impairing the capacity and resilience of communities to respond to the burden of malaria.

**FIG. 11.1.**

**Distribution of malaria cases in 2019 by human development index in 2018** Sources: WHO estimates, UNDP.



UNDP: United Nations Development Programme; WHO: World Health Organization.




About 90% of the burden of malaria occurs in countries where health expenditure as a percentage of GDP is less than 7%, and 75% of the burden is in countries where health expenditure is less than 5% of GDP (Fig. 11.2). In these countries, more than 70% of funding for malaria is from external sources, mainly from the

Global Fund and PMI (Section 6). Among moderate to high transmission countries in sub-Saharan Africa, progress towards the target of 15% expenditure on health as a percentage of GDP by 2015 committed to by countries under the Abuja Declaration (1) remains elusive, with no country achieving it by 2017 (161).

**FIG. 11.2.**

**Distribution of malaria cases in 2019 by current health expenditure as a percentage of GDP in 2017** Sources: WHO estimates, World Bank.



GDP: gross domestic product; WHO: World Health Organization.


# 11 Key results, context and conclusion

There are no reliable data measuring the status of health system governance. **Fig. 11.3** presents the distribution of burden by level of general governance effectiveness, as analysed by the World Bank (162). The index of governance effectiveness reflects respondent perceptions of the quality of public services, the quality of the civil service and its degree of independence from political pressures, the quality of policy formulation and implementation, and the

credibility of the government's commitment to policies. Information on governance effectiveness for malaria endemic countries was extracted and countries were grouped into qualitative categories by government effectiveness as very low, low, moderate or high (**Fig. 11.3**). About 77% of all malaria case burden is accounted for by countries with very low or low governance effectiveness.

**FIG. 11.3.**

**Distribution of malaria cases in 2019 by category of governance effectiveness in 2019** Sources: WHO estimates, World Bank.



WHO: World Health Organization.




An analysis of the UHC service coverage index by country was undertaken by WHO for the period 2000–2017 (163). This index was computed using information on 16 tracer indicators across four service coverage categories: reproductive, maternal, newborn and child health; infectious diseases; noncommunicable diseases; and service capacity and access, and health security (164). The burden of malaria and access to malaria interventions were also included in the

composite index of effective service coverage. The potential circularity notwithstanding, there is a clear pattern in the relationship between the UHC service coverage index and malaria burden (Fig. 11.4). About 90% of the burden of malaria globally in 2019 was concentrated in countries that were classified as having a low UHC service coverage index (i.e. <50).

**FIG. 11.4.**

**Distribution of malaria cases in 2019 by category of UHC service coverage index in 2017** Sources: WHO estimates, World Bank.



UHC: universal health coverage; WHO: World Health Organization.



# 11 Key results, context and conclusion

Reliable health information is critical for developing sound strategic and operational plans, efficiently and equitably targeting resources and reliably measuring the impact of interventions (**Section 3, Section 5**). Considerable improvements have been made in recent years, building on the introduction of parasitological diagnosis, which have improved the value of the data on malaria cases, and the use of digital solutions (e.g. DHIS2), which in turn have improved data transmission, validation and analysis. In many moderate to high burden countries, especially in sub-Saharan Africa, the available routine data are increasing in volume, but there are still considerable issues with data quality. Consequently, for 30 countries in this region – which account for over 85% of the burden of malaria cases for this report – malaria case totals are computed using a method that derives case incidence from intermittent community parasite prevalence data (**Section 3, Annex 1**). Mortality estimation also relies on verbal autopsy data to define causes of death; however, such data have been shown to be unreliable in identifying malaria deaths (165). Facility-level electronic data entry is non-existent in most of the countries in sub-Saharan Africa, making data transmission and aggregation labour intensive, and increasing the likelihood of transcription errors and significant delays. These weaknesses have been most starkly demonstrated by the difficulties in tracking service disruptions during the COVID-19 pandemic (**Section 10**).

Over the past 2 decades, malaria endemic countries have also had to deal with numerous complex emergencies – both natural and human made – undermining progress in these countries and resulting in a heavy toll on already fragile health and livelihoods. As recently as 2018–2020, many high burden malaria endemic countries have been afflicted with major storms or flooding, including, for example, Burkina Faso, Cameroon, Central African Republic, Chad, Côte d'Ivoire, Ethiopia, Ghana, Kenya, Malawi, Mali,

Mauritania, Mozambique, Niger, Nigeria, Senegal, Sierra Leone, Somalia, South Sudan, Sudan, Uganda and the United Republic of Tanzania (166–168). Many countries are also dealing with local active conflicts (170) that limit the population's access to care, and the ability of government and stakeholders to reach people. In addition, frequent outbreaks and epidemics of non-malaria diseases in malaria endemic settings have resulted in major disruptions to malaria services (**Section 10**). Despite their frequency and impact, these emergencies are unpredictable; in fact, they are missing entirely from quantitative global projections of the future trajectory of malaria (113, 115).

Between 2007 and 2018, almost US\$ 7.3 billion was invested in basic research and product development for malaria, rising from about US\$ 500 million in 2007 to slightly over US\$ 650 million in 2018 (**Section 6**). A lot of knowledge has been generated and many tools are in the pipeline. However, progress against malaria in the past 2 decades has been delivered by the continued dependence of countries on a combination of several imperfect tools delivered to communities through relatively expensive mechanisms (**Section 7**), resulting in persistent gaps in coverage. Many of the tools currently in use were developed in the 1980s and 1990s. There have been progressive improvements, such as new ITNs/LLINs, new ACTs, and new formulations of existing ACTs and the advent of RDTs (an important innovation that enhances case management), the targeted use of ACTs the value of routine malaria case data. The next major innovation may be a malaria vaccine, introduced as part of routine control efforts. Pilot implementation of RTS,s/AS01 in three African countries started in 2019. In late 2021, WHO is expected to review evaluation data from the pilots together with the results of several studies conducted since 2015, and consider the advisability of broader use of this vaccine. This would open a new paradigm in the approach to malaria control.

## 11.3 CONSEQUENCES OF THE COVID-19 PANDEMIC

COVID-19 has exposed the fragility of today's society and systems, shaken the global economy and begun to reverse the progress made in reducing poverty and fighting disease (171). It is estimated that COVID-19 will push about 100 million people into extreme poverty in 2020 and will have a prolonged economic legacy (172). At the time of writing, almost 50 million cases of COVID-19 have been reported to WHO, and more than 1.2 million people have lost their lives. millions more are likely to have died due to disruption of essential health services.

Health sectors across the world are facing a triple challenge: minimizing the immediate health impact of

COVID-19, reducing disruption to other essential services and managing the health of their nation while reorienting their economies for recovery. The limited fiscal space in many parts of sub-Saharan Africa has compromised spending on COVID-19 and continues to threaten other health priorities. Early lockdown measures in many malaria endemic countries may have protected people from COVID-19, but they have also affected people's access to health care and other services. On the demand side, fewer patients are presenting to outpatient care (**Section 10**), fearing the risk of becoming infected with COVID-19, and hindered by lockdowns and lack of transport. On the supply side,



elective care has frequently been cancelled, and commodity supply chains both within and beyond malaria endemic countries have been disrupted. COVID-19 highlighted the severe shortages in the health workforce in LMICs, compromising clinical and social care and public health services. Health worker redeployment, fear of returning to work without PPE, sickness and death have further hampered service delivery (173).

The lack of infection prevention in facilities, including PPE, has had dire personal and public health consequences. A disproportionate number of health workers have been infected with COVID-19, compromising the capacity to deliver essential services, putting patients at risk of COVID-19 and deterring people from seeking care. Based on reports from key informants, the most frequently disrupted areas included routine immunization-outreach services (70%) and facility-based services (61%), non-communicable diseases diagnosis and treatment (69%), family planning and contraception (68%), treatment for mental health disorders (61%), and cancer diagnosis and treatment (55%) (152). Thirty-seven (58%) of 64

malaria endemic countries surveyed have also reported disruptions to malaria diagnosis and treatment (Section 10). Although disrupted or delayed, many of the campaigns for ITNs and SMC were conducted safely. However, the analysis suggests that even if malaria prevention campaigns are completed in 2020 as planned, disruptions to access to effective antimalarial treatment could lead to considerable loss of life (Section 10).

The pandemic is clearly a global crisis that requires a concerted global response. The sheer scale of the pandemic and the broader disruptions it has caused requires strong leadership and citizenship to chart a new way forward. In an interconnected world, this pandemic has highlighted the critical importance of global solidarity in addressing the divisions, fragilities and inequities that COVID-19 and other infectious diseases thrive upon. The ACT Accelerator (174) is a good example of the collective resolve necessary to rapidly develop quality assured vaccines, diagnostics and therapeutics, and to allocate them fairly. Building on the GTS principles, these positive lessons from COVID-19 need to be extended to the malaria response.

## 11.4 BUILDING A MORE PROSPEROUS FUTURE

The challenge of getting back on track during such difficult times is daunting, but there are reasons to be hopeful. Over the past 2 decades the malaria community has shown what it can do when faced with adversity. Looking forward, as we learn from COVID-19 and the early progress on HBHI, the principles outlined in the GTS become even more relevant for the challenges we are facing today.

### 11.4.1 Country ownership and leadership, with involvement and participation of communities, are essential to accelerating progress through a multisectoral approach

The major public health challenges, including malaria, require a whole of government, whole of society approach. Trusted, accountable national political leadership is essential, using the best knowledge and science to galvanize the many actors around a common narrative and unified response. Their political commitment will need to translate into resources and actions to ensure that all those in need have access to the appropriate mix of interventions for malaria prevention and quality health care, without financial hardship. As with other health priorities, this relies upon the inclusion and participation of many stakeholders, including the most vulnerable communities, women

and children. Empowered and incentivized individuals are at the heart of primary health care, as people and their communities are advocates for policies that promote and protect health and well-being, are co-developers of health and social services, and act as self-carers and caregivers to others (114).

### 11.4.2 Improved surveillance, monitoring and evaluation, as well as stratification by malaria disease burden, are required to optimize the implementation of malaria interventions

Effective and efficient malaria programming and the containment of outbreaks such as the COVID-19 pandemic rely on effective data and surveillance systems. Data and local intelligence are critical for adapting to constantly evolving local disease patterns, and for optimizing the choice and delivery of interventions. Data are also needed to ensure that no one is left behind, helping to identify the least served and to understand and overcome the barriers they face. This data-driven approach is at the heart of the HBHI approach and is applicable to all malaria endemic countries. As the COVID-19 pandemic takes a toll on global economies, the data-driven approach will be even more critical in achieving more with less.

# 11 Key results, context and conclusion

Bold actions are needed to ensure that surveillance systems are ready for efficient routine operations and future epidemics. Seven broad areas require investment: **i)** assessing the status of surveillance systems to understand bottlenecks and use evidence to guide investments in the system; **ii)** ensuring availability of parasitological tests in all health facilities and increasing adherence to test results; **iii)** moving away from aggregate tallying of cases by hand from registers to using personal electronic records in all malaria endemic countries, thus improving the efficiency, quality and value of surveillance – this will apply to the broader health information system, including the private sector, and will be achieved if gains in electrification, renewable energy, increased connectivity and reduced costs of computing hardware are optimized; **iv)** developing integrated databases that are governed by national authorities and analytical capacity at all programmatic levels, to ensure countries can use and act on their data; **v)** adapting surveillance systems and analytics to the changing socioeconomic and demographic environment – in particular, to respond to malaria in an increasingly urbanized population; **vi)** using the data to inform communities about the services that are available to them, their rights to access those services and the risks they are exposed to; and **vii)** enhancing innovation in the use of digital solutions, data science and genomics in malaria surveillance.

## **11.4.3 Equity in access to health services especially for the most vulnerable and hard-to-reach populations is essential**

All citizens, wherever malaria is present, should have access to quality services to prevent, diagnose and treat the disease without facing financial hardship. However, as this report documents, many people living in countries where malaria remains a major public health challenge still lack access to essential health services, and some people are still pushed into extreme poverty by paying the costs of malaria prevention and treatment. Well-functioning, resilient health systems based on primary health care are critical for progress towards the interrelated goals of health security and UHC. The global commitments on UHC made in September 2019, at the UN high-level meeting on UHC (116) need to be translated into resources for implementing high-impact health interventions to combat malaria and other diseases, protecting women's and children's health, and ensuring no one suffers financial hardship because they have had to pay for their health care. This will

require strengthening of integrated frontline delivery channels – primary care and emergency care, equipped with essential medicines and commodities to provide people with diagnosis and treatment when and wherever they need it. These platforms deliver benefits across a range of conditions and reap economies of scale.

## **11.4.4 Strengthen health workforce and malaria expert base**

In most countries where malaria is endemic, there is a chronic shortage of skilled health professionals. Robust expansion of malaria interventions requires significantly expanded human resource capacities at national, district and community levels, and the deployment of health workers to cater for remote and underserved populations. A strengthening of the workforce across a variety of technical and service delivery areas based on a sound analysis and national plan should be recognized as an essential part of health systems strengthening.

## **11.4.5 Innovation in tools and implementation approaches will enable countries to accelerate their progression along the path to elimination**

Continued investment in R&D is needed to develop the tools required to stay ahead of resistance and other effects of biological selection pressures. Periodic reviews of unmet public health needs and the types of products required to address those needs should be set alongside the development pipelines. This will make it easier to identify opportunities to intensify effort and investments to accelerate the availability of products where they are needed. Finance is required to generate solid evidence of the epidemiological and public health benefits of new interventions. Only with such information can programmes tailor the introduction of new technologies and be confident that they are maximizing the impact of available resources. Innovation is needed in the global financing architecture to incentivize R&D of products intended primarily for LMICs. Similarly, forethought is needed to avoid the bottlenecks that prevent production and delivery at scale of newly developed products. To keep product developers and donors engaged, it is essential to show a path to market and public health impact. For too long, operational and implementation research has been too neglected. Additional investments can help to unlock the full potential impact of the tools that are already available.



## 11.5 CONCLUDING REMARKS

The malaria problem is evolving, dynamic and diverse. The lessons of the past 2 decades show that success in malaria is possible when the world pulls together. They also show that there are enormous data, biological, political, governance, socioeconomic and financial challenges. It is now understood that a one-size-fits-all approach cannot be expected to address the problem in any one country. Compounding the challenges are weak coordination structures that do not always put the national decision-making processes at the core of public health governance. Both within endemic countries and across the broader malaria architecture, we need to take stock of and improve our approaches to responding to malaria.

The GTS principles, agreed by Member States and the wider malaria community, remain as relevant to the future of malaria control and elimination as they have been in the past. WHO promotes the GTS milestones as staging posts that help us to reflect on our past – and plan our future – contributions to the malaria response. The GTS recognizes that there are needs specific to malaria while acknowledging that success is only achievable through strong primary health care. Now that we are at the first milestone in the GTS, we must commit to doing a better job or delivering on its promise through our collective resolve.

# References

1. The Abuja Declaration and the plan of action. An extract from the African Summit on Roll Back Malaria, Abuja, 25 April 2000. Geneva: World Health Organization; 2000 ([http://whqlibdoc.who.int/hq/2003/WHO\\_CDS\\_RBM\\_2003.46.pdf](http://whqlibdoc.who.int/hq/2003/WHO_CDS_RBM_2003.46.pdf)).
2. Millennium Development Goals [website]. New York City: United Nations (<https://www.un.org/millenniumgoals/>).
3. Sustainable Development Goals: 17 goals to transform our world [website]. United Nations 2015 (<https://www.un.org/sustainabledevelopment/sustainable-development-goals>).
4. Global technical strategy for malaria 2016–2030 (last update: 22 January 2016). Geneva: World Health Organization; 2015 ([http://apps.who.int/iris/bitstream/10665/176712/1/9789241564991\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/176712/1/9789241564991_eng.pdf)).
5. Roll Back Malaria Partnership Secretariat. Action and investment to defeat malaria 2016–2030. For a malaria-free world. Geneva: World Health Organization; 2015 ([https://endmalaria.org/sites/default/files/RBM\\_AIM\\_Report\\_0.pdf](https://endmalaria.org/sites/default/files/RBM_AIM_Report_0.pdf)).
6. Declarations and decisions adopted by the Thirty-Third Assembly of Heads of States and Government: Assembly of Heads of State and Government, Thirty-Third Ordinary Session, June. OAU Secretariat; 1997 ([https://au.int/sites/default/files/decisions/9542-1997\\_ahg\\_dec\\_120-123\\_xxxiii\\_e.pdf](https://au.int/sites/default/files/decisions/9542-1997_ahg_dec_120-123_xxxiii_e.pdf)).
7. Kidson C. Global malaria challenge: the Amsterdam summit. *Southeast Asian J Trop Med Public Health*. 1992;23(4):635–40 (<https://pubmed.ncbi.nlm.nih.gov/1298069/>).
8. Snow RW, Amratia P, Kabaria CW, Noor AM, Marsh K. The changing limits and incidence of malaria in Africa: 1939–2009. *Adv Parasitol*. 2012;78:169–262 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3521063/>).
9. Alonso P, Noor AM. The global fight against malaria is at crossroads. *Lancet*. 2017;390(10112):2532–4 (<https://pubmed.ncbi.nlm.nih.gov/29195688>).
10. Trape J-F. The public health impact of chloroquine resistance in Africa. *Am J Trop Med Hyg*. 2001;64(1-2 Suppl):12–7 (<https://pubmed.ncbi.nlm.nih.gov/11425173/>).
11. Ministerial Conference on Malaria: report by the Director-General. Geneva: World Health Organization; 1992 ([https://apps.who.int/iris/bitstream/handle/10665/171067/EB91\\_4\\_eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/171067/EB91_4_eng.pdf)).
12. Forty-ninth World Health Assembly: Resolutions and decisions annexes. Geneva: World Health Organization; 1996 ([https://apps.who.int/iris/bitstream/handle/10665/178941/WHA49\\_1996-REC-1\\_eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/178941/WHA49_1996-REC-1_eng.pdf)).
13. Accelerated implementation of malaria control. Geneva: Roll Back Malaria, World Health Organization; 2000 ([https://apps.who.int/iris/bitstream/handle/10665/66635/WHO\\_CDS\\_RBM\\_2000.27.pdf](https://apps.who.int/iris/bitstream/handle/10665/66635/WHO_CDS_RBM_2000.27.pdf)).
14. Hedding A, Keusch GT, Davies CS. The multilateral initiative on malaria: past, present, and future. *Am J Trop Med Hyg*. 2004;71:279–82 ([https://www.ncbi.nlm.nih.gov/books/NBK3741/pdf/Bookshelf\\_NBK3741.pdf](https://www.ncbi.nlm.nih.gov/books/NBK3741/pdf/Bookshelf_NBK3741.pdf)).
15. Nabarro DN, Tayler EM. The “Roll Back Malaria” campaign. *Science*. 1998;280(5372):2067–8 (<https://pubmed.ncbi.nlm.nih.gov/9669961/>).
16. TDR (Special Programme for Research and Training in Tropical Diseases) [website]. Geneva: United Nations Children’s Fund, United Nations Development Programme, World Bank, World Health Organization; 2020 (<https://www.who.int/tdr/about/en/>).
17. Alonso PL, Lindsay SW, Armstrong J, De Francisco A, Shenton F, Greenwood B et al. The effect of insecticide-treated bed nets on mortality of Gambian children. *Lancet*. 1991;337(8756):1499–502 (<https://pubmed.ncbi.nlm.nih.gov/1675368/>).
18. Snow R, Lindsay S, Hayes R, Greenwood B. Permethrin-treated bed nets (mosquito nets) prevent malaria in Gambian children. *Trans R Soc Trop Med Hyg*. 1988;82(6):838–42.
19. White N. Antimalarial drug resistance and combination chemotherapy. *Philos Trans R Soc Lond B Biol Sci*. 1999;354(1384):739–49 (<https://pubmed.ncbi.nlm.nih.gov/10365399/>).
20. White N. Delaying antimalarial drug resistance with combination chemotherapy. *Parassitologia*. 1999;41(1–3):301–8 (<https://pubmed.ncbi.nlm.nih.gov/10697872/>).
21. Schellenberg JRA, Maokola W, Shirima K, Manzi F, Mrisho M, Mushi A et al. Cluster-randomized study of intermittent preventive treatment for malaria in infants (IPTi) in southern Tanzania: evaluation of impact on survival. *Malar J*. 2011;10(1):387.
22. Greenwood BM, Greenwood AM, Bradley AK, Snow RW, Byass P, et al. (1988) Comparison of two strategies for control of malaria within a primary health care programme in The Gambia. *Lancet*. 1: 1121–1127.
23. Steketee RW, Wirima JJ, Hightower AW, Slutsker L, Heymann DL, Breman JG. The effect of malaria and malaria prevention in pregnancy on offspring birthweight, prematurity, and intrauterine growth retardation in rural Malawi. *Am J Trop Med Hyg*. 1996;55(1\_Suppl):33–41.
24. Alonso PL, Sacarlal J, Aponte JJ, Leach A, Macete E, Milman J et al. Efficacy of the RTS,S/AS02A vaccine against *Plasmodium falciparum* infection and disease in young African children: randomised controlled trial. *Lancet*. 2004;364(9443):1411–20 (<https://pubmed.ncbi.nlm.nih.gov/15488216/>).
25. INDEPTH Network. <http://www.indepth-network.org/about-us>

26. Guidelines for the treatment of malaria, third edition. Geneva: World Health Organization; 2015 (<https://www.who.int/publications/item/9789241549127>).
27. Safety of pyrethroid-treated mosquito nets: WHOPES (WHO Pesticide Evaluation Scheme) fact sheet (WHO/CDS/CPE/WHOPES/99.5). Geneva: World Health Organization/Department of Communicable Disease Prevention, Control and Eradication; 1999 ([https://www.who.int/whopes/resources/who\\_cds\\_cpe\\_whopes\\_99.5/en/](https://www.who.int/whopes/resources/who_cds_cpe_whopes_99.5/en/)).
28. Lengeler C. Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Sys Rev.* 2004;(2):CD000363 (<https://www.ncbi.nlm.nih.gov/pubmed/15106149>).
29. Chandre F, Darrier F, Manga L, Akogbeto M, Faye O, Mouchet J et al. Status of pyrethroid resistance in *Anopheles gambiae sensu lato*. *Bull World Health Org.* 1999;77(3):230 (<https://pubmed.ncbi.nlm.nih.gov/10212513/>).
30. Sibley CH, Hyde JE, Sims PF, Plowe CV, Kublin JG, Mberu EK et al. Pyrimethamine-sulfadoxine resistance in *Plasmodium falciparum*: what next? *Trends Parasitol.* 2001;17(12):570–1 (<https://pubmed.ncbi.nlm.nih.gov/11756042/>).
31. Developing antimalarials to save lives [website]. Geneva: Medicines for Malaria Venture (<https://www.mmv.org/about-us/what-we-do/our-history>).
32. G8 Summit [website]. Okinawa, 2000 (<http://www.g8.utoronto.ca/summit/2000okinawa/>).
33. Gallup JL, Sachs JD. The economic burden of malaria. *Am J Trop Med Hyg.* 2001;64:85–96 (<https://www.ncbi.nlm.nih.gov/books/NBK2624/>).
34. MDG 6: combat HIV/AIDS, malaria and other diseases. Geneva: World Health Organization; 2014 ([https://www.who.int/topics/millennium\\_development\\_goals/diseases/en/](https://www.who.int/topics/millennium_development_goals/diseases/en/)).
35. The use of antimalarial drugs: report of a WHO informal consultation. Geneva: World Health Organization; 2000 ([https://apps.who.int/iris/bitstream/handle/10665/66961/WHO\\_CDS\\_RBM\\_2001.33.pdf](https://apps.who.int/iris/bitstream/handle/10665/66961/WHO_CDS_RBM_2001.33.pdf)).
36. Global report on antimalarial efficacy and drug resistance: 2000–2010 [website]. Geneva: World Health Organization; 2010 ([http://whqlibdoc.who.int/publications/2010/9789241500470\\_eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241500470_eng.pdf)).
37. New perspectives: malaria diagnosis – report of a joint WHO/USAID informal consultation 25–27 October 1999. Geneva: World Health Organization; 2000 ([https://apps.who.int/iris/bitstream/handle/10665/66321/WHO\\_CDS\\_RBM\\_2000.14.pdf](https://apps.who.int/iris/bitstream/handle/10665/66321/WHO_CDS_RBM_2000.14.pdf)).
38. Bill & Melinda Gates Foundation: who we are [website]. Seattle: Bill & Melinda Gates Foundation; 2020 (<https://www.gatesfoundation.org/Who-We-Are>).
39. EDCTP: about us [website]. The Hague: European & Developing Countries Clinical Trials Partnership (EDCTP); 2020 (<https://www.edctp.org/get-know-us/#>).
40. EDCTP malaria portfolio: large investments in treatment and vaccines [website]. The Hague: European & Developing Countries Clinical Trials Partnership (EDCTP); 2019 (<http://www.edctp.org/news/edctp-malaria-portfolio-large-investments-in-treatment-and-vaccine/>).
41. FIND – At a glance [website]. Geneva: Foundation for Innovative New Diagnostics (FIND); 2020 (<https://www.finddx.org/at-a-glance/>).
42. The Global Fund overview [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria (<https://www.theglobalfund.org/en/overview/>).
43. Bruno J-M, Feacham R, Godal T, Nchinda T, Ogilvie B, Mons B et al. The spirit of Dakar: a call for action on malaria (correspondence). *Nature.* 1997;386:541 (<https://www.nature.com/articles/386541a0.pdf>).
44. World malaria report 2005 – UNICEF’s role. Geneva: United Nations Children’s Fund; 2005 (<https://www.unicef.org/media/files/UNICEFinmalaria.pdf>).
45. Malaria [website]. Geneva: United Nations Children’s Fund; 2020 (<https://www.unicef.org/topics/malaria>).
46. Scaling up insecticide-treated netting programmes in Africa (2nd edition, Revision 23). Geneva: Roll Back Malaria, World Health Organization; 2005 (<https://www.who.int/publications/m/item/scaling-up-insecticide-treated-netting-programmes-in-africa>).
47. The Africa malaria report 2003. Geneva: World Health Organization, United Nations Children’s Fund; 2003 ([https://apps.who.int/iris/bitstream/handle/10665/67869/WHO\\_CDS\\_MAL\\_2003.1093.pdf](https://apps.who.int/iris/bitstream/handle/10665/67869/WHO_CDS_MAL_2003.1093.pdf)).
48. World malaria report 2005. Geneva: World Health Organization; 2005 ([http://whqlibdoc.who.int/publications/2005/9241593199\\_eng.pdf](http://whqlibdoc.who.int/publications/2005/9241593199_eng.pdf)).
49. The Global Fund’s first replenishment 2006–2007 [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; 2005 ([https://www.theglobalfund.org/media/1489/replenishment\\_pledges2006and2007\\_report\\_en.pdf](https://www.theglobalfund.org/media/1489/replenishment_pledges2006and2007_report_en.pdf)).
50. World malaria report 2008. Geneva: World Health Organization; 2008 ([http://whqlibdoc.who.int/publications/2008/9789241563697\\_eng.pdf](http://whqlibdoc.who.int/publications/2008/9789241563697_eng.pdf)).
51. PMI: About page [website]. U.S. President’s Malaria Initiative (PMI) (<https://www.pmi.gov/about>).
52. PMI. 14th annual report to congress. Washington, DC: U.S. President’s Malaria Initiative (PMI); 2020 (<https://www.pmi.gov/docs/default-source/default-document-library/pmi-reports/2020-pmi-fourteenth-annual-report.pdf>).
53. The Roll Back Malaria strategy for improving access to treatment through home management of malaria. Geneva: World Health Organization; 2005 ([http://whqlibdoc.who.int/hq/2005/WHO\\_HTM\\_MAL\\_2005.1101.pdf](http://whqlibdoc.who.int/hq/2005/WHO_HTM_MAL_2005.1101.pdf)).
54. Guidelines for the treatment of malaria. Geneva: World Health Organization; 2006 ([https://archives.who.int/publications/2006/9241546948\\_eng.pdf](https://archives.who.int/publications/2006/9241546948_eng.pdf)).
55. Malaria indicator surveys [website]. Rockville, MD: RBM Partnership to End Malaria (<https://www.malariasurveys.org/>).

56. The DHS Program: demographic and health surveys [website]. Washington, DC: United States Agency for International Development (<https://dhsprogram.com/>).
57. MICS surveys [website]. Geneva: United Nations Children's Fund (<https://mics.unicef.org/surveys>).
58. About IVCC [website]. Liverpool, United Kingdom: IVCC (<https://www.ivcc.com/about/>).
59. Market access [website]. Liverpool, United Kingdom: IVCC; 2020 (<https://www.ivcc.com/market-access/>).
60. Unitaid constitution. Geneva: Unitaid; 2018 (<https://unitaid.org/assets/UNITAID-Constitution-revised-version-15-June-2018.pdf>).
61. Unitaid's malaria portfolio is expanding to support world elimination targets [website]. Geneva: Unitaid; 2018 (<https://unitaid.org/news-blog/unitaids-malaria-portfolio-is-expanding-to-support-world-elimination-targets/#en>).
62. Willey BA, Paintain LS, Mangham L, Car J, Schellenberg JA. Strategies for delivering insecticide-treated nets at scale for malaria control: a systematic review. *Bull World Health Organ.* 2012;90:672–84 (<https://www.who.int/bulletin/volumes/90/9/11-094771/en/>).
63. Noor AM, Amin AA, Akhwale WS, Snow RW. Increasing coverage and decreasing inequity in insecticide-treated bed net use among rural Kenyan children. *PLoS Med.* 2007;4(8):e255 (<https://pubmed.ncbi.nlm.nih.gov/17713981/>).
64. Insecticide-treated mosquito nets: a WHO position statement [website]. Geneva: World Health Organization (<http://medi-guide.meditool.cn/ympdf/FBACA171-AF95-DCFA-673E-3FA3AFEB79AE.pdf>).
65. Bill and Melinda Gates call for new global commitment to chart a course for malaria eradication. Seattle: Bill & Melinda Gates Foundation; 2007 (<https://www.gatesfoundation.org/Media-Center/Press-Releases/2007/10/Chart-a-Course-for-Malaria-Eradication>).
66. Global Malaria Action Plan for a malaria-free world. Roll Back Malaria Partnership; 2008 (<https://www.afro.who.int/sites/default/files/2017-06/Gmapfull.pdf>).
67. Malaria [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria (<https://www.theglobalfund.org/en/malaria/>).
68. Talisuna AO, Adibaku S, Amojah CN, Amofah GK, Aubyn V, Doodoo A et al. The affordable medicines facility-malaria – a success in peril. *Malar J.* 2012;11(1):370 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3506461/pdf/1475-2875-11-370.pdf>).
69. AMFm Independent Evaluation Team. Independent Evaluation of Phase 1 of the Affordable Medicines Facility - malaria (AMFm), Multi-country independent evaluation report: final report. Calverton, Maryland and London: ICF International and London School of Hygiene and Tropical Medicine; 2012.
70. Independent evaluation of the affordable medicines facility – malaria (AMFm) Phase 1. Calverton, Maryland and London: International Classification of Functioning, Disability and Health (ICF) & London School of Hygiene and Tropical Medicine; 2012 ([https://www.theglobalfund.org/media/3011/terg\\_evaluation2013-2014thematicreviewamfm2012iephase1\\_report\\_en.pdf](https://www.theglobalfund.org/media/3011/terg_evaluation2013-2014thematicreviewamfm2012iephase1_report_en.pdf)).
71. Gamboa D, Ho MF, Bendezu J, Torres K, Chiodini PL, Barnwell JW et al. A large proportion of *P. falciparum* isolates in the Amazon region of Peru lack pfhrp2 and pfhrp3: implications for malaria rapid diagnostic tests. *PLoS One.* 2010;5(1):e8091 (<https://www.ncbi.nlm.nih.gov/pubmed/20111602>).
72. About ALMA [website]. African Leaders Malaria Alliance (ALMA); 2020 (<https://alma2030.org/about-alma/>).
73. Intermittent preventive treatment for infants using sulfadoxine-pyrimethamine (SP-IPTi) for malaria control in Africa: implementation field guide. Geneva: World Health Organization; 2011 ([http://whqlibdoc.who.int/hq/2011/WHO\\_IVB\\_11.07\\_eng.pdf](http://whqlibdoc.who.int/hq/2011/WHO_IVB_11.07_eng.pdf)).
74. Rao M. The International Centers of Excellence for Malaria Research. *Am J Trop Med Hyg.* 2015;93(3 Suppl):1–4 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4574268/>).
75. World malaria report 2011. Geneva: World Health Organization; 2011 ([http://apps.who.int/iris/bitstream/10665/44792/2/9789241564403\\_eng\\_full.pdf](http://apps.who.int/iris/bitstream/10665/44792/2/9789241564403_eng_full.pdf)).
76. Malaria Policy Advisory Committee [website]. Geneva: World Health Organization (<https://www.who.int/groups/malaria-policy-advisory-committee>).
77. World malaria report 2019. Geneva: World Health Organization; 2019 (<https://www.who.int/publications/i/item/9789241565721>).
78. Dondorp AM, Yeung S, White L, Nguon C, Day NP, Socheat D et al. Artemisinin resistance: current status and scenarios for containment. *Nat Rev Microbiol.* 2010;8(4):272–80 (<https://pubmed.ncbi.nlm.nih.gov/20208550/>).
79. Global Plan for Artemisinin Resistance Containment (GPARC). Geneva: Roll Back Malaria, World Health Organization; 2011 ([https://apps.who.int/iris/bitstream/handle/10665/44482/9789241500838\\_eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/44482/9789241500838_eng.pdf)).
80. Countries of the Greater Mekong zero in on falciparum malaria [website]. Geneva: World Health Organization; 2019 ([https://www.who.int/malaria/areas/greater\\_mekong/en/](https://www.who.int/malaria/areas/greater_mekong/en/)).
81. Focus on Regional Artemisinin-resistance Initiative (RAI). Geneva: The Global Fund to Fight AIDS, Tuberculosis and Malaria; 2019 ([https://www.theglobalfund.org/media/6509/publication\\_regionalartemisininresistanceinitiative\\_focuson\\_en.pdf](https://www.theglobalfund.org/media/6509/publication_regionalartemisininresistanceinitiative_focuson_en.pdf)).
82. The Global Fund Strategy 2012–2016: investing for impact. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; 2011 ([https://www.theglobalfund.org/media/1179/core\\_globalfundstrategy2012-2016\\_strategy\\_en.pdf](https://www.theglobalfund.org/media/1179/core_globalfundstrategy2012-2016_strategy_en.pdf)).
83. Board approves integration of AMFm into core global fund grant processes. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; 2012 (<https://www.theglobalfund.org/en/news/2012-11-15-board-approves-integration-of-amfm-into-core-global-fund-grant-processes/>).
84. Seasonal malaria chemoprevention (SMC) [website]. Geneva: World Health Organization; 2017 (<https://www.who.int/teams/global-malaria-programme/prevention/preventive-chemotherapies>).
85. Unitaid. The project [website]. 2018 (<https://www.access-smc.org/pages/access-smc/context>).
86. Overview: funding model [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; (<https://www.theglobalfund.org/en/funding-model/>).

87. PMI. Seasonal malaria chemoprevention. Washington, DC: U.S. President's Malaria Initiative (PMI); 2020 (<https://www.pmi.gov/how-we-work/technical-areas/seasonal-malaria-chemoprevention>).
88. Global plan for insecticide resistance management in malaria vectors [website]. Geneva: World Health Organization; 2012 ([https://www.who.int/iris/bitstream/10665/44846/1/9789241564472\\_eng.pdf](https://www.who.int/iris/bitstream/10665/44846/1/9789241564472_eng.pdf)).
89. Faulde MK, Rueda LM, Khairah BA. First record of the Asian malaria vector *Anopheles stephensi* and its possible role in the resurgence of malaria in Djibouti, Horn of Africa. *Acta Trop.* 2014;139:39–43.
90. Ashine T, Teka H, Esayas E, Messenger LA, Chali W, Meerstein-Kessel L et al. *Anopheles stephensi* as an emerging malaria vector in the Horn of Africa with high susceptibility to Ethiopian *Plasmodium vivax* and *Plasmodium falciparum* isolates. *bioRxiv.* 2020.
91. malERA – A research agenda for malaria eradication [website]. San Francisco: PLOS (<https://collections.plos.org/malera>).
92. Malaria elimination [website]. *The Lancet*; 2010 (<https://www.thelancet.com/malaria-elimination>).
93. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature.* 2015;526(7572):207–11 (<https://pubmed.ncbi.nlm.nih.gov/26375008>).
94. Hay SI, Guerra CA, Gething PW, Patil AP, Tatem AJ, Noor AM et al. A world malaria map: *Plasmodium falciparum* endemicity in 2007. *PLoS Med.* 2009;6(3):e1000048 (<https://pubmed.ncbi.nlm.nih.gov/19323591/>).
95. Noor AM, Kinyoki DK, Mundia CW, Kabaria CW, Mutua JW, Alegana VA et al. The changing risk of *Plasmodium falciparum* malaria infection in Africa: 2000–10: a spatial and temporal analysis of transmission intensity. *Lancet.* 2014;383(9930):1739–47 (<https://pubmed.ncbi.nlm.nih.gov/24559537/>).
96. Snow RW, Sartorius B, Kyalo D, Maina J, Amratia P, Mundia CW et al. The prevalence of *Plasmodium falciparum* in sub-Saharan Africa since 1900. *Nature.* 2017;550(7677):515–8 (<https://pubmed.ncbi.nlm.nih.gov/29019978/>).
97. World malaria report 2015. Geneva: World Health Organization; 2015 ([http://apps.who.int/iris/bitstream/10665/200018/1/9789241565158\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/200018/1/9789241565158_eng.pdf)).
98. Sustainable Development Goals: the 17 goals [website]. New York City: United Nations, Department of Economic and Social Affairs – Sustainable Development; (<https://sdgs.un.org/goals>).
99. Global indicator framework for the Sustainable Development Goals and targets of the 2030 Agenda for Sustainable Development. New York City: United Nations; 2020 ([https://unstats.un.org/sdgs/indicators/Global%20Indicator%20Framework%20after%202020%20review\\_Eng.pdf](https://unstats.un.org/sdgs/indicators/Global%20Indicator%20Framework%20after%202020%20review_Eng.pdf)).
100. Malaria threats map [website]. Geneva: World Health Organization (<https://apps.who.int/malaria/maps/threats/>).
101. Strategy for malaria elimination in the Greater Mekong subregion: 2015–2030. Geneva: World Health Organization; 2015 (<https://apps.who.int/iris/handle/10665/208203>).
102. RBM Partnership. RBM Partnership to End Malaria overview [website]. 2020 (<https://endmalaria.org/about-us/overview>).
103. The Global Fund Strategy 2017–2022: investing to end epidemics. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; 2016 (<https://www.theglobalfund.org/en/strategy/>).
104. Policy Cures Research: G-FINDER data portal [website]. Sydney, Australia: Policy Cures Research (<https://gfinderdata.policycuresresearch.org>).
105. Eliminating malaria: 21 countries, a common goal [website]. Geneva: World Health Organization (<https://www.who.int/teams/global-malaria-programme/elimination/e-2020-initiative-of-malaria-eliminating-countries>).
106. Ministerial Declaration on Accelerating and Sustaining Malaria Elimination in the South-East Asia Region [website]. New Delhi: WHO South-East Asia Region; 2017 ([http://origin.searo.who.int/entity/malaria/sea\\_declaration\\_malaria\\_elimination.pdf](http://origin.searo.who.int/entity/malaria/sea_declaration_malaria_elimination.pdf)).
107. World malaria report 2017. Geneva: World Health Organization; 2017 (<https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2017>).
108. High burden to high impact: a targeted malaria response. Geneva: World Health Organization; 2019 (<http://apps.who.int/iris/bitstream/handle/10665/275868/WHO-CDS-GMP-2018.25-eng.pdf>).
109. Compendium of WHO malaria guidance – prevention, diagnosis, treatment, surveillance and elimination. Geneva: World Health Organization; 2019 (<https://apps.who.int/iris/bitstream/handle/10665/312082/WHO-CDS-GMP-2019.03-eng.pdf>).
110. WHO technical brief for countries preparing malaria funding requests for the Global Fund (2020–2022). Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/9789240004139>).
111. Sixth Global Fund's Replenishment Conference [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; 2019 (<https://www.theglobalfund.org/en/specials/2019-10-09-global-fund-sixth-replenishment-conference/>).
112. The use of artesunate-pyronaridine for the treatment of uncomplicated malaria. Geneva: World Health Organization; 2019 (<https://www.who.int/publications/i/item/use-of-artesunate-pyronaridine-for-the-treatment-of-uncomplicated-malaria>).
113. Malaria eradication: benefits, future scenarios and feasibility. Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/malaria-eradication-benefits-future-scenarios-feasibility>).
114. Astana Declaration on Primary Health Care. Geneva: World Health Organization; 2018 (<https://www.who.int/teams/primary-health-care/conference/declaration>).
115. Report overview and technical briefs [website]. The Lancet Commission on Malaria Eradication (<http://malariaeradicationcommission.com/report/report-overview-and-technical-briefs>).
116. Universal health coverage: moving together to build a healthier world – UN high-level meeting on universal health coverage, 23 September 2019, New York [website]. New York City: United Nations (<https://www.un.org/pga/73/event/universal-health-coverage/>).



117. The potential impact of health service disruptions on the burden of malaria. A modelling analysis for countries in sub-Saharan Africa. Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/the-potential-impact-of-health-service-disruptions-on-the-burden-of-malaria>).
118. Guyatt HL, Snow RW. Impact of malaria during pregnancy on low birth weight in sub-Saharan Africa. *Clin Microbiol Rev.* 2004;17(4):760–9 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC523568/>).
119. Guyatt HL, Snow RW. Malaria in pregnancy as an indirect cause of infant mortality in sub-Saharan Africa. *Trans R Soc Trop Med Hyg.* 2001;95(6):569–76 (<https://academic.oup.com/trstmh/article-abstract/95/6/569/1905057?redirectedFrom=fulltext>).
120. Malaria in pregnant women [website]. Geneva: World Health Organization; 2017 ([https://www.who.int/malaria/areas/high\\_risk\\_groups/pregnancy/en/](https://www.who.int/malaria/areas/high_risk_groups/pregnancy/en/)).
121. Walker PG, ter Kuile FO, Garske T, Menendez C, Ghani AC. Estimated risk of placental infection and low birthweight attributable to *Plasmodium falciparum* malaria in Africa in 2010: a modelling study. *Lancet Glob Health.* 2014;2(8):e460–e7 (<https://pubmed.ncbi.nlm.nih.gov/25103519/>).
122. Walker PG, Griffin JT, Cairns M, Rogerson SJ, Van Eijk AM, Ter Kuile F et al. A model of parity-dependent immunity to placental malaria. *Nat Commun.* 2013;4(1):1–11 (<https://www.nature.com/articles/ncomms2605>).
123. Total fertility rate (live births per woman) [website]. United Nations Statistics Division (<http://data.un.org/Data.aspx?d=PopDiv&f=variableID%3A54>).
124. Global call to boost access to lifesaving malaria treatment for pregnant women in Africa. Geneva: RBM Partnership to End Malaria; 2020 (<https://endmalaria.org/news/global-call-boost-access-lifesaving-malaria-treatment-pregnant-women-africa>).
125. Zero Malaria Starts with Me [website]. RBM Partnership to End Malaria and the African Union Commission (<https://zeromalaria.africa/>).
126. Funding model: allocation [website]. Global Fund to Fight AIDS, Tuberculosis and Malaria; 2019 (<https://www.theglobalfund.org/en/funding-model/before-applying/allocation/>).
127. Mass action for a malaria free Uganda – handbook for leaders. Government of Uganda; 2020 (<https://www.afro.who.int/publications/mass-action-malaria-free-uganda-handbook-leaders>).
128. Improving WHO’s malaria policy-making and dissemination processes. Geneva: World Health Organization; 2020 (<https://www.who.int/teams/global-malaria-programme/policy-making-process>).
129. Meeting report of the WHO Evidence Review Group on malaria burden estimation methods [website]. Geneva: World Health Organization; 2018 (<https://www.who.int/publications/m/item/WHO-CDS-GMP-MPAC-2018.10>).
130. Statistics on international development: final UK aid spend 2019. United Kingdom: Foreign, Commonwealth & Development Office; 2019 (<https://www.gov.uk/government/statistics/statistics-on-international-development-final-uk-aid-spend-2019>).
131. Malaria Atlas Project [website]. (<https://malariaatlas.org>).
132. False-negative RDT results and implications of new reports of *P. falciparum* histidine-rich protein 2/3 gene deletions. Geneva: World Health Organization; 2017 (<https://apps.who.int/iris/bitstream/10665/258972/1/WHO-HTM-GMP-2017.18-eng.pdf>).
133. Thomson R, Parr JB, Cheng Q, Chenet S, Perkins M, Cunningham J. Prevalence of *Plasmodium falciparum* lacking histidine-rich proteins 2 and 3: a systematic review. *Bull World Health Organ.* 2020;98(8):558 (<http://dx.doi.org/10.2471/BLT.20.250621>).
134. Template protocol for therapeutic efficacy studies. Geneva: World Health Organization; 2019 (<https://www.who.int/teams/global-malaria-programme/case-management/drug-efficacy-and-resistance/tools-for-monitoring-antimalarial-drug-efficacy>).
135. Malaria surveillance, monitoring & evaluation: a reference manual. Geneva: World Health Organization; 2018 (<http://apps.who.int/iris/bitstream/handle/10665/272284/9789241565578-eng.pdf>).
136. Insecticide resistance [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/teams/global-malaria-programme/prevention/insecticide-resistance>).
137. Test procedures for insecticide resistance monitoring in malaria vector mosquitoes (second edition). Geneva: World Health Organization; 2018 (<http://apps.who.int/iris/bitstream/10665/250677/1/9789241511575-eng.pdf>).
138. Insecticide resistance in malaria vectors [website]. Geneva: World Health Organization (<https://www.who.int/teams/global-malaria-programme/prevention/insecticide-resistance/insecticide-resistance-in-malaria-vectors>).
139. DHIS2-based entomology and vector control data collection and collation tools [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/teams/global-malaria-programme/prevention/vector-control/dhis-data-collection-and-collation-tools>).
140. Framework for a national plan for monitoring and management of insecticide resistance in malaria vectors. Geneva: World Health Organization; 2017 (<https://apps.who.int/iris/bitstream/handle/10665/254916/9789241512138-eng.pdf>).
141. Timeline: WHO’s COVID-19 response [website]. Geneva: World Health Organization; (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/interactive-timeline#event-5>).
142. International Health Regulations (2005). Geneva: World Health Organization; 2014 (<https://www.who.int/publications/i/item/9789241580496>).
143. Coronavirus disease (COVID-19) dashboard [website]. Geneva: World Health Organization; 2020 (<https://covid19.who.int/>).
144. Q&As on COVID-19 and related health topics [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/question-and-answers-hub>).
145. Massinga Loembe M, Tshangela A, Salyer SJ, Varma JK, Ouma AEO, Nkengasong JN. COVID-19 in Africa: the spread and response. *Nat. Med.* 2020;26(7):999–1003 (<https://pubmed.ncbi.nlm.nih.gov/32528154/>).
146. COVID-19: examining theories for Africa’s low death rates [website]. Australia: The Conversation; (<https://theconversation.com/covid-19-examining-theories-for-africas-low-death-rates-147393>).

147. WHO urges countries to ensure the continuity of malaria services in the context of the COVID-19 pandemic [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/news/item/25-03-2020-who-urges-countries-to-ensure-the-continuity-of-malaria-services-in-the-context-of-the-covid-19-pandemic>).
148. Tailoring malaria interventions in the COVID-19 response. Geneva: World Health Organization; 2020 (<https://www.who.int/docs/default-source/documents/publications/gmp/tailoring-malaria-interventions-covid-19.pdf>).
149. COVID-19 response mechanism [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria (<https://www.theglobalfund.org/en/covid-19/response-mechanism/>).
150. Country/Regional Support Partner Committee (CRSPC) [website]. United Nations Office for Project Services, RBM Partnership to End Malaria (<https://endmalaria.org/about-us-governance-partner-committees/countryregional-support-partner-committee-crspc>).
151. Monitoring and evaluation [website]. RBM Partnership to End Malaria; 2020 (<https://endmalaria.org/our-work-working-groups/monitoring-and-evaluation>).
152. Pulse survey on continuity of essential health services during the COVID-19 pandemic: interim report, 27 August 2020 [website]. World Health Organization; 2020 ([https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS\\_continuity-survey-2020.1](https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS_continuity-survey-2020.1)).
153. Data updates [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria; 2020 (<https://www.theglobalfund.org/en/updates/data/>).
154. World population prospects [website]. New York City: United Nations; 2020 (<https://population.un.org/wpp/>).
155. GDP per capita (current US%) [website]. Washington DC: World Bank; 2020 (<https://data.worldbank.org/indicator/NY.GDP.PCAP.CD>).
156. Poverty [website]. Washington DC: World Bank; 2020 (<https://data.worldbank.org/topic/11>).
157. Access to electricity, rural (% of rural population) [website]. Washington DC: World Bank; 2020 (<https://data.worldbank.org/indicator/EG.ELC.ACCS.RU.ZS>).
158. Mobile cellular subscriptions (per 100 people) [website]. Washington DC: World Bank; 2020 (<https://data.worldbank.org/indicator/IT.CEL.SETS.P2>).
159. Human development index (HDI) [website]. Geneva: United Nations Development Programme (UNDP); 2020 (<http://www.hdr.undp.org/en/content/human-development-index-hdi>).
160. WHO guide to identifying the economic consequences of disease and injury. Geneva: World Health Organization; 2009 ([https://www.who.int/choice/publications/d\\_economic\\_impact\\_guide.pdf](https://www.who.int/choice/publications/d_economic_impact_guide.pdf)).
161. Current health expenditure (% of GDP) [website]. Washington DC: World Bank; 2020 (<https://data.worldbank.org/indicator/SH.XPD.CHEX.GD.ZS>).
162. Kaufmann D, Kraay A, Mastruzzi M. The worldwide governance indicators: methodology and analytical issues: World Bank Policy Research Working Paper No. 5430. Washington, DC: The World Bank; 2010 ([https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=1682130](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=1682130)).
163. The Global Health Observatory: UHC service coverage index [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/data/gho/indicator-metadata-registry/imr-details/4834>).
164. Hogan D, Hosseinpoor, AR, Boerma T. Technical note: developing an index for the coverage of essential health services. Geneva: World Health Organization; 2005 ([https://www.who.int/healthinfo/universal\\_health\\_coverage/UHC\\_WHS2016\\_TechnicalNote\\_May2016.pdf](https://www.who.int/healthinfo/universal_health_coverage/UHC_WHS2016_TechnicalNote_May2016.pdf)).
165. Snow R, Winstanley M, Marsh V, Newton C, Waruiru C, Mwangi I et al. Childhood deaths in Africa: uses and limitations of verbal autopsies. *Lancet*. 1992;340(8815):351–5.
166. Information Bulletin Africa: Floods. Geneva: International Federation of the Red Cross; 2020 ([https://reliefweb.int/sites/reliefweb.int/files/resources/IB%20Africa%20Region\\_Floods.pdf](https://reliefweb.int/sites/reliefweb.int/files/resources/IB%20Africa%20Region_Floods.pdf)).
167. West and Central Africa: Weekly regional humanitarian snapshot [website]. United Nations Office for the Coordination of Humanitarian Affairs (OCHA); 2020 (<https://reliefweb.int/sites/reliefweb.int/files/resources/External%20Weekly%2011-17%20August%202020.pdf>).
168. Africa under water: heightened “mega” crises [website]. United Nations Office for the Coordination of Humanitarian Affairs (OCHA); 2020 (<https://unocha.exposure.co/africa-under-water-heightened-mega-crises>).
169. West Africa – floods hit Burkina Faso and Northern Ghana [website]. Floodlist; 2020 (<http://floodlist.com/africa/west-africa-burkinafaso-ghana-september-2020>).
170. UCDP dataset download center [website]. Uppsala: Uppsala Conflict Data Program; 2020 (<https://ucdp.uu.se/downloads/>).
171. Global economic prospects [website]. Washington DC: World Bank; 2020 (<https://www.worldbank.org/en/publication/global-economic-prospects>).
172. A UN framework for the immediate socio-economic response to COVID-19. Geneva: United Nations; 2020 (<https://unsdg.un.org/resources/un-framework-immediate-socio-economic-response-covid-19>).
173. Talisuna A, Yoti Z, Lee C, et al. The urgent need for a global commitment to protect healthcare workers. *BMJ Global Health* 2020;5:e004077. doi:10.1136/bmjgh-2020-004077
174. ACT Accelerator: an economic investment case & financing requirements [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/an-economic-investment-case-financing-requirements>).



# Annexes

## **Annex 1 - Data sources and methods**

### **Annex 2 - Regional profiles**

- > A. WHO African Region
  - a. West Africa
  - b. Central Africa
  - c. Countries with high transmission in East and Southern Africa
  - d. Countries with low transmission in East and Southern Africa
- > B. WHO Region of the Americas
- > C. WHO Eastern Mediterranean Region
- > D. WHO South-East Asia Region
- > E. WHO Western Pacific Region

### **Annex 3 - Data tables**

- > A. Policy adoption, 2019
- > B. Antimalarial drug policy, 2019
- > C. Funding for malaria control, 2017–2019
- > D. Commodities distribution and coverage, 2017–2019
- > Ea. Household survey results, 2015–2019, compiled through STATcompiler
- > Eb. Household survey results, 2015–2019, compiled through WHO calculations
- > F. Population denominator for case incidence and mortality rate, and estimated malaria cases and deaths, 2000–2019
- > G. Population denominator for case incidence and mortality rate, and reported malaria cases by place of care, 2019
- > H. Reported malaria cases by method of confirmation, 2010–2019
- > I. Reported malaria cases by species, 2010–2019
- > J. Reported malaria deaths, 2010–2019

## Annex 1 – Data sources and methods

### Table 2.1. GTS: global targets for 2030 and milestones for 2020 and 2025

Targets and milestones are as described in the *Global technical strategy for malaria 2016–2030 (GTS) (1)* and *Action and investment to defeat malaria 2016–2030 (AIM) (2)*.

### Fig. 2.1. Key milestones in the fight against malaria in the past 2 decades

An overview presentation of key milestones over the past 2 decades in the fight against malaria. Information was obtained from published and grey literature. Relevant original information sources are provided in the reference list.

### Fig. 3.1. Countries with indigenous cases in 2000 and their status by 2019

Data on the number of indigenous cases (an indicator of whether countries are endemic for malaria) were as reported to the World Health Organization (WHO) by national malaria programmes (NMPs). Countries with 3 consecutive years of zero indigenous cases are considered to have eliminated malaria.

### Table 3.1. Global estimated malaria cases and deaths, 2000–2019

#### a) Global estimated malaria cases

The number of malaria cases was estimated by one of the two methods described below.

#### Method 1

Method 1 was used for countries and areas outside Africa, and for low-transmission countries and areas in Africa: Afghanistan, Bangladesh, Bolivia (Plurinational State of), Botswana, Brazil, Cambodia, Colombia, Dominican Republic, Eritrea, Ethiopia, French Guiana, Gambia, Guatemala, Guyana, Haiti, Honduras, India, Indonesia, Lao People's Democratic Republic, Madagascar, Mauritania, Myanmar, Namibia, Nepal, Nicaragua, Pakistan, Panama, Papua New Guinea, Peru, Philippines, Rwanda, Senegal, Solomon Islands, Timor-Leste, Vanuatu, Venezuela (Bolivarian Republic of), Viet Nam, Yemen and Zimbabwe.

Estimates were made by adjusting the number of reported malaria cases for completeness of reporting, the likelihood that cases were parasite positive, and the extent of health service use. The procedure, which is described in the *World malaria report 2008 (3)*, combines data reported by NMPs (i.e. reported cases, reporting completeness and likelihood that cases are parasite positive) with data obtained from nationally representative household surveys on health service use. Briefly:

$$T = (a + (c \times e))/d \times (1+f/g+(1-g-f)/2/g)$$

where:

a is malaria cases confirmed in public sector

b is suspected cases tested

c is presumed cases (not tested but treated as malaria)

d is reporting completeness

e is test positivity rate (malaria positive fraction) = a/b

f is fraction seeking treatment in private sector

g is fraction seeking treatment in public sector

No treatment seeking factor: (1-g-f)

Cases in public sector: (a + (c x e))/d

Cases in private sector: (a + (c x e))/d x f/g

To estimate the uncertainty around the number of cases, the *test positivity rate* was assumed to have a normal distribution centred on the test positivity rate value and standard deviation – defined as  $0.244 \times f^{0.5547}$ , and truncated to be in the range 0, 1. *Reporting completeness (d)*, when reported as a range or below 80%, was assumed to have one of three distributions, depending on the value reported by the NMP. If the value was greater than 80%, the distribution was assumed to be triangular, with limits of 0.8 and 1.0, and the peak at 0.8. If the value was greater than 50% but less than 80%, the distribution was assumed to be rectangular, with limits of 0.5 and 0.8. Finally, if the value was lower than 50%, the distribution was assumed to be triangular, with limits of 0 and 0.5, and the peak at 0.5 (4). If the reporting completeness was reported as a value and was greater than 80%, a beta distribution was assumed with a mean value of the reported value (maximum of 95%) and confidence intervals (CIs) of 5% around the mean value. The fraction of children brought for care in the public sector and in the private sector was assumed to have a beta distribution, with the mean value being the estimated value in the survey and the standard deviation calculated from the range of the estimated 95% CIs. The fraction of children not brought for care was assumed to have a rectangular distribution, with the lower limit being 0 and the upper limit calculated as 1 minus the proportion that were brought for care in the public and private sectors. The three distributions (fraction seeking treatment in public sector, fraction seeking treatment in private sector only and fraction not seeking treatment) were constrained to add up to 1.

Values for the fractions seeking care were linearly interpolated between the years that had a survey, and were extrapolated for the years before the first or after the last survey. Missing values for the distributions were imputed in a similar way or, if there was no value for any year in the country or area, were imputed as a mixture of the distribution of the region for that year. CIs were obtained from 10 000 draws of the convoluted distributions. The data were analysed using R statistical software (5).

For India, the values were obtained at subnational level using the same methodology, but adjusting the private sector for an additional factor because of the active case detection, estimated as the ratio of the test positivity rate in active case detection over the test positivity rate for passive case detection. This factor was assumed to have a normal distribution, with mean value and standard deviation calculated from the values reported in 2010.

No adjustment for private sector treatment seeking was made for the following countries and areas, because they report cases from the private and public sector together: Bangladesh, Bolivia (Plurinational State of), Botswana, Brazil, Colombia, Dominican Republic, French Guiana, Guatemala, Guyana, Haiti, Honduras, Myanmar (since 2013), Nicaragua, Panama, Peru, Rwanda, Senegal (70% of private sector reported together with public sector in 2018) and Venezuela (Bolivarian Republic of).

## Method 2

Method 2 was used for high-transmission countries in Africa and for countries in the WHO Eastern Mediterranean Region in which the quality of surveillance data did not permit a robust estimate from the number of reported cases: Angola, Benin, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad, Congo, Côte d'Ivoire, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Ghana, Guinea, Guinea-Bissau, Kenya, Liberia, Malawi, Mali, Mozambique, Niger, Nigeria, Sierra Leone, Somalia, South Sudan, Sudan, Togo, Uganda, United Republic of Tanzania and Zambia. In this method, estimates of the number of malaria cases were derived from information on parasite prevalence obtained from household surveys.

First, data on parasite prevalence from nearly 60 000 survey records were assembled within a spatio-temporal Bayesian geostatistical model, along with environmental and sociodemographic covariates, and data distribution on interventions such as insecticide-treated mosquito nets (ITNs), antimalarial drugs and indoor residual spraying (IRS) (6). The geospatial model enabled predictions of *Plasmodium falciparum* prevalence in children aged 2–10 years, at a resolution of  $5 \times 5$  km<sup>2</sup>, throughout all malaria endemic African countries for each year from 2000 to 2019. Second, an ensemble model was developed to predict malaria incidence as a function of parasite prevalence (7). The model was then applied to the estimated parasite prevalence in order to obtain estimates of the malaria case incidence at  $5 \times 5$  km<sup>2</sup> resolution for each year from 2000 to 2019.<sup>1</sup> Data for each  $5 \times 5$  km<sup>2</sup> area were then aggregated within country and regional boundaries, to obtain both national and regional estimates of malaria cases (9).

## Other methods

For most of the elimination countries and countries at the stage of prevention of reintroduction, the number of indigenous cases registered by NMPs are reported without further adjustments. The countries in this category were Algeria, Argentina, Armenia, Azerbaijan, Belize, Bhutan, Cabo Verde, China, Comoros, Costa Rica, Democratic People's Republic of Korea, Djibouti, Ecuador, Egypt, El Salvador, Eswatini, Georgia, Iran (Islamic Republic of), Iraq, Kazakhstan, Kyrgyzstan, Malaysia, Mexico, Morocco, Oman, Paraguay, Republic of Korea, Sao Tome and Principe, Saudi Arabia, South Africa, Sri Lanka, Suriname, Syrian Arab Republic, Tajikistan, Thailand, Turkey, Turkmenistan, United Arab Emirates and Uzbekistan.

For some years, information was not available or was not of sufficient quality to be used. For those countries, the number of cases was imputed from other years where the quality of the data was better, adjusting for population growth, as follows: for Afghanistan, values for 2000 and 2001 were imputed from 2002–2003; and for Bangladesh, values for 2001–2005 were imputed from 2006–2008. For Ethiopia, the values for 2000–2019 were taken from a mixed distribution between values from Method 1 and Method 2 (50% from each method). For Gambia, values for 2000–2010 were imputed from 2011–2013; for Haiti, values for 2000–2005, 2009 and 2010 were imputed from 2006–2008; for Indonesia, values for 2000–2003 and 2007–2009 were imputed from 2004–2006; for Mauritania, values for 2000–2010 were imputed from a mixture of Method 1 and Method 2, starting with 100% values from Method 2 for 2001 and 2002, and increasing to 90% values from Method 1 in 2010. For Myanmar, values for 2000–2005 were imputed from 2007–2009; for Namibia, values for 2000 were imputed from 2001–2003, and for 2012 from 2011 and 2013. For Pakistan, values for 2000 were imputed from 2001–2003; for Papua New Guinea, values for 2012 were imputed from 2009–2011. For Rwanda, values for 2000–2006 were imputed from a mixture of Method 1 and Method 2, starting with 100% values from Method 2 in 2000, with that percentage decreasing to 10% in 2006. For Senegal, values for 2000–2006 were imputed from a mixture of Method 1 and Method 2, with 90% of Method 2 in 2000, decreasing to 10% of Method 2 in 2006. For Thailand, values for 2000 were imputed from 2001–2003; for Timor-Leste, values for 2000–2001 were imputed from 2002–2004; and for Zimbabwe, values for 2000–2006 were imputed from 2007–2009. For Burkina Faso, Mali and Niger, values for 2000–2019 were imputed from the estimated series in the *World malaria report 2019* (10). For Côte d'Ivoire and Uganda, values were obtained from a combination of the values from the *World malaria report 2019* (10) and the current series, extrapolated as the trend from the most

<sup>1</sup> See the Malaria Atlas Project website for methods on the development of maps (8).

## Annex 1 – Data sources and methods

recent years for the 2019 estimation for Côte d'Ivoire and from the last incidence value for Uganda.

The number of malaria cases caused by *P. vivax* in each country was estimated by multiplying the country's reported proportion of *P. vivax* cases (computed as  $1 - P. falciparum$ ) by the total number of estimated cases for the country. For countries where the estimated proportion was not 0 or 1, the proportion of *P. falciparum* cases was assumed to have a beta distribution and was estimated from the proportion of *P. falciparum* cases reported by NMPs.

To transform malaria cases into incidence, an estimate of population at risk was used. The proportion of the population at high, low or no risk of malaria was provided by NMPs. This was applied to United Nations (UN) population estimates, to compute the number of people at risk of malaria.

### b) Global estimated malaria deaths

Numbers of malaria deaths were estimated using methods from Category 1, 2 or 3, as outlined below.

#### Category 1 method

The Category 1 method was used for low-transmission countries and areas, both within and outside Africa: Afghanistan, Bangladesh, Bolivia (Plurinational State of), Botswana, Cambodia, Comoros, Djibouti, Eritrea, Eswatini, Ethiopia, French Guiana, Guatemala, Guyana, Haiti, Honduras, India, Indonesia, Lao People's Democratic Republic, Madagascar, Myanmar, Namibia, Nepal, Pakistan, Papua New Guinea, Peru, Philippines, Solomon Islands, Somalia, Sudan, Timor-Leste, Vanuatu, Venezuela (Bolivarian Republic of), Viet Nam, Yemen and Zimbabwe. A case fatality rate of 0.256% was applied to the estimated number of *P. falciparum* cases, which represents the average of case fatality rates reported in the literature (11–13) and rates from unpublished data from Indonesia, 2004–2009.<sup>1</sup> The proportion of deaths then follows a categorical distribution of 0.01%, 0.19%, 0.30%, 0.38% and 0.40%, each one with equal probability.

A case fatality rate of 0.0375% was applied to the estimated number of *P. vivax* cases, representing the midpoint of the range of case fatality rates reported in a study by Douglas et al. (14), following a rectangular distribution between 0.012% and 0.063%. Following the nonlinear association explained for the Category 2 method below, the proportion of deaths in children aged under 5 years was estimated as:

$$\text{Proportion of deaths}_{\text{under 5}} = -0.2288 \times \text{Mortality}_{\text{overall}}^2 + 0.823 \times \text{Mortality}_{\text{overall}} + 0.2239$$

where  $\text{Mortality}_{\text{overall}}$  is the number of estimated deaths over the estimated population at risk per 1000 (see **Annex 3.F** for national estimates of population at risk).

#### Category 2 method

The Category 2 method was used for countries in Africa with a high proportion of deaths due to malaria: Angola, Benin, Burkina Faso, Burundi, Cameroon, Central African Republic, Chad, Congo, Côte d'Ivoire, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Gambia, Ghana, Guinea, Guinea-Bissau, Kenya, Liberia, Malawi, Mali, Mauritania, Mozambique, Niger, Nigeria, Rwanda, Senegal, Sierra Leone, South Sudan, Togo, Uganda, United Republic of Tanzania and Zambia.

In this method, child malaria deaths were estimated using a verbal autopsy multicausal model that was developed by the WHO Maternal and Child Health Epidemiology Estimation Group (MCEE) to estimate causes of death in children aged 1–59 months (15). Mortality estimates (and 95% CI) were derived for seven causes of post-neonatal death (pneumonia, diarrhoea, malaria, meningitis, injuries, pertussis and other disorders), four causes arising in the neonatal period (prematurity, birth asphyxia and trauma, sepsis, and other conditions of the neonate), and other causes (e.g. malnutrition). Deaths due to measles, unknown causes and HIV/AIDS were estimated separately. The resulting cause-specific estimates were adjusted, country by country, to fit the estimated mortality envelope of 1–59 months (excluding HIV/AIDS and measles deaths) for corresponding years. Estimated prevalence of malaria parasites (see methods notes for **Table 3.1**) was used as a covariate within the model. It was assumed that the number of deaths follows a rectangular distribution, with limits being the estimated 95% CI. The malaria mortality rate in children aged under 5 years estimated with this method was then used to infer malaria-specific mortality in those aged over 5 years, using the relationship between levels of malaria mortality in a series of age groups and the intensity of malaria transmission (16), and assuming a nonlinear association between under-5-years mortality and over-5-years mortality, as follows:

$$\text{Proportion of deaths}_{\text{over 5}} = -0.293 \times \text{Mortality}_{\text{under 5}}^2 + 0.8918 \times \text{Mortality}_{\text{under 5}} + 0.2896$$

where  $\text{Mortality}_{\text{under 5}}$  is estimated from the number of deaths from the MCEE model over the population at risk per 1000.

#### Category 3 method

For the Category 3 method, the number of indigenous malaria deaths registered by NMPs is reported without further adjustments. This category is used in the following countries: Algeria, Argentina, Armenia, Azerbaijan, Belize, Bhutan, Brazil, Cabo Verde, China, Colombia, Costa Rica, Democratic People's Republic of Korea, Dominican Republic, Ecuador, Egypt, El Salvador, Georgia, Iran (Islamic Republic of), Iraq, Kazakhstan, Kyrgyzstan,

<sup>1</sup> Dr Ric Price, Menzies School of Health Research, Australia, personal communication (November 2014).

Malaysia, Mexico, Morocco, Nicaragua, Oman, Panama, Paraguay, Republic of Korea, Sao Tome and Principe, Saudi Arabia, South Africa, Sri Lanka, Suriname, Syrian Arab Republic, Tajikistan, Thailand, Turkey, Turkmenistan, United Arab Emirates and Uzbekistan.

**Fig. 3.2. Global trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019, c) distribution of malaria cases and d) deaths by country, 2019**

See methods notes for Table 3.1.

**Table 3.2. Estimated malaria cases and deaths in the WHO African Region, 2000–2019**

See methods notes for Table 3.1.

**Fig. 3.3. Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO African Region, 2019**

See methods notes for Table 3.1.

**Table 3.3. Estimated malaria cases and deaths in the WHO South-East Asia Region, 2000–2019**

See methods notes for Table 3.1.

**Fig. 3.4. Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO South-East Asia Region, 2019**

See methods notes for Table 3.1.

**Table 3.4. Estimated malaria cases and deaths in the WHO Eastern Mediterranean Region, 2000–2019**

See methods notes for Table 3.1.

**Fig. 3.5. Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO Eastern Mediterranean Region, 2019**

See methods notes for Table 3.1.

**Table 3.5. Estimated malaria cases and deaths in the WHO Western Pacific Region, 2000–2019**

See methods notes for Table 3.1.

**Fig. 3.6. Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO Western Pacific Region, 2019**

See methods notes for Table 3.1.

**Table 3.6. Estimated malaria cases and deaths in the WHO Region of the Americas, 2000–2019**

See methods notes for Table 3.1.

**Fig. 3.7. Trends in a) malaria case incidence rate (cases per 1000 population at risk), b) mortality rate (deaths per 100 000 population at risk), 2000–2019 and c) malaria cases by country in the WHO Region of the Americas, 2019**

See methods notes for Table 3.1.

**Fig. 3.8. Cumulative number of cases and deaths averted globally and by WHO region, 2000–2019**

See methods for information on estimation of cases and deaths. Estimated cases and deaths averted were computed by comparing current estimates for each year since 2000 with estimates computed by holding the 2000 case incidence and mortality rates constant throughout the period 2000–2019.

**Fig. 3.9. Percentage of a) cases and b) deaths averted by WHO region, 2000–2019**

See methods for information on estimation of cases and deaths. See Fig. 3.8 for methods to estimate cases and deaths averted. The percentage of cases and deaths averted was estimated using overall global cases and deaths averted as denominator, and regional cases and deaths averted as numerator.

**Fig. 3.10. Estimated prevalence of exposure to malaria infection during pregnancy, overall and by subregion in 2019, in moderate to high transmission countries in the WHO African Region**

Estimates of malaria-exposed pregnancies and preventable malaria-attributable low birthweight (LBW) deliveries in the absence of pregnancy-specific malaria prevention (i.e. long-lasting insecticidal net [LLIN] delivery based on intermittent preventive treatment in pregnancy [IPTp] or antenatal care [ANC]) were obtained using a model of the relationship between these outcomes with slide microscopy prevalence in the general population and age- and gravidity-specific fertility patterns. This model



## Annex 1 – Data sources and methods

was developed by fitting an established model of the relationship between malaria transmission and malaria infection by age (17) to patterns of infection in placental histology (18) and attributable LBW risk by gravidity in the absence of IPTp or other effective chemoprevention (19). The model was run across a 0.2 degree (5 km<sup>2</sup>) longitude/latitude grid for 100 realizations of the Malaria Atlas Project (MAP) joint posterior estimated slide prevalence in children aged 2–10 years in 2018 (9). Country-specific, age-specific or gravidity-specific fertility rates, stratified by urban rural status, were obtained from demographic health surveys (DHS) and malaria indicator surveys (MIS), where such surveys had been carried out since 2014 and were available from the DHS program website (20). Countries where surveys were not available were allocated fertility patterns from a survey from another country, matched on the basis of total fertility rate (21) and geography. Fertility patterns of individual women within simulations at each grid-point were simulated according to the proportion of women estimated to be living in urban or rural locations. Urban or rural attribution at a 1 km<sup>2</sup> scale was conducted based on WorldPop 1 km<sup>2</sup> population estimates from 2018 (22) and an urban/rural threshold of 386/km<sup>2</sup> (23); the estimates were then aggregated to the 0.2 degree (5 km<sup>2</sup>) resolution of the MAP surfaces. This provided a risk of malaria infection and malaria-attributable LBW in the absence of prevention, along with a modelled per capita pregnancy rate for each grid-point, which was aggregated to country level (using WorldPop population estimates) to provide a per pregnancy risk of malaria infection and per livebirth estimate of malaria-attributable LBW in the absence of prevention. These were then multiplied by country-level estimates of pregnancies and estimates of LBW in 2019 (Fig. 3.11).

### **Fig. 3.11. Estimated number of low birthweights due to exposure to malaria infection during pregnancy overall and by subregion in 2019, in moderate to high transmission countries in sub-Saharan Africa**

Methods for estimating malaria infection in pregnancy and malaria-attributable LBWs are described in Walker et al. (19). Numbers of pregnancies were estimated from the latest UN population-estimated number of births and adjusted for the rate of abortion, miscarriage and stillbirths (24, 25). The underlying *P. falciparum* parasite prevalence estimates were from the updated MAP series, using methods described in Bhatt et al. (2015) (9).

### **Fig. 3.12. Estimated number of low birthweights averted if current levels of IPTp coverage are maintained and the additional number averted if coverage of first dose of IPTp was optimized to match levels of coverage of first ANC visit in 2019, in moderate to high transmission countries in the WHO African Region**

Efficacy of IPTp was modelled as a per-sulfadoxine-pyrimethamine (SP) dose reduction in the attributable risk of LBW and fitted to data from trials of IPTp-SP efficacy before the implementation of the intervention as policy; thus, they reflect impact on drug-sensitive parasites, with our central estimate being based on an assumed malaria-attributable LBW fraction of 40% within these trials. The modelling produced estimates of 48.5%, 73.5% and 86.3% efficacy in preventing malaria-attributable LBW for women receiving one, two or three doses of SP through IPTp, respectively. See the methods for Fig. 3.11.

### **Fig. 4.1. Number of countries that were malaria endemic in 2000, with fewer than 10, 100, 1000 and 10 000 indigenous malaria cases between 2000 and 2019**

The figure is based on the countries where malaria was endemic in 2000 and had cases of malaria in 2000. The number of estimated cases was tabulated.

### **Table 4.1. Countries eliminating malaria since 2000**

Countries are shown by the year in which they attained zero indigenous cases for 3 consecutive years, according to reports submitted by NMPs.

### **Table 4.2. Number of indigenous malaria cases in E-2020 countries, 2010–2019**

Data were derived from NMP reports.

### **Fig. 4.2. Total malaria and *P. falciparum* cases in the GMS, 2000–2019**

Data were derived from NMP reports to the Greater Mekong subregion (GMS) Malaria Elimination Database (MEDB).

### **Fig. 4.3. Regional map of malaria incidence in the GMS by area, 2012–2019**

Data were derived from NMP reports to the GMS MEDB.

### **Fig. 5.1. HBHI: a targeted malaria response to get countries back on track to achieve the GTS 2025 milestones**

This figure on high burden high impact (HBHI) was taken from a recent WHO publication (26).

### **Table 5.1. HBHI Response Element 2: work areas and status update**

The work areas shown in the table were developed by WHO and the RBM Partnership in consultation with countries and stakeholders as part of the HBHI response (26).

### **Fig. 5.2. Example of subnational tailoring of malaria intervention mixes and their projected impacts implemented as part of the HBHI response (in Nigeria)**

This is an example from Nigeria of analysis resulting from the HBHI Response Element 2 support involving subnational tailoring of malaria interventions using granular data on epidemiology and other factors developed by GMP. A mathematical model developed by the Institute for Disease Modeling<sup>1</sup> was used to assess the impact of various scenarios, with different mixes of interventions.

### **Fig. 5.3. Estimated malaria a) cases, b) cases per 1000 population at risk, c) deaths and d) deaths per 100 000 population at risk, 2018 and 2019, in HBHI countries**

See methods notes for Table 3.1.

### **Table 5.2. Comparisons of estimated malaria cases (millions) using the parasite rate-to-incidence model (Annex 1) and the reported data from the routine public health sector in high-burden countries of the WHO African Region, 2019**

See methods notes for Table 3.1. The analysis compares, for 10 HBHI countries in Africa, the estimated number of malaria cases in 2019 if results from Method 2 (officially used to estimate cases in these countries) were compared with those in Method 1.

### **Fig. 6.1. Funding for malaria control and elimination, 2010–2019 (% of total funding), by source of funds (constant 2019 US\$)**

Total funding for malaria control and elimination over the period 2000–2019 was estimated using data obtained from several sources, where available. The methodology below describes the collection and analysis for all available domestic and international funding for Figs. 6.1–6.5. For Figs. 6.1–6.5, data are represented for the years 2010–2019, because the Organisation for Economic Co-operation and Development (OECD) use of the multilateral system and the country-specific unit cost estimates were not available before 2010. Figs. 6.3–6.5 reflect data available for 2000–2019, where, when there are no data available for a specific funder, no imputation

was conducted and thus the trends presented in the main text should be interpreted carefully.

Contributions from governments of endemic countries were estimated as the sum of government contributions reported by NMPs for the world malaria report of the relevant year plus the estimated costs of patient care delivery services at public health facilities. If NMP contributions were missing for 2019, data reported from previous years were used after conversion to constant 2019 US\$. The number of reported malaria cases attending public health facilities was sourced from NMP reports, adjusted for diagnosis and reporting completeness. Between 1% and 3% of uncomplicated reported malaria cases were assumed to have moved to the severe stage of disease, and 50–80% of these severe cases were assumed to have been hospitalized. Costs of outpatient visits and inpatient bed-stays were estimated from the perspective of the public health care provider, using unit cost estimates from WHO-CHOosing Interventions that are Cost-Effective (WHO-CHOICE) (27). For each country, the 2010 unit cost estimates from WHO-CHOICE, expressed in the national currency, were estimated for the period 2011–2019 using the gross domestic product (GDP) annual price deflator published by the World Bank (28) on 7 July 2020, and converted in base year 2010. Country-specific unit cost estimates were then converted from national currency to constant 2019 US\$ for each year during 2010–2019. For each country, the number of adjusted reported malaria cases attending public health facilities was then multiplied by the estimated unit costs. In the absence of information on the level of care at which malaria patients attend public health facilities, uncertainty around unit cost estimates was handled through probabilistic uncertainty analysis. The mean total cost of patient care service delivery was calculated from 1000 estimations. Contributions from governments of endemic countries as reported by NMPs were available for 2000–2019.

International bilateral funding data were obtained from several sources. Data on planned funding from the government of the United States of America (USA) were sourced from the US government Foreign Assistance website (29), with the technical assistance of the Kaiser Family Foundation. Country-level funding data were available for the US Agency for International Development (USAID) for the period 2006–2019. Country-specific planned funding data from other agencies, such as the US Centers for Disease Control and Prevention (CDC) and the US Department of Defense, were not available; therefore, data on total annual planned funding from each of these two agencies were used for the period 2001–2019, as well as total annual planned funding from USAID for 2001–2005 until the introduction of country-specific funding from 2006 through 2019. For the government of the United Kingdom of Great Britain and Northern Ireland

<sup>1</sup> <https://idmod.org/documentation>

## Annex 1 – Data sources and methods

(United Kingdom), funding data towards malaria control for 2017, 2018 and 2019 were sourced from the *Statistics on International Development: final UK aid spend 2019 (30)* (*UK aid spend*) with the technical assistance of the United Kingdom Department for International Development. The *UK aid spend* data do not capture all spending from the United Kingdom that may affect malaria outcomes. The United Kingdom supports malaria control and elimination through a broad range of interventions; for example, via support to overall health systems in malaria endemic countries, and through research and development (R&D), which are not included in these data. For the period 2010–2016, United Kingdom spending data were sourced from the OECD creditor reporting system (CRS) database on aid activity (31). For all other donors, disbursement data were also obtained from the OECD CRS database on aid activity for the period 2002–2018. For each year and each funder, the country- and regional-level project-type interventions and other technical assistance were extracted. All data were converted to constant 2019 US\$. For years with no data available for a particular funder, no imputation was conducted so trends presented in the main text figures should be interpreted carefully.

Malaria-related annual funding from donors through multilateral agencies was estimated from data on (i) donors' contributions published by the Global Fund to Fight AIDS, Tuberculosis and Malaria (Global Fund) (32) from 2010 to 2019, and annual disbursements by the Global Fund to malaria endemic countries between 2003 and 2019, as reported by the Global Fund; and (ii) donors' disbursements to malaria endemic countries published in the OECD CRS and in the OECD Development Assistance Committee (DAC) members' total use of the multilateral system from 2011 through 2018 (31). All funding flows were converted to constant 2019 US\$.

For (i), the amount of funding contributed by each donor was estimated as the proportion of funding paid by each donor out of the total amount received by the Global Fund in a given year, multiplied by the total amount disbursed by the Global Fund in that same year.

For (ii), contributions from donors to multilateral channels were estimated by calculating the proportion of the core contributions received by a multilateral agency each year by each donor, then multiplying that amount by the multilateral agency's estimated investment in malaria control in that same year.

Contributions from malaria endemic countries to multilateral agencies were allocated to governments of endemic countries under the "funding source" category. Contributions from non-DAC countries and other sources to multilateral agencies were not available and were therefore not included.

Annual estimated investments were summed to estimate the total amount each funder contributed to malaria control and elimination over the period 2010–2019, and the relative percentage of the total spending contributed by each funder was calculated for the period 2010–2019.

**Fig. 6.1** excludes household spending on malaria prevention and treatment in malaria endemic countries.

### **Fig. 6.2. Funding for malaria control and elimination, 2010–2019, by source of funds (constant 2019 US\$)**

See methods notes for **Fig. 6.1** for sources of information on total funding for malaria control and elimination from governments of malaria endemic countries and on international funding flows. **Fig. 6.2** excludes household spending on malaria prevention and treatment in malaria endemic countries.

### **Fig. 6.3. Funding for malaria control and elimination, 2000–2019, by World Bank 2019 income group and source of funding (constant 2019 US\$)**

See methods notes for **Fig. 6.1** for sources of information on total funding for malaria control and elimination from governments of malaria endemic countries and on international funding flows. Data on income group classification for 2019 were sourced from the World Bank (33). For years with no data available for a particular funder, no imputation was conducted so trends presented in the main text figures should be interpreted carefully. **Fig. 6.3** excludes household spending on malaria prevention and treatment in malaria endemic countries.

### **Fig. 6.4. Funding for malaria control and elimination, 2000–2019, by channel (constant 2019 US\$)**

See methods notes for **Fig. 6.1** for sources of information on total funding for malaria control and elimination from governments of malaria endemic countries and on international funding flows. For years with no data available for a particular funder, no imputation was conducted so trends presented in the main text figures should be interpreted carefully. **Fig. 6.4** excludes household spending on malaria prevention and treatment in malaria endemic countries.

### **Fig. 6.5. Funding for malaria control and elimination, 2000–2019, by WHO region (constant 2019 US\$)**

See methods notes for **Fig. 6.1** for sources of information on total funding for malaria control and elimination from governments of malaria endemic countries and on

international funding flows. The “Unspecified” category includes all funding data for which there was no geographical information on the recipient. For years with no data available for a particular funder, no imputation was conducted so trends presented in the main text figures should be interpreted carefully. **Fig. 6.5** excludes household spending on malaria prevention and treatment in malaria endemic countries.

**Fig. 6.6. Funding for malaria-related R&D, 2007–2018, by product type (constant 2019 US\$)**

Data on funding for malaria-related R&D for 2007–2018 were sourced directly from Policy Cures Research through the G-FINDER data portal (34).

**Fig. 6.7. Malaria R&D funding from 2007 to 2018, by sector (constant 2019 US\$)**

See methods notes for **Fig. 6.6**.

**Fig. 7.1. Number of ITNs delivered by manufacturers and distributed by NMPs, 2010–2019**

Data on the number of ITNs delivered by manufacturers to countries were provided to WHO by Milliner Global Associates. Data from NMP reports were used for the number of ITNs distributed within countries.

**Fig. 7.2. Indicators of population-level coverage of ITNs, sub-Saharan Africa, 2000–2019: a) percentage of households with at least one ITN, b) percentage of households with one ITN for every two people, c) percentage of population with access to an ITN, d) percentage of population using an ITN, e) percentage of children aged under 5 years using an ITN and f) percentage of pregnant women sleeping under an ITN**

Estimates of ITN coverage were derived from a model developed by MAP (8), using a two-stage process. First, a mechanism was designed for estimating net crop (i.e. the total number of ITNs in households in a country at a given time), taking into account inputs to the system (e.g. deliveries of ITNs to a country) and outputs (e.g. loss of ITNs from households). Second, empirical modelling was used to translate estimated net crops (i.e. total number of ITNs in a country) into resulting levels of coverage (e.g. access within households, use in all ages and use among children aged under 5 years).

The model incorporates data from three sources:

- the number of ITNs delivered by manufacturers to countries, as provided to WHO by Milliner Global Associates;

- the number of ITNs distributed within countries, as reported to WHO by NMPs; and
- data from nationally representative household surveys from 39 countries in sub-Saharan Africa, from 2001 to 2018.

**Countries for analysis**

The main analysis covered 40 of the 47 malaria endemic countries or areas of sub-Saharan Africa. The islands of Mayotte (for which no ITN delivery or distribution data were available) and Cabo Verde (which does not distribute ITNs) were excluded, as were the low-transmission countries of Eswatini, Namibia, Sao Tome and Principe, and South Africa, for which ITNs comprise a small proportion of vector control. Analyses were limited to populations categorized by NMPs as being at risk.

**Estimating national net crops through time**

As described by Flaxman et al. (35), national ITN systems were represented using a discrete-time stock-and-flow model. Nets delivered to a country by manufacturers were modelled as first entering a “country stock” compartment (i.e. stored in-country but not yet distributed to households). Nets were then available from this stock for distribution to households by the NMP or through other distribution channels. To accommodate uncertainty in net distribution, the number of nets distributed in a given year was specified as a range, with all available country stock (i.e. the maximum number of nets that could be delivered) as the upper end of the range and the NMP-reported value (i.e. the assumed minimum distribution) as the lower end. The total household net crop comprised new nets reaching households plus older nets remaining from earlier times, with the duration of net retention by households governed by a loss function. However, rather than the loss function being fitted to a small external dataset – as per Flaxman et al. (35) – the loss function was fitted directly to the distribution and net crop data within the stock-and-flow model itself. Loss functions were fitted on a country-by-country basis, were allowed to vary through time, and were defined separately for conventional ITNs (cITNs) and LLINs. The fitted loss functions were compared with existing assumptions about rates of net loss from households. The stock-and-flow model was fitted using Bayesian inference and Markov chain Monte Carlo methods, which provided time-series estimates of national household net crop for cITNs and LLINs in each country, and an evaluation of under-distribution, all with posterior credible intervals.

**Estimating indicators of national ITN access and use from the net crop**

Rates of ITN access within households depend not only on the total number of ITNs in a country (i.e. the net crop), but also on how those nets are distributed among households. One factor that is known to strongly influence the

## Annex 1 – Data sources and methods

relationship between net crop and net distribution patterns among households is the size of households, which varies among countries, particularly across sub-Saharan Africa. Many recent national surveys report the number of ITNs observed in each household surveyed. Hence, it is possible not only to estimate net crop, but also to generate a histogram that summarizes the household net ownership pattern (i.e. the proportion of households with 0, 1 or 2 nets, etc). In this way, the size of the net crop was linked to distribution patterns among households while accounting for household size, making it possible to generate ownership distributions for each stratum of household size. The bivariate histogram of net crop to distribution of nets among households by household size made it possible to calculate the proportion of households with at least one ITN. Also, because the numbers of both ITNs and people in each household were available, it was possible to directly calculate two additional indicators: the proportion of households with at least one ITN for every two people, and the proportion of the population with access to an ITN within their household. For the final ITN indicator – the proportion of the population who slept under an ITN the previous night – the relationship between ITN use and access was defined using 62 surveys in which both these indicators were available ( $ITN_{use\ all\ ages} = 0.8133 \times ITN\ access_{all\ ages} + 0.0026$ ,  $R^2 = 0.773$ ). This relationship was applied to the MAP's country-year estimates of household access, to obtain ITN use among all ages. The same method was used to obtain the country-year estimates of ITN use in children aged under 5 years ( $ITN_{use\ children\ under\ 5} = 0.9327 \times ITN\ access_{children\ under\ 5} + 0.0282$ ,  $R^2 = 0.754$ ).

### **Fig. 7.3. Concentration index of ITN use by children aged under 5 years, sub-Saharan Africa at administrative level 1**

The distribution of ITN usage related to the distribution of wealth index was analysed from household surveys using the `conindex` command in Stata (36). The concentration index (37) has a value of 0 if there is no difference in the distribution of the usage related to the distribution of wealth, a positive value if the usage is concentrated among the high-wealth population and a negative value if the usage is concentrated among the low-wealth population.

### **Fig. 7.4. Percentage of the population at risk protected by IRS, by WHO region, 2010–2019**

The number of people protected by IRS was reported to WHO by NMPs. The total population of each country was taken from the 2017 revision of the *World population prospects* (27), and the proportion at risk of malaria was derived from NMP reports.

### **Fig. 7.5. Subnational areas where SMC was delivered in implementing countries in sub-Saharan Africa, 2019**

Data were provided by the Seasonal Malaria Chemoprevention (SMC) Working Group.

### **Table 7.1. Average number of children treated with at least one dose of SMC by year in countries implementing SMC, 2012–2019**

Data were provided by the London School of Hygiene & Tropical Medicine (LSHTM) and MMV.

### **Table 7.2. Average number of children targeted and treated, and total treatment doses targeted and delivered, in countries implementing SMC, 2019**

Data were provided by LSHTM and MMV.

### **Fig. 7.6. Percentage of pregnant women attending an ANC clinic at least once and receiving IPTp, by dose, sub-Saharan Africa, 2010–2019**

The total number of pregnant women eligible for IPTp was calculated by adding total live births calculated from UN population data and spontaneous pregnancy loss (specifically, miscarriages and stillbirths) after the first trimester (24). Spontaneous pregnancy loss has previously been calculated by Dellicour et al. (25). Country-specific estimates of IPTp coverage were calculated as the ratio of pregnant women receiving IPTp at ANC clinics to the estimated number of pregnant women eligible for IPTp in a given year. ANC attendance rates were derived in the same way, using the number of initial ANC visits reported through routine information systems. Local linear interpolation or information for national representative surveys was used to compute missing values. Annual aggregate estimates exclude countries for which a report or interpolation was not available for the specific year. Dose coverage could be calculated for 34 of the 38 countries with an IPTp policy.

### **Diagnostic testing and treatment**

The analysis is based on the latest nationally representative household surveys (DHS and MIS) conducted between 2015 and 2019, and surveys (latest from 2000–2005) considered baseline surveys from sub-Saharan African countries where data on malaria case management were available. Data are only available for children aged under 5 years because DHS and MIS focus on the most vulnerable population groups. Interviewers ask caregivers whether the child has had fever in the 2 weeks preceding the interview and, if so, where care was sought; whether the child received a finger or heel stick as part of the care; what treatment was received for the fever and when; and, in particular, whether the child received an artemisinin-based combination

therapy (ACT) or other antimalarial medicine. In addition to self-reported data, DHS and MIS also include biomarker testing for malaria, using rapid diagnostic tests (RDTs) that detect *P. falciparum* histidine-rich protein 2 (HRP2). Percentages and 95% CIs were calculated for each country

each year, taking into account the survey design. Median values and interquartile ranges (IQRs) were calculated using country percentages for the latest and baseline surveys.

The following indicators are presented in **Table 7.3**:

Indicator	Numerator	Denominator
Median prevalence of fever in the past 2 weeks	Children aged under 5 years with a history of fever in the past 2 weeks	Children aged under 5 years
Median prevalence of fever in the past 2 weeks for whom treatment was sought	Children aged under 5 years with a history of fever in the past 2 weeks for whom treatment was sought	Children aged under 5 years with fever in the past 2 weeks
Median prevalence of treatment seeking by source of treatment for fever (public health facility, private health facility or community health worker)	Children aged under 5 years with a history of fever in the past 2 weeks for whom treatment was sought in the public sector or private sector or community health worker	Children aged under 5 years with fever in the past 2 weeks for whom treatment was sought
Median prevalence of receiving finger or heel prick	Children aged under 5 years with a history of fever in the past 2 weeks for whom treatment was sought and who received a finger or heel prick	Children aged under 5 years with fever in the past 2 weeks for whom treatment was sought
Median prevalence of treatment with ACTs	Children aged under 5 years with a history of fever in the past 2 weeks for whom treatment was sought and who were treated with ACTs	Children aged under 5 years with fever in the past 2 weeks for whom treatment was sought in public, private or community health services
Median prevalence of treatment with ACTs among those who received a finger or heel prick	Received ACT treatment	Children aged under 5 years with fever in the past 2 weeks for whom treatment was sought and who received a finger or heel prick

The use of household survey data has several limitations. One issue is that, because of difficulty recalling past events, respondents may not provide reliable information, especially on episodes of fever and the identity of prescribed medicines, resulting in a misclassification of drugs. Also, because respondents can choose more than one source of care for one episode of fever, and because the diagnostic test and treatment question is asked broadly and hence is not linked to any specific source of care, it has been assumed that the diagnostic test and treatment were received in all the selected sources of care. However, only a low percentage (<5%) of febrile children were brought for care in more than one source of care. Data may also be biased by the seasonality of survey data collection, because DHS are carried out at various times during the year and MIS are usually timed to correspond with the high malaria transmission season. Another limitation, when undertaking trend analysis, is that DHS and MIS are done intermittently, or not at all in some countries, resulting in a relatively small number of countries in sub-Saharan Africa or for any particular 4-year period. Countries are also not the same across each 4-year period. In addition, depending on the sample size of the survey, the denominator for some indicators can be small – countries where the number of children in the denominator was less than 30 were excluded from the calculation.

**Fig. 7.7. Number of RDTs sold by manufacturers and distributed by NMPs for use in testing suspected malaria cases, 2010–2019**

The numbers of RDTs distributed by WHO region are the annual totals reported as having been distributed by NMPs. Numbers of RDT sales between 2010 and 2019 reflect sales by companies eligible for procurement. From 2010 to 2017, WHO received reports from up to 44 (cumulative number; figure differs from year to year) manufacturers that participated in the RDT Product Testing Programme by WHO, the Foundation for Innovative New Diagnostics (FIND), the CDC, and the Special Programme for Research and Training in Tropical Diseases. Since WHO Prequalification became a selection criterion for procurement, 2018 and 2019 sales data mainly focus on sales by the 11 eligible companies. The number of RDTs reported by manufacturers represents total sales to the public and private sectors worldwide.

**Fig. 7.8. Number of ACT treatment courses delivered by manufacturers and distributed by NMPs to patients, 2010–2019**

Data on ACT sales were provided by 10 manufacturers eligible for procurement by WHO and United Nations Children’s Fund (UNICEF). ACT sales were categorized as being to either the public sector or the private sector, also

## Annex 1 – Data sources and methods

taking into account the Global Fund co-payment mechanism and the Affordable Medicines Facility–malaria (AMFm) initiative. Data on ACTs distributed within countries through the public sector were taken from NMP reports.

### **Table 7.3. Summary of coverage of treatment seeking for fever, diagnosis and use of ACTs for children aged under 5 years, from household surveys in sub-Saharan Africa, at baseline (2005–2011) and most recent (2015–2019)**

See the information provided in the section titled Diagnostic testing and treatment (above).

### **Fig. 7.9. Concentration index of a) prevalence of fever in, and b) care seeking for children aged under 5 years at administrative level 1, sub-Saharan Africa**

The distribution of prevalence of fever in, and care seeking for children aged under 5 years related to the distribution of wealth index was analysed from DHS using the `conindex` command in Stata (36); see Fig. 7.3 for details.

### **Fig. 8.1. Comparison of global progress in malaria: a) case incidence and b) mortality rate, considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)**

The GTS target is a 90% reduction of malaria incidence and mortality rate by 2030, with milestones of 40% and 75% reductions in both indicators for the years 2020 and 2025, respectively (7). A curve based on a quadratic fit is used for the malaria incidence milestones. For projection of malaria incidence under current estimated trends, the same year-on-year trend observed from recent years (2017–2019) is forecast up to 2030.

### **Fig. 8.2. Map of malaria endemic countries showing progress towards the GTS 2020 malaria case incidence milestone of at least 40% reduction from a 2015 baseline**

See methods notes for Fig. 8.1.

### **Fig. 8.3. Map of malaria endemic countries showing progress towards the GTS 2020 malaria mortality rate milestone of at least 40% reduction from a 2015 baseline**

See methods notes for Fig. 8.1.

### **Fig. 8.4. Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO African Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)**

See methods notes for Fig. 8.1.

### **Fig. 8.5. Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO Region of the Americas considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)**

See methods notes for Fig. 8.1.

### **Fig. 8.6. Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO Eastern Mediterranean Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)**

See methods notes for Fig. 8.1.

### **Fig. 8.7. Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO South-East Asia region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)**

See methods notes for Fig. 8.1.

### **Fig. 8.8. Comparison of progress in malaria: a) case incidence and b) mortality rate in the WHO Western Pacific Region considering two scenarios: current trajectory maintained (blue) and GTS targets achieved (green)**

See methods notes for Fig. 8.1.

### **Fig. 9.1. Treatment failure rates among patients with *P. falciparum* malaria, WHO African Region, 2010–2019**

The box-and-whisker plots show the distribution of values for each drug, with the boxes extending from the 25th to the 75th percentile, and the middle line indicating the median. The whiskers denote adjacent values extending from the top of the box to the largest data element, which is  $\leq 1.5$  times the IQR (i.e. the distance from the 25th to the 75th percentile), and down from the bottom of the box to the smallest data element, which is  $\geq 1.5$  times the IQR. The dots denote observations outside the range of adjacent values.

### **Fig. 9.2. Treatment failure rates among patients with *P. vivax* malaria, WHO Region of the Americas, 2010–2019**

See methods notes for Fig. 9.1.

### **Fig. 9.3. Treatment failure rates among patients with *P. falciparum* malaria, WHO South-East Asia Region, 2010–2019**

See methods notes for Fig. 9.1.

**Fig. 9.4. Treatment failure rates among patients with *P. falciparum* malaria, WHO Eastern Mediterranean Region, 2010–2019**

See methods notes for Fig. 9.1.

**Fig. 9.5. Treatment failure rates among patients with *P. falciparum* malaria, WHO Western Pacific Region, 2010–2019**

See methods notes for Fig. 9.1.

**Fig. 9.6. Number of classes to which resistance was confirmed in at least one malaria vector in at least one monitoring site, 2010–2019**

Resistance to an insecticide class was considered to be confirmed in a country if at least one vector species exhibited resistance to one insecticide in the class in at least one collection site in the country, as measured by standard WHO tube tests or CDC bottle bioassays conducted with validated discriminating concentrations in 2010–2019. The map was developed based on data contained in the WHO global database for insecticide resistance in malaria vectors. These data were reported to WHO by NMPs, national public health institutes, universities and research centres, the African Network for Vector Resistance, MAP (8), VectorBase and the US President's Malaria Initiative (PMI), or extracted from scientific publications.

**Fig. 9.7. Reported insecticide resistance status as a proportion of sites for which monitoring was conducted, by WHO region, 2010–2019: pyrethroids, organochlorines, carbamates and organophosphates**

The status of resistance at each mosquito collection site for each insecticide class was assessed based on the lowest mosquito mortality reported across all standard WHO tube tests or CDC bottle bioassays conducted at the site during 2010–2019, with validated discriminating concentrations of the insecticides in the class. If multiple insecticides and mosquito species were tested between 2010 and 2019 at the collection site, the lowest mosquito mortality was considered. If the lowest mosquito mortality was below 90%, resistance was considered to be confirmed at the site; if the lowest mosquito mortality was 90% or more but below 98%, resistance was considered to be possible at the site; if the lowest mortality was more than 98%, vectors at the site were considered to be susceptible to the insecticide class. The figure was developed based on data in the WHO global database for insecticide resistance in malaria vectors. These data were reported to WHO by NMPs, national public health institutes, universities and research centres, the African Network for Vector Resistance, MAP, VectorBase and PMI, or extracted from scientific publications.

**Fig. 10.1. Trends in COVID-19 cases and deaths in malaria endemic countries globally and by WHO region (as of 23 November 2020)**

This graph is built on daily numbers of COVID-19 cases and deaths as reported to WHO (38).

**Fig. 10.2. Malaria seasonality and trends of COVID-19 cases in malaria endemic countries and areas, 2020 (as of 23 November 2020)**

For each country, the monthly average of seasonality at administrative level 1 and the daily number of COVID-19 cases reported to WHO are presented (38). To compare both trends over time, each series has been scaled to have similar maximum values in every country.

**Table 10.1. The global workstreams on the malaria response during the COVID-19 pandemic**

The table summarizes the various WHO-convened workstreams on the malaria response during the COVID-19 pandemic (39).

**Fig. 10.3. Potential RDT stockouts forecast in June 2020, if country orders were not delivered**

The figure shows forecast RDT needs and potential stockouts developed by PMI and the Global Fund as part of activities under the workstream on supplies and commodities (see Table 10.1).

**Fig. 10.4 Results from WHO surveys on disruptions of malaria related services during the COVID-19 pandemic: a) ANC services and b) diagnosis and treatment**

Data were obtained from surveys conducted in May–September 2020 by the WHO Department of Integrated Health Services. Structured online questionnaires were sent to each country office for completion by relevant national respondents (40).

**Fig. 10.5. Monthly trends in all-cause outpatients attendances in 23 countries in sub-Saharan Africa in 2019 and 2020**

Graphs of all-cause outpatient attendances were developed using data submitted by NMPs.

**Fig. 10.6. Monthly trends in malaria outpatients attendances in 24 countries in sub-Saharan Africa in 2019 and 2020**

Graphs of malaria outpatient attendances were developed using data submitted by NMPs.



## Annex 1 – Data sources and methods

### **Fig. 10.7. Estimated potential increase in malaria deaths in sub-Saharan Africa (excluding Botswana, Eswatini, Namibia and South Africa) corresponding to varying levels of disruptions of access to effective antimalarial treatment**

The figure shows projected estimates of the impact of disruptions on effective treatment with antimalarial services, using methods described by WHO (41).

### **Fig. 11.1. Distribution of malaria cases in 2019 by human development index in 2018**

For malaria cases see method for **Table 3.1**. The human development index estimates were obtained from the United Nations Development Programme (UNDP) (42).

### **Fig. 11.2. Distribution of malaria cases in 2019 by current health expenditure as percentage of GDP in 2017**

For malaria cases, see method for **Table 3.1**. The information on current health expenditure as percentage of GDP in 2017 was obtained from the World Bank data on health expenditure (43).

### **Fig. 11.3. Distribution of malaria cases in 2019 by category of governance effectiveness in 2019**

For malaria cases, see method for **Table 3.1**. The governance effectiveness estimates were obtained from the World Bank data on governance (43).

### **Fig. 11.4. Distribution of malaria cases in 2019 by category of UHC service coverage index in 2017**

For malaria cases see method for **Table 3.1**. The universal health coverage (UHC) service coverage index was obtained from the WHO Global Health Observatory (44); methods for its estimation are also provided online (45).



# References for Annex 1

1. Global technical strategy for malaria 2016–2030 (last update: 22 January 2016). Geneva: World Health Organization; 2015 ([https://www.who.int/malaria/areas/global\\_technical\\_strategy/en/](https://www.who.int/malaria/areas/global_technical_strategy/en/)).
2. Roll Back Malaria Partnership Secretariat. Action and investment to defeat malaria 2016–2030. For a malaria-free world. Geneva: World Health Organization; 2015 ([https://endmalaria.org/sites/default/files/RBM\\_AIM\\_Report\\_0.pdf](https://endmalaria.org/sites/default/files/RBM_AIM_Report_0.pdf)).
3. World malaria report 2008. Geneva: World Health Organization; 2008 (<https://www.who.int/malaria/publications/atoz/9789241563697/en/>).
4. Cibulskis RE, Aregawi M, Williams R, Otten M, Dye C. Worldwide incidence of malaria in 2009: estimates, time trends, and a critique of methods. *PLoS Med*. 2011;8(12):e1001142.
5. The R Core Team. R: A language and environment for statistical computing: reference index. Vienna: R Foundation for Statistical Computing.
6. Weiss DJ, Mappin B, Dalrymple U, Bhatt S, Cameron E, Hay SI et al. Re-examining environmental correlates of *Plasmodium falciparum* malaria endemicity: a data-intensive variable selection approach. *Malar J*. 2015;14(1):68.
7. Cameron E, Battle KE, Bhatt S, Weiss DJ, Bisanzio D, Mappin B et al. Defining the relationship between infection prevalence and clinical incidence of *Plasmodium falciparum* malaria. *Nat Commun*. 2015;6:8170.
8. Malaria Atlas Project [website]. (<https://malariaatlas.org>).
9. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature*. 2015;526(7572):207–11 (<https://pubmed.ncbi.nlm.nih.gov/26375008>).
10. World malaria report 2019. Geneva: World Health Organization; 2019 (<https://www.who.int/publications/i/item/9789241565721>).
11. Alles HK, Mendis KN, Carter R. Malaria mortality rates in South Asia and in Africa: implications for malaria control. *Parasitol Today*. 1998;14(9):369–75.
12. Luxemburger C, Ricci F, Nosten F, Raimond D, Bathet S, White NJ. The epidemiology of severe malaria in an area of low transmission in Thailand. *Trans R Soc Trop Med Hyg*. 1997;91(3):256–62.
13. Meek SR. Epidemiology of malaria in displaced Khmers on the Thai–Kampuchean border. *Southeast Asian J Trop Med Public Health*. 1988;19(2):243–52.
14. Douglas NM, Pontororing GJ, Lampah DA, Yeo TW, Kenangalem E, Poespoprodjo JR et al. Mortality attributable to *Plasmodium vivax* malaria: a clinical audit from Papua, Indonesia. *BMC Med*. 2014;12(1):217.
15. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE et al. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet*. 2015;385(9966):430–40.
16. Ross A, Maire N, Molineaux L, Smith T. An epidemiologic model of severe morbidity and mortality caused by *Plasmodium falciparum*. *Am J Trop Med Hyg*. 2006;75(2 Suppl):63–73.
17. Griffin JT, Ferguson NM, Ghani AC. Estimates of the changing age-burden of *Plasmodium falciparum* malaria disease in sub-Saharan Africa. *Nat Commun*. 2014;5(1):1–10.
18. Walker PG, Griffin JT, Cairns M, Rogerson SJ, Van Eijk AM, Ter Kuile F et al. A model of parity-dependent immunity to placental malaria. *Nat Commun*. 2013;4(1):1–11 (<https://www.nature.com/articles/ncomms2605>).
19. Walker PG, ter Kuile FO, Garske T, Menendez C, Ghani AC. Estimated risk of placental infection and low birthweight attributable to *Plasmodium falciparum* malaria in Africa in 2010: a modelling study. *Lancet Glob Health*. 2014;2(8):e460–e7 (<https://pubmed.ncbi.nlm.nih.gov/25103519>).
20. The DHS Program: demographic and health surveys [website]. Washington, DC: United States Agency for International Development (<https://dhsprogram.com/>).
21. World population prospects [website]. New York City: United Nations; 2020 (<https://population.un.org/wpp/>).
22. WorldPop [website]. (<https://www.worldpop.org/>).
23. Cairns M, Roca-Feltrer A, Garske T, Wilson AL, Diallo D, Milligan PJ et al. Estimating the potential public health impact of seasonal malaria chemoprevention in African children. *Nat Commun*. 2012;3:881.

24. Adekanbi AO, Olayemi OO, Fawole AO, Afolabi KA. Scourge of intra-partum foetal death in Sub-Saharan Africa. *World J Clin Cases*. 2015;3(7):635 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4517338/>).
25. Dellicour S, Tatem AJ, Guerra CA, Snow RW, ter Kuile FO. Quantifying the number of pregnancies at risk of malaria in 2007: a demographic study. *PLoS Med*. 2010;7(1):e1000221.
26. High burden to high impact: a targeted malaria response. Geneva: World Health Organization; 2019 (<https://www.who.int/malaria/publications/atoz/high-impact-response/en/>).
27. Cost effectiveness and strategic planning (WHO-CHOICE) [website]. Geneva: World Health Organization; (<https://www.who.int/choice/en/>).
28. Indicators [website]. Washington (DC): The World Bank; 2020 (<https://data.worldbank.org/indicator>).
29. Foreign assistance [website]. US government (<https://foreignassistance.gov/>).
30. Statistics on international development: final UK aid spend 2019. United Kingdom: Foreign, Commonwealth & Development Office; 2019 (<https://www.gov.uk/government/statistics/statistics-on-international-development-final-uk-aid-spend-2019>).
31. Creditor reporting system (CRS) [website]. Organisation for Economic Co-operation and Development (<https://stats.oecd.org/Index.aspx?DataSetCode=CRS1>).
32. Financials [website]. Geneva: Global Fund to Fight AIDS, Tuberculosis and Malaria (<https://www.theglobalfund.org/en/financials/>).
33. World Bank country and lending groups [website]. Washington DC: World Bank (<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>).
34. Policy Cures Research: G-FINDER data portal [website]. Sydney, Australia: Policy Cures Research (<https://gfinderdata.policycuresresearch.org>).
35. Flaxman AD, Fullman N, Otten MW, Menon M, Cibulskis RE, Ng M et al. Rapid scaling up of insecticide-treated bed net coverage in Africa and its relationship with development assistance for health: a systematic synthesis of supply, distribution, and household survey data. *PLoS Med*. 2010;7(8):e1000328.
36. O'Donnell O, O'Neill S, Van Ourti T, Walsh B. Conindex: estimation of concentration indices. *Stata J*. 2016;16(1):112–38.
37. O'Donnell O, Van Doorslaer E, Wagstaff A, Lindelow M. Analyzing health equity using household survey data: a guide to techniques and their implementation. The World Bank; 2007 (<https://elibrary.worldbank.org/doi/abs/10.1596/978-0-8213-6933-3>).
38. WHO Coronavirus Disease (COVID-19) Dashboard [website]. Geneva: World Health Organization (<https://covid19.who.int/>).
39. Jointly addressing endemic malaria and pandemic COVID-19. Geneva: World Health Organization; 2019 ([https://www.who.int/malaria/areas/epidemics\\_emergencies/covid-19/who-gmp-response-covid-19.pdf](https://www.who.int/malaria/areas/epidemics_emergencies/covid-19/who-gmp-response-covid-19.pdf)).
40. Pulse survey on continuity of essential health services during the COVID-19 pandemic: interim report, 27 August 2020. Geneva: World Health Organization; 2020 ([https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS\\_continuity-survey-2020.1](https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS_continuity-survey-2020.1)).
41. The potential impact of health service disruptions on the burden of malaria. A modelling analysis for countries in sub-Saharan Africa. Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/the-potential-impact-of-health-service-disruptions-on-the-burden-of-malaria>).
42. Human development index (HDI) [website]. Geneva: United Nations Development Programme (UNDP); 2020 (<http://www.hdr.undp.org/en/content/human-development-index-hdi>).
43. Kaufmann D, Kraay A, Mastruzzi M. The worldwide governance indicators: methodology and analytical issues: World Bank Policy Research Working Paper No. 5430. Washington, DC: The World Bank; 2010 ([https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=1682130](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=1682130)).
44. WHO Global Health Observatory data repository: Index of service coverage (<https://apps.who.int/gho/data/view.main.INDEXOFESSENTIALSERVICECOVERAGE>).
45. Hogan D, Hosseinpoor, AR, Boerma T. Technical note: developing an index for the coverage of essential health services. Geneva: World Health Organization; 2005 ([https://www.who.int/healthinfo/universal\\_health\\_coverage/UHC\\_WHS2016\\_TechnicalNote\\_May2016.pdf](https://www.who.int/healthinfo/universal_health_coverage/UHC_WHS2016_TechnicalNote_May2016.pdf)).

# Annex 2 – A. WHO African Region, a. West Africa

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 394 million

**Parasites:** *P. falciparum* (almost 100%) and other (<1%)

**Vectors:** *An. arabiensis*, *An. coluzzii*, *An. funestus s.l.*, *An. gambiae s.l.*, *An. hispaniola*, *An. labranchiae*, *An. melas*, *An. moucheti*, *An. multicolor*, *An. nili s.l.*, *An. pharoensis* and *An. sergentii s.l.*

## FUNDING (US\$), 2010–2019

557.1 million (2010), 568.6 million (2015), 792.0 million (2019); increase 2010–2019: 42%

**Proportion of domestic source<sup>a</sup> in 2019:** 10%

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2019

**Countries with ≥80% coverage with either LLIN or IRS in 2019:** Cabo Verde and Ghana

**Countries with ≥50% coverage with either LLIN or IRS in 2019:** Burkina Faso, Côte d'Ivoire, Guinea, Liberia, Mali, Niger, Senegal, Sierra Leone and Togo

**Countries that implemented IPTp in 2019:** Benin, Burkina Faso, Côte d'Ivoire, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali, Mauritania, Niger, Nigeria, Senegal, Sierra Leone and Togo

**Countries with >30% IPTp<sup>3+</sup> in 2019:** Burkina Faso, Côte d'Ivoire, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali, Niger, Senegal, Sierra Leone and Togo

**Percentage of suspected cases tested (reported):** 44% (2010), 71% (2015), 98% (2019)

**Number of ACT courses distributed:** 32.2 million (2010), 47.4 million (2015), 65.1 million (2019)

**Number of any antimalarial treatment courses (incl. ACT) distributed:** 32.2 million (2010), 49.3 million (2015), 66.9 million (2019)

## REPORTED CASES AND DEATHS IN PUBLIC SECTOR, 2010–2019

**Total (presumed and confirmed) cases:** 29.4 million (2010), 52.3 million (2015), 65.6 million (2019)

**Confirmed cases:** 7.1 million (2010), 33.3 million (2015), 57.0 million (2019)

**Percentage of total cases confirmed:** 24.3% (2010), 63.6% (2015), 87.0% (2019)

**Deaths:** 39 000 (2010), 23 000 (2015), 18 700 (2019)

**Children aged under 5 years, presumed and confirmed cases:** 11.9 million (2010), 21.0 million (2015), 27.7 million (2019)

**Children aged under 5 years, percentage of total cases:** 40.6% (2010), 40.2% (2015), 42.3% (2019)

**Children aged under 5 years, deaths:** 214 100 (2010), 22 100 (2015), 38 700 (2019)

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:** 116.1 million (2010), 105.5 million (2015), 112.1 million (2019); decrease 2010–2019: 3%

**Deaths:** 306 000 (2010), 224 500 (2015), 196 100 (2019); decrease 2010–2019: 36%

## ACCELERATION TO ELIMINATION

**Countries with subnational/territorial elimination programme:** Senegal

**Countries with nationwide elimination programme:** Cabo Verde

**Zero indigenous cases for 3 consecutive years (2017, 2018 and 2019):** Algeria

**Zero indigenous cases in 2019:** Cabo Verde


**Certified as malaria free since 2010:** Algeria (since May 2019)

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH *P. FALCIPARUM* MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2010–2019	113	0.0	0.0	11.9	0.0	2.2
AS–AQ	2010–2019	91	0.0	0.0	8.0	0.0	1.8
AS–PY	2011–2016	7	0.0	0.5	1.2	0.0	0.6
DHA–PPQ	2010–2018	27	0.0	0.0	2.4	0.0	0.0

AL: artemether-lumefantrine; AS–AQ: artesunate-amodiaquine; AS–PY: artesunate-pyronaridine; DHA–PPQ: dihydroartemisinin-piperazine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. *P. falciparum* parasite prevalence (PfPP), 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland; USAID: United States Agency for International Development.


<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

## C. Malaria funding<sup>a</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.

## D. Share of estimated malaria cases, 2019



## E. Percentage of population with access to either LLINs or IRS, 2019


Source: ITN coverage model from MAP




IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; LLIN: long-lasting insecticidal net; MAP: Malaria Atlas Project.

<sup>a</sup> Cabo Verde is an E-2020 country; vector control targeted at foci.


## F. Estimated number of cases in countries on track to reduce case incidence by ≥40% by 2020




## G. Estimated number of cases in countries likely to reduce case incidence by <40% by 2020




## H. Estimated number of cases in countries with an increase in case incidence, 2015–2019




## I. Change in estimated malaria incidence and mortality rates, 2015–2019



## J. Incidence in 2019 compared to baseline (2016–2018)



## K. Reported indigenous cases in countries with national elimination activities, 2015 versus 2019



## KEY MESSAGES

- About 394 million people living in the 17 countries of West Africa are at high risk of malaria. Algeria was certified malaria free in May 2019, following 3 consecutive years with zero indigenous cases. Cabo Verde has had zero indigenous cases since February 2018 and since then has started its preparation for the certification process. The high burden to high impact (HBHI) initiative was initiated in Burkina Faso, Ghana, Niger and Nigeria in 2019, leading to evidence-based national strategic plans and funding requests. In countries of this subregion, except for Algeria and Cabo Verde, malaria transmission is year-round and almost exclusively due to *P. falciparum*, with strong seasonality in the Sahelian countries.
- The subregion had about 112 million estimated cases and about 196 000 estimated deaths – a 3% and 36% decrease compared with 2010, respectively. Five countries accounted for over 80% of the estimated cases: Nigeria (54%), Côte d'Ivoire (7%), Niger (7%), Burkina Faso (7%) and Mali (6%). More than 65 million cases were reported in the public and private sectors, and in the community, of which 42.3% were in children aged under 5 years, and 57 million (87%) were confirmed. The proportion of total cases that were confirmed has improved substantially over time, being only 24.3% in 2010. A total of 38 697 malaria deaths were reported in children aged under 5 years; this figure exceeded the total malaria deaths, indicating challenges in the surveillance of malaria mortality in some countries.
- In nine of the 17 countries in this subregion, where routine distribution of LLINs or use of IRS is still applicable, 50% or more of the population had access to the interventions. Five countries are on track to meet the GTS target by reducing case incidence by at least 40% by 2020 compared with 2015 (Algeria, which is already certified malaria free, Cabo Verde, the Gambia, Senegal and Togo). In nine countries, although there is progress towards meeting the target, efforts need to be

- accelerated to achieve the 40% reduction (Benin, Burkina Faso, Ghana, Guinea, Mali, Niger, Nigeria, Mauritania and Sierra Leone). In Côte d'Ivoire, Guinea-Bissau and Liberia, incidence increased in 2019 compared with 2015. After a large increase in indigenous cases in Cabo Verde between 2016 and 2017, the country has been reporting zero indigenous cases since February 2018. In addition to Algeria and Cabo Verde, Burkina Faso and Mali are on track to reduce malaria mortality rates by at least 40%. However, the estimation from Burkina Faso is affected by the decline in reporting completeness, from 98% in 2018 to 60% in 2019.
- The Nouakchott Declaration was adopted in 2013 and the new Sahel Malaria Elimination Initiative (SaME) was launched in 2018 by ministers of the eight Sahelian countries (Burkina Faso, Cabo Verde, Chad, the Gambia, Mali, Mauritania, Niger and Senegal) to accelerate implementation of high-impact strategies towards eliminating malaria by 2030. In line with these initiatives, an action plan was adopted in 2019. In addition to Cabo Verde as an eliminating country, the Gambia, Mauritania, Niger and Senegal have reoriented their programmes towards malaria subnational elimination.
- Vector resistance to pyrethroids was confirmed in 91% of the sites, to organochlorines in 95%, to carbamates in 42% and to organophosphates in 24%. Eight countries have developed their insecticide resistance monitoring and management plans.
- Challenges include inadequate political commitment and leadership, weak malaria programme management, insufficient prioritization and sustainability of interventions, inappropriate application of larviciding, inadequate domestic financing and weak surveillance systems, including a lack of well-functioning vital registration systems.

# Annex 2 – A. WHO African Region, b. Central Africa

## EPIDEMIOLOGY

Population denominator used to compute incidence and mortality rate: 186 million

Parasites: *P. falciparum* (100%)

Vectors: *An. arabiensis*, *An. funestus s.l.*, *An. gambiae s.l.*, *An. melas*, *An. moucheti*, *An. nili s.l.* and *An. pharoensis*.

## FUNDING (US\$), 2010–2019

250.5 million (2010), 376.4 million (2015), 422.5 million (2019); increase 2010–2019: 69%

Proportion of domestic source<sup>a</sup> in 2019: 17%

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2019

Countries with ≥80% coverage with either LLIN or IRS in 2019: Sao Tome and Principe

Countries with ≥50% coverage with either LLIN or IRS in 2019: Burundi, Cameroon, Central African Republic, Congo and Democratic Republic of the Congo

Countries that implemented IPTp in 2019: Angola, Burundi, Cameroon, Central African Republic, Chad, Congo, Democratic Republic of the Congo, Equatorial Guinea, Gabon and Sao Tome and Principe

Countries with >30% IPTp<sup>3+</sup> in 2019: Burundi, Cameroon, Chad, Democratic Republic of the Congo and Gabon

Percentage of suspected cases tested (reported): 41% (2010), 92% (2015), 95% (2019)

Number of ACT courses distributed: 18.2 million (2010), 22.4 million (2015), 34.0 million (2019)

Number of any antimalarial treatment courses (incl. ACT) distributed: 19.0 million (2010), 22.4 million (2015), 34.2 million (2019)

## REPORTED CASES AND DEATHS IN PUBLIC SECTOR, 2010–2019

Total (presumed and confirmed) cases: 20.4 million (2010), 24.6 million (2015), 48.8 million (2019)

Confirmed cases: 6.6 million (2010), 22.2 million (2015), 47.0 million (2019)

Percentage of total cases confirmed: 32.6% (2010), 90.1% (2015), 96.3% (2019)

Deaths: 40 400 (2010), 58 200 (2015), 45 400 (2019)

Children aged under 5 years, presumed and confirmed cases: 9.1 million (2010), 11.3 million (2015), 22.8 million (2019)

Children aged under 5 years, percentage of total cases: 44.9% (2010), 46.1% (2015), 46.8% (2019)

Children aged under 5 years, deaths: 26 000 (2010), 37 100 (2015), 22 500 (2019)

Children aged under 5 years, percentage of total deaths: 64% (2010), 64% (2015), 50% (2019)

## ESTIMATED CASES AND DEATHS, 2010–2019

Cases: 43.4 million (2010), 42.1 million (2015), 52.3 million (2019); increase 2010–2019: 21%

Deaths: 118 200 (2010), 92 100 (2015), 89 300 (2019); decrease 2010–2019: 24%

## ACCELERATION TO ELIMINATION


Countries with subnational/territorial elimination programme: Sao Tome and Principe

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH *P. FALCIPARUM* MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2010–2019	40	0.0	1.7	13.6	0.0	3.5
AS-AQ	2010–2019	44	0.0	1.7	8.2	0.0	4.4
DHA-PPQ	2010–2017	12	0.0	0.0	5.2	0.0	2.6

AL: artemether-lumefantrine; AS-AQ: artesunate-amodiaquine; DHA-PPQ: dihydroartemisinin-piperazine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. *P. falciparum* parasite prevalence (PfPP), 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland; USAID: United States Agency for International Development.


<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

## C. Malaria funding<sup>a</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.

#### D. Share of estimated malaria cases, 2019




#### E. Percentage of population with access to either LLINs or IRS, 2019

Source: ITN coverage model from MAP




IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; LLIN: long-lasting insecticidal net; MAP: Malaria Atlas Project.


#### F. Estimated number of cases in countries likely to reduce case incidence by <40% by 2020



#### G. Estimated number of cases in countries with an increase in case incidence, 2015–2019




#### H. Change in estimated malaria incidence and mortality rates, 2015–2019




<sup>a</sup> Sao Tome and Principe already achieved the 40% reduction in mortality rate in 2015; since then, there has been no change.

#### I. Incidence in 2019 compared to baseline (2016–2018)



#### J. Reported indigenous cases in countries with national elimination activities, 2015 versus 2019



## KEY MESSAGES

- About 186 million people living in the 10 countries of Central Africa are at high risk of malaria. Malaria transmission, almost exclusively due to *P. falciparum*, occurs throughout the year except in the north of Cameroon, northern Chad and the southern part of the Democratic Republic of the Congo. The HBHI initiative has been initiated in Cameroon and the Democratic Republic of the Congo.
- In 2019, the subregion had over 51 million estimated cases and almost 90 000 estimated deaths – a 12% increase and a 24% decrease compared with 2010, respectively. Five countries in the region accounted for 80% of the estimated cases: the Democratic Republic of the Congo accounted for 55.5% of estimated cases, followed by Angola (14.9%), Cameroon (12.8%), Burundi (5.8%) and Chad (5.2%). A similar distribution was seen for estimated malaria deaths, which were mainly observed in the Democratic Republic of the Congo (49%), Angola (15%), Cameroon (13%) and Chad (10%). More than 48 million cases were reported in the public and private sector, and in the community; of these, 46.8% were in children aged under 5 years and 46.9 million (96.3%) were confirmed. The proportion of total cases that were confirmed has improved substantially over time, being only 32.6% in 2010.
- Progress has been made towards achieving the GTS target of a 40% reduction in incidence by 2020 in Cameroon, Central African Republic, Chad, Equatorial Guinea and Gabon, but greater efforts are needed to ensure these countries meet the target. Five countries saw an increase in estimated

- malaria incidence between 2015 and 2019; Burundi had the largest increase (54%), followed by Angola (18%), Sao Tome and Principe (10%), the Democratic Republic of the Congo (5%) and the Congo (4%). Sao Tome and Principe also saw a slight increase in reported cases, although there have been zero deaths reported since 2018. Coverage of preventive vector control measures remains low in the region, except for Sao Tome and Principe with more than 80% coverage. In 2019, Angola, Burundi, Cameroon, the Congo and the Democratic Republic of the Congo conducted LLIN mass campaigns. Additionally, Cameroon and Chad are implementing SMC in targeted areas of the country.
- Vector resistance to pyrethroids was confirmed in 86% of the sites, to organochlorines in 94%, to carbamates in 20% and to organophosphates in 5%. Vector resistance to organochlorines was confirmed in all countries, and to pyrethroids in all countries except Sao Tome and Principe. Four countries have developed their insecticide resistance monitoring and management plans.
- The performance of the surveillance system varies across countries in the region, as can be seen through the completeness of public sector data reported for 2019. All countries except Sao Tome and Principe reported a public sector completeness rate below 100%. Additional challenges include insufficient domestic and international funding, and frequent malaria outbreaks.



# Annex 2 – A. WHO African Region, c. Countries with high transmission in East and Southern Africa

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 360 million

**Parasites:** *P. falciparum* (almost 100%), *P. vivax* (<1%) and other (<1%)

**Vectors:** *An. arabiensis*, *An. funestus s.l.*, *An. gambiae s.l.*, *An. gambiae s.s.*, *An. leesonii*, *An. nili*, *An. pharoensis*, *An. rivulorum*, *An. stephensi s.l.*<sup>a</sup> and *An. vaneedeni*.

<sup>a</sup> A potential vector identified.

## FUNDING (US\$), 2010–2019

758.6 million (2010), 733.7 million (2015), 698.1 million (2019); decrease 2010–2019: 8%

**Proportion of domestic source<sup>a</sup> in 2019:** 9%

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2019

**Countries with ≥80% coverage with either LLIN or IRS in 2019:** none

**Countries with ≥50% coverage with either LLIN or IRS in 2019:** Kenya, Madagascar, Malawi, Mozambique, Uganda and United Republic of Tanzania

**Countries that implemented IPTp in 2019:** Kenya, Madagascar, Malawi, Mozambique, South Sudan, Uganda, United Republic of Tanzania (mainland), Zambia and Zimbabwe

**Countries with >30% IPTp<sup>3+</sup> in 2019:** Madagascar, Malawi, Mozambique, Uganda, United Republic of Tanzania and Zambia

**Percentage of suspected cases tested (reported):**<sup>a</sup> 30% (2010), 80% (2015), 91% (2019)

**Number of ACT courses distributed:**<sup>b</sup> 84.5 million (2010), 108.2 million (2015), 79.7 million (2019)

**Number of any antimalarial treatment courses (incl. ACT) distributed:** 84.7 million (2010), 109.9 million (2015), 87.6 million (2019)

<sup>a</sup> Uganda did not report any suspected cases in 2019.

<sup>b</sup> Malawi, South Sudan and Zimbabwe did not report on treatment courses distributed in 2019.

## REPORTED CASES AND DEATHS IN PUBLIC SECTOR, 2010–2019

**Total (presumed and confirmed) cases:** 53.2 million (2010), 54.3 million (2015), 61.0 million (2019)

**Confirmed cases:** 19.9 million (2010), 40.2 million (2015), 56.8 million (2019)

**Percentage of total cases confirmed:** 37.5% (2010), 74.1% (2015), 93.0% (2019)

**Deaths:** 70 700 (2010), 38 300 (2015), 17 700 (2019)

**Children aged under 5 years, presumed and confirmed cases:** 21.6 million (2010), 17.6 million (2015), 21.3 million (2019)

**Children aged under 5 years, percentage of total cases:** 40.5% (2010), 32.5% (2015), 34.9% (2019)

**Children aged under 5 years, deaths:** 25 300 (2010), 10 400 (2015), 7000 (2019)

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:**<sup>a</sup> 55.8 million (2010), 51.4 million (2015), 50.0 million (2019);

decrease 2010–2019: 10%

**Deaths:** 117 000 (2010), 100 800 (2015), 98 500 (2019); decrease 2010–2019: 16%

<sup>a</sup> Estimated cases are derived from the *PPPr*-to-incidence model, which means that estimated cases are lower than reported by the country.

## ACCELERATION TO ELIMINATION


**Countries with subnational/territorial elimination programme:** United Republic of Tanzania (Zanzibar)

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH *P. FALCIPARUM* MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2010–2019	131	0.0	1.4	19.5	0.0	3.7
AS-AQ	2011–2018	30	0.0	0.0	2.0	0.0	1.0
DHA-PPQ	2010–2019	24	0.0	0.7	6.0	0.0	1.4

AL: artemether-lumefantrine; AS-AQ: artesunate-amodiaquine; DHA-PPQ: dihydroartemisinin-piperazine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. *P. falciparum* parasite prevalence (PfPP), 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland; USAID: United States Agency for International Development.


<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

## C. Malaria funding<sup>a</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.

### D. Share of estimated malaria cases, 2019




### E. Percentage of population with access to either LLINs or IRS, 2019

Source: ITN coverage model from MAP




IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; LLIN: long-lasting insecticidal net; MAP: Malaria Atlas Project.


### F. Estimated number of cases in countries on track to reduce case incidence by ≥40% by 2020




### G. Estimated number of cases in countries likely to reduce case incidence by <40% by 2020




### H. Estimated number of cases in countries with an increase in case incidence, 2015–2019



### I. Change in estimated malaria incidence and mortality rates, 2015–2019



### J. Incidence in 2019 compared to baseline (2016–2018)



## KEY MESSAGES

- About 360 million people in the 11 countries with high transmission in East and Southern Africa are at high risk of malaria. Malaria transmission is almost exclusively due to *P. falciparum* (except in Ethiopia), and is highly seasonal in Ethiopia, Madagascar and Zimbabwe, and in coastal and highland areas of Kenya. Malaria transmission is stable in most of Malawi, Mozambique, South Sudan, Uganda, the United Republic of Tanzania and Zambia. The HBHI initiative has been initiated in Mozambique and Uganda.
- The subregion had 50 million estimated cases and about 98 500 estimated deaths, representing a 10% and 16% decrease compared with 2010, respectively. Three countries accounted for over 50% of the estimated cases: Uganda (23.2%), Mozambique (18.7%) and the United Republic of Tanzania (12.9%). In the public and private sector and the community, 61 million cases were reported, of which 34.9% were in children aged under 5 years and over 56 million (93%) were confirmed. The proportion of total cases that were confirmed improved substantially over time, from only 37.5% in 2010. A significantly lower number of deaths were reported in 2019 (17 700) compared with 2010 (70 700) and 2015 (38 300).
- In 2019, Ethiopia had already achieved the GTS target of a 40% reduction in incidence by 2020. Zambia and Zimbabwe were closely approaching the target with a reduction in incidence of 33% and 30%, respectively, between 2015 and 2019, whereas all other countries in the region reported either small reductions in incidence, or increases (countries that reported increases were

- Madagascar, Rwanda, Uganda and, to a lesser extent, the United Republic of Tanzania). In more than half of the countries, 50% or more of the population had access to LLINs or IRS in 2019.
- Reported cases in Rwanda increased from 2.5 million in 2015 to 3.6 million in 2019, an increase of 42.6%. Madagascar also reported an increase of 38.1% during the period 2015–2019. Causes of such increases can include inadequate vector control, climatic factors and improved reporting. Uganda reported a 21% increase compared with 2015, which may have resulted from the rapid public health response to the 25% increase in cases that was reported between 2016 and 2017. Zanzibar (United Republic of Tanzania) reported 6963 cases in 2019, over 4.5 times higher than the number of cases reported in 2018 (1532).
- Vector resistance to pyrethroids was confirmed in 74% of the sites, to organochlorines in 42%, to carbamates in 26% and to organophosphates in 14%. Vector resistance to pyrethroids, organochlorines and carbamates was confirmed in all countries except South Sudan, which did not report resistance monitoring. Eleven countries have developed their insecticide resistance monitoring and management plans.
- Challenges include frequent epidemics, emergencies, inadequate response (South Sudan), inadequate funding, delays in critical commodities and weak surveillance systems in several countries.

# Annex 2 – A. WHO African Region, d. Countries with low transmission in East and Southern Africa

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 14 million

**Parasites:** *P. falciparum* (96%), *P. vivax* (4%) and other (<1%)

**Vectors:** *An. arabiensis*, *An. funestus s.l.*, *An. funestus s.s.*, *An. gambiae s.l.* and *An. gambiae s.s.*

## FUNDING (US\$), 2010–2019

68.8 million (2010), 25.9 million (2015), 47.0 million (2019); decrease 2010–2019: 32%

**Proportion of domestic source<sup>a</sup> in 2019:** 73%

**Regional funding mechanisms:** Southern Africa Malaria Elimination Eight Initiative

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2019

**Countries with ≥80% coverage of at-risk population with either LLIN or IRS in 2019:** None

**Countries with ≥80% coverage of high risk population with either LLIN or IRS in 2019:** Botswana

**Countries with >30% IPTp3+ in 2019:** none

**Percentage of suspected cases tested (reported):<sup>a</sup>** 100% (2010), 100% (2015), 96% (2019)

**Number of ACT courses distributed:<sup>b</sup>** 575 000 (2010), 366 000 (2015), 224 000 (2019)

**Number of any antimalarial treatment courses (incl. ACT) distributed:** 575 000 (2010), 366 000 (2015), 224 000 (2019)

<sup>a</sup> Comoros and South Africa did not report any suspected cases in 2019.

<sup>b</sup> Comoros and Eswatini did not report on treatment courses distributed in 2019.

## REPORTED CASES AND DEATHS IN PUBLIC SECTOR, 2010–2019

**Total (presumed and confirmed) cases:** 205 300 (2010), 47 800 (2015), 132 500 (2019)

**Confirmed cases:** 82 400 (2010), 33 900 (2015), 132 500 (2019)

**Percentage of total cases confirmed:** 40.2% (2010), 70.8% (2015), 100% (2019)

**Deaths:<sup>a</sup>** 242 (2010), 178 (2015), 99 (2019)

**Children aged under 5 years, presumed and confirmed cases:** 56 400 (2010), 7300 (2015), 43 900 (2019)

**Children aged under 5 years, percentage of total cases:** 27.5% (2010), 15.2% (2015), 33.2% (2019)

**Children aged under 5 years, deaths:** 37 (2010), 16 (2015), 1 (2019)

<sup>a</sup> No report for Comoros in 2019.

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:** 133 200 (2010), 90 500 (2015), 224 900 (2019); increase 2010–2019: 69%

**Deaths:** 344 (2010), 293 (2015), 569 (2019); increase 2010–2019: 65%

## ACCELERATION TO ELIMINATION


**Countries with nationwide elimination programme:** Botswana, Comoros, Eswatini, Namibia and South Africa

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH *P. FALCIPARUM* MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2011–2017	18	0.0	0.0	2.5	0.0	0.0
AS–AQ	2010–2016	18	0.0	2.4	7.9	0.0	5.2

AL: artemether–lumefantrine; AS–AQ: artesunate–amodiaquine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. Confirmed malaria cases per 1000 population, 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland.


<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

## C. Malaria funding<sup>a</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.

#### D. Share of estimated malaria cases, 2019



#### E. Percentage of population with access to either LLINs or IRS, 2019

Source: ITN coverage model from MAP




IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; LLIN: long-lasting insecticidal net; MAP: Malaria Atlas Project.


<sup>a</sup> Comoros and Eritrea have ITN coverage estimated by a model from MAP.

<sup>b</sup> LLIN and IRS coverage is combined in Namibia and South Africa because there is no overlap in the areas where they are used.


#### F. Estimated number of cases in countries on track to reduce case incidence by ≥40% by 2020



#### G. Estimated number of cases in countries with an increase in case incidence, 2015–2019




#### H. Change in estimated malaria incidence and mortality rates, 2015–2019




<sup>a</sup> Eswatini already achieved the 40% reduction in mortality rate in 2015; since then there has been no change.

#### I. Percentage of total confirmed cases investigated, 2019



#### J. Reported indigenous cases in countries with national elimination activities, 2015 versus 2019



## KEY MESSAGES

- About 14 million people in the six countries with low transmission in East and Southern Africa are at high risk of malaria. Around 132 000 cases were reported, of which 33.2% were in children aged under 5 years and 100% were confirmed. The proportion of total cases that were confirmed improved substantially over time, from only 40.2% in 2010.
- Progress has been made towards achieving the GTS target of a 40% reduction in incidence by 2020 in Botswana, Namibia and South Africa. Estimated cases in Namibia increased significantly, from 2590 in 2010 to 61 564 in 2018, then declined greatly, falling to 5618 in 2019. Estimated indigenous cases in Botswana declined from 2229 in 2010 to 257 in 2019.
- Between 2015 and 2019, Botswana, Comoros, Eswatini and South Africa recorded an increase in reported indigenous, imported and unclassified cases: Botswana (8%, from 326 to 352), Comoros (1086%, from 1300 to 15 421), Eswatini (270%, from 195 to 722) and South Africa (2392%, from 555 to 13 833). Nevertheless, the number of indigenous cases in Botswana declined over the same period, from 326 to 169 cases. Between 2018 and 2019, increases in cases were reported in Eswatini (10%) and South Africa (45%), whereas decreases were reported in Botswana (40%) and Comoros (1%). Conversely, Namibia reported a 72% reduction in cases between 2015 (12 050) and 2019 (3404).

- Vector resistance to pyrethroids was confirmed in 29% of the sites, to organochlorines in 18%, to carbamates in 0% and to organophosphates in 7%. There remain significant gaps in standard resistance monitoring for carbamates and organophosphates. Five countries have developed their insecticide resistance monitoring and management plans.
- Challenges include inadequate coverage of vector control, bottlenecks in procurement and supply management, importation of cases from neighbouring countries and resurgence during the past 3 years.

# Annex 2 – B. WHO Region of the Americas

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 139 million

**Parasites:** *P. vivax* (76%), *P. falciparum* and mixed (24%), and other (<1%)

**Vectors:** *An. albimanus*, *An. albirtarsis*, *An. aquasalis*, *An. argyritarsis*, *An. braziliensis*, *An. cruzii*, *An. darlingi*, *An. neivai*, *An. nuneztovari*, *An. pseudopunctipennis* and *An. punctimacula*.

## FUNDING (US\$), 2010–2019

220.5 million (2010), 197.4 million (2015), 139.2 million (2019); decrease 2010–2019: 37%

**Proportion of domestic source<sup>a</sup> in 2019:** 86%

**Regional funding mechanisms:** Regional Malaria Elimination Initiative

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2010–2019

**Number of people protected by IRS:** 2.78 million (2010), 2.81 million (2015), 1.35 million (2019)

**Total LLINs distributed:<sup>a</sup>** 363 000 (2010), 875 000 (2015), 1 122 000 (2019)

**Number of RDTs distributed:** 83 700 (2010), 533 900 (2015), 1 232 700 (2019)

**Number of ACT courses distributed:** 148 400 (2010), 209 400 (2015), 136 100 (2019)

**Number of any first-line antimalarial treatment courses (incl. ACT) distributed:** 1.25 million (2010), 669 000 (2015), 1 110 000 (2019)

<sup>a</sup> Number of piperonyl butoxide (PBO) nets distributed is reported in 2019.

## REPORTED CASES AND DEATHS IN PUBLIC SECTOR, 2010–2019

**Total (presumed and confirmed) cases:** 677 100 (2010), 434 000 (2015), 723 000 (2019)

**Confirmed cases:** 677 100 (2010), 434 000 (2015), 723 000 (2019)

**Percentage of total cases confirmed:** 100% (2010), 100% (2015), 100% (2019)

**Deaths:** 190 (2010), 98 (2015), 197 (2019)

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:** 821 000 (2010), 561 000 (2015), 889 000 (2019); increase 2010–2019: 8%

**Deaths:** 510 (2010), 400 (2015), 550 (2019); increase 2010–2019: 9%

## ACCELERATION TO ELIMINATION

**Countries part of the E-2020 initiative:** Belize, Costa Rica, Ecuador, El Salvador, Mexico, Paraguay and Suriname

**Zero indigenous cases for 3 consecutive years (2017, 2018 and 2019):** El Salvador

**Zero indigenous cases in 2019:** Belize and El Salvador


**Certified as malaria free since 2010:** Argentina (2019) and Paraguay (2018)

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH P. FALCIPARUM MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2011–2019	7	0.0	0.0	0.0	0.0	0.0
AS-MQ	2010–2017	6	0.0	0.0	0.0	0.0	0.0

AL: artemether-lumefantrine; AS-MQ: artesunate-mefloquine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. Confirmed malaria cases per 1000 population, 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; USAID: United States Agency for International Development.


<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

## C. Malaria funding<sup>a</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.


#### D. Share of estimated malaria cases, 2019




#### E. Percentage of *Plasmodium* species from indigenous cases, 2010 and 2019




#### F. Estimated number of cases in countries and areas on track to reduce case incidence by $\geq 40\%$ by 2020



#### G. Estimated number of cases in countries with an increase in case incidence, 2015–2019




#### H. Change in estimated malaria incidence and mortality rates, 2015–2019




<sup>a</sup> These countries and areas (plus Costa Rica) already achieved the 40% reduction in mortality rate in 2015 and since then, there has been no change.  
<sup>b</sup> These countries used reported deaths for mortality.

#### I. Percentage of total confirmed cases investigated, 2019



Note: Countries and areas with no reported case investigation: Bolivia (Plurinational State of), French Guiana, Guyana and Haiti.

#### J. Number of reported indigenous cases in countries with national elimination activities, 2015 versus 2019



## KEY MESSAGES

- About 139 million people in 18 countries in the WHO Region of the Americas are at risk of malaria, most of which (almost 75%) is caused by *P. vivax*. In 2019, the region reported 723 025 malaria cases – a 7% increase from 2010 – and 197 deaths – a 4% increase from 2010. Three countries accounted for almost 90% of all reported cases: Venezuela (Bolivarian Republic of) (55%), Brazil (22%) and Colombia (11%). Malaria prevention in most of the countries relies on IRS, or mass or routine distribution of bed nets. Bolivia (Plurinational State of) was the only country that introduced the distribution of piperonyl butoxide (PBO) nets in 2019.
- Seven of the endemic countries in the region are on target to reduce estimated case incidence by more than 40% by 2020. Twelve countries – Bolivia (Plurinational State of), Brazil, Colombia, Costa Rica, Dominican Republic, Ecuador, Guyana, Mexico, Nicaragua, Panama, Suriname and Venezuela (Bolivarian Republic of) – saw increases in incidence in 2019 compared with 2015. Additionally, Colombia, Guatemala, Haiti, Honduras and Peru experienced a reduction in the number of estimated deaths larger than 40%, while another nine countries reported zero malaria deaths.
- Seven countries experienced a reduction in the number of reported cases between 2015 and 2019: Belize (100% reduction), El Salvador (100% reduction), French Guiana (53% reduction), Guatemala (63% reduction), Haiti (39% reduction), Honduras (91% reduction) and Peru (63% reduction). All other countries experienced varying levels of increases in reported cases. Nevertheless, transmission in countries was focal – in particular, in Choco in Colombia, Loreto in Peru and Bolivar in Venezuela (Bolivarian Republic of) – with more than one third of all cases in the region in 2018 being from

- eight municipalities. Increases in other countries in 2019 are attributed to improved surveillance and focal outbreaks.
- All the local cases reported by Guatemala, Mexico and Suriname were due to *P. vivax*. Additionally, between 60% and 99% of the local cases were due to *P. vivax* in Bolivia (Plurinational State of), Brazil, Ecuador, French Guiana, Guyana, Honduras, Nicaragua, Panama, Peru and Venezuela (Bolivarian Republic of). Conversely, all local cases reported by the Dominican Republic and Haiti were due to *P. falciparum*, and 71% of the local cases reported in Colombia in 2019 were due to *P. falciparum*.
- Seven countries in this region are part of the E-2020 initiative: Belize, Costa Rica, Ecuador, El Salvador, Mexico, Paraguay and Suriname. Paraguay and Argentina were certified malaria free by WHO in 2018 and 2019, respectively. In 2019, imported cases accounted for 100% of the cases in Belize (2/2) and El Salvador (3/3), 52% of the cases in Suriname (111/215), 31% of the cases in Costa Rica (45/145), 5% of the cases in Ecuador (106/1906) and 3% of the cases in Mexico (22/641). Additionally, nine countries in Central America and Hispaniola are taking part in the subregional initiative to eliminate malaria by 2020.
- Vector resistance to pyrethroids was confirmed in 26% of the sites, to organochlorines in 4%, to carbamates in 17% and to organophosphates in 19%. Significant gaps remain in standard resistance monitoring for all of the five insecticide classes commonly used for vector control. Nine countries have developed insecticide resistance monitoring and management plans.

# Annex 2 – C. WHO Eastern Mediterranean Region

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 324 million

**Parasites:** *P. falciparum* and mixed (73%), *P. vivax* (27%) and other (<1%)

**Vectors:** *An. annularis*, *An. arabiensis*, *An. culicifacies s.l.*, *An. d'thali*, *An. fluviatilis s.l.*, *An. funestus s.l.*, *An. gambiae s.s.*, *An. maculipennis s.l.*, *An. merus*, *An. pulcherrimus*, *An. sacharovi*, *An. sergentii*, *An. stephensi* and *An. superpictus s.l.*

## FUNDING (US\$), 2010–2019

130.1 million (2010), 160.2 million (2015), 128.9 million (2019); decrease 2010–2019: 1%

**Proportion of domestic source<sup>a,b</sup> in 2019:** 29%

**Regional funding mechanisms:** none

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

<sup>b</sup> No domestic funding data reported for Afghanistan, Sudan and Yemen in 2019.

## INTERVENTIONS, 2010–2019

**Number of people protected by IRS:<sup>a</sup>** 10.5 million (2010), 27.8 million (2015), 7.9 million (2019)

**Total LLINs distributed:<sup>a</sup>** 2.8 million (2010), 5.7 million (2015), 13.5 million (2019)

**Number of RDTs distributed:** 2.0 million (2010), 6.1 million (2015), 14.2 million (2019)

**Number of ACT courses distributed:** 2.6 million (2010), 3.2 million (2015), 4.7 million (2019)

**Number of any first-line antimalarial treatment courses (incl. ACT) distributed:** 2.6 million (2010), 4.0 million (2015), 5.4 million (2019)

<sup>a</sup> No data reported for Pakistan in 2010.

## REPORTED CASES AND DEATHS IN PUBLIC SECTOR, 2010–2019

**Total (presumed and confirmed) cases:<sup>a</sup>** 6.4 million (2010), 5.4 million (2015), 4.5 million (2019)

**Confirmed cases:** 1.2 million (2010), 1.0 million (2015), 2.6 million (2019)

**Percentage of total cases confirmed:** 18.3% (2010), 18.5% (2015), 57.8% (2019)

**Deaths:<sup>b</sup>** 1140 (2010), 1020 (2015), 1690 (2019)

<sup>a</sup> Figures include imported cases. In 2019, 0 and 38 indigenous cases were reported in Iran (Islamic Republic of) and Saudi Arabia, respectively.

<sup>b</sup> In 2019, there was no report on malaria deaths in Pakistan.

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:** 5.0 million (2010), 4.1 million (2015), 5.2 million (2019); increase 2010–2019: 15%

**Deaths:** 8720 (2010), 7880 (2015), 10 130 (2019); increase 2010–2019: 16%

## ACCELERATION TO ELIMINATION

**Countries with nationwide elimination programme:** Iran (Islamic Republic of) and Saudi Arabia

**Zero indigenous cases in 2019:** Iran (Islamic Republic of)


**Certified as malaria free since 2010:** Morocco (2010)

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH *P. FALCIPARUM* MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2010–2018	32	0.0	0.0	7.9	0.0	2.0
AS+SP	2010–2017	42	0.0	1.0	22.2	0.0	4.4
DHA-PPQ	2015–2017	8	0.0	0.0	2.5	0.0	1.4

AL: artemether-lumefantrine; AS+SP: artesunate-sulfadoxine-pyrimethamine; DHA-PPQ: dihydroartemisinin-piperazine.



## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. *P. falciparum* parasite prevalence (PfPP)/confirmed malaria cases per 1000 population, 2019



## B. Malaria funding<sup>a,b</sup> by source, 2010–2019




Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland; USAID: United States Agency for International Development.

<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

<sup>b</sup> No domestic funding data reported for Afghanistan, Sudan and Yemen in 2019.


## C. Malaria funding<sup>a,b</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.

<sup>b</sup> No domestic funding data reported for Afghanistan, Sudan and Yemen in 2019.


### D. Share of estimated malaria cases, 2019




### E. Percentage of *Plasmodium* species from indigenous cases, 2010 and 2019




### F. Countries with an increase in reported cases, 2015–2019




### G. Change in estimated malaria incidence and mortality rates, 2015–2019



### H. Percentage of total confirmed cases investigated, 2019



### I. Reported indigenous cases in countries with national elimination activities, 2015 versus 2019



## KEY MESSAGES

- Fourteen countries in the WHO Eastern Mediterranean Region are free of indigenous malaria and are at the stage of prevention of re-establishment. There are eight malaria endemic countries in the region, and *P. falciparum* is responsible for 73% of all detected infections. Estimated malaria incidence in the region declined between 2010 and 2015 but increased over the past 4 years, translating into a 15% increase between 2010 and 2019. The number of estimated malaria deaths also increased, in this case by 16% between 2010 and 2019.
- Sudan and Yemen accounted for about two thirds of the cases estimated for the region. In 2019, the region reported that about 2.6 million of the 4.5 million cases reported were confirmed (57.8%), which represented an increase from the 46% confirmation rate reported in 2018 and the 18% in 2010. The reported number of deaths increased from 1143 in 2010 to 1690 in 2019.
- The Islamic Republic of Iran and Saudi Arabia are both targeting elimination by 2020. The Islamic Republic of Iran reported zero indigenous cases for the past 2 years (and until October 2020). In Saudi Arabia, the number of indigenous malaria cases declined from 272 in 2016 to 38 in 2019. These countries undertake continued vigilance for malaria in the general health services, and provide diagnosis and treatment free of charge to all imported cases.

- Vector resistance to pyrethroids, organochlorines and organophosphates was confirmed in 76%, 66% and 46% of the sites tested, respectively, in all countries except for Saudi Arabia. Also, 25% of the sites in the region confirmed resistance to carbamates in all countries except for Saudi Arabia and Somalia. Seven countries have developed their insecticide resistance monitoring and management plans.
- Challenges include low coverage of essential interventions (below universal target) in most malaria endemic countries, inadequate funding and dependence on external resources, humanitarian emergencies, difficult operational environments and population displacements, a shortage of skilled technical staff (particularly at subnational level), and weak surveillance and health information systems. Frequent floods – particularly in Somalia, Sudan and Yemen – and the increasing presence of invasive *An. stephensi* in Djibouti, Somalia and Sudan have increased the risk of malaria, particularly in urban and suburban areas. The confirmed presence of HRP2/3 gene deletions in Djibouti and the high probability of the presence of this mutation in Somalia is another threat for the region. These challenges may have led to an overall increase in cases during the period 2015–2019 in some countries of the region.



# Annex 2 – D. WHO South-East Asia Region

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 1.64 billion

**Parasites:** *P. falciparum* and mixed (53%), *P. vivax* (46%) and other (<1%)

**Vectors:** *An. albimanus*, *An. annularis*, *An. balabacensis*, *An. barbirostris*, *An. culicifacies* s.l., *An. dirus* s.l., *An. farauti* s.l., *An. fluviatilis*, *An. leteri*, *An. maculatus* s.l., *An. minimus* s.l., *An. peditaeniatus*, *An. philippinensis*, *An. pseudowillmori*, *An. punctulatus* s.l., *An. sinensis* s.l., *An. stephensi* s.l., *An. subpictus* s.l., *An. sundaicus* s.l., *An. tessellatus*, *An. vagus*, *An. varuna* and *An. yatsushiroensis*.

## FUNDING (US\$), 2010–2019

250.9 million (2010), 201.8 million (2015), 259.9 million (2019); increase 2010–2019: 4%

**Proportion of domestic source<sup>a</sup> in 2019:** 61%

**Regional funding mechanisms:** Mekong Malaria Elimination (MME) initiative in the Greater Mekong subregion: Myanmar and Thailand

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2010–2019

**Number of people protected by IRS:** 76.4 million (2010), 57.2 million (2015), 31.6 million (2019)

**Total LLINs distributed:** 7.4 million (2010), 7.3 million (2015), 34.8 million (2019)

**Number of RDTs distributed:**<sup>a</sup> 11.4 million (2010), 23.5 million (2015), 6.6 million (2019)

**Number of ACT courses distributed:**<sup>b</sup> 3.5 million (2010), 2.8 million (2015), 1.0 million (2019)

**Number of any first-line antimalarial treatment courses (incl. ACT) distributed:**<sup>b</sup>

4.1 million (2010), 2.9 million (2015), 1.1 million (2019)

<sup>a</sup> Data for India were not available for 2019.

<sup>b</sup> Distribution numbers were not reported in India in 2019. Numbers for India were assigned based on the total number of cases treated in the country.

## REPORTED CASES AND DEATHS, 2010–2019

**Total (presumed and confirmed) cases:** 3.1 million (2010), 1.6 million (2015), 672 000 (2019)

**Confirmed cases:** 2.6 million (2010), 1.6 million (2015), 671 000 (2019)

**Percentage of total cases confirmed:** 84.8% (2010), 98.4% (2015), 99.9% (2019)

**Deaths:** 2421 (2010), 620 (2015), 162 (2019)

<sup>a</sup> Bangladesh, Bhutan, Indonesia, Nepal, Thailand and Timor-Leste included imported cases in 2019.

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:** 24.6 million (2010), 13.3 million (2015), 6.3 million (2019); decrease 2010–2019: 74%

**Deaths:** 38 300 (2010), 24 100 (2015), 9000 (2019); decrease 2010–2019: 76%

## ACCELERATION TO ELIMINATION

**Countries with subnational/territorial elimination programme:** Bangladesh, India, Indonesia, Myanmar and Thailand

**Countries with nationwide elimination programme:** Bhutan, Democratic People's Republic of Korea, Nepal and Timor-Leste

**Zero indigenous cases in 2019:** Timor-Leste


**Certified as malaria free since 2010:** Maldives (2015) and Sri Lanka (2016)

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH P. FALCIPARUM MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile 25	Percentile 75
AL	2010–2019	88	0.0	0.0	14.3	0.0	1.9
AS+SP	2010–2017	56	0.0	0.0	25.9	0.0	1.5
AS-MQ	2010–2016	23	0.0	2.1	49.1	0.0	15.6
DHA-PPQ	2010–2018	33	0.0	0.0	100.0	0.0	2.0

AL: artemether-lumefantrine; AS-MQ: artesunate-mefloquine; AS+SP: artesunate+sulfadoxine-pyrimethamine; DHA-PPQ: dihydroartemisinin-piperaquine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. Confirmed malaria cases per 1000 population, 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland; USAID: United States Agency for International Development.


<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.

## C. Malaria funding<sup>a</sup> per person at risk, average 2017–2019




<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.



#### D. Share of estimated malaria cases, 2019




#### E. Percentage of *Plasmodium* species from indigenous cases, 2010 and 2019




#### F. Estimated number of cases in countries on track to reduce case incidence by ≥40% by 2020




#### G. Change in estimated malaria incidence and mortality rates, 2015–2019



#### H. Percentage of total confirmed cases investigated,<sup>a</sup> 2019



#### I. Reported indigenous cases in countries with national elimination activities, 2015 versus 2019



<sup>a</sup> These countries already achieved the 40% reduction in mortality rate in 2015; since then, there has been no change.

<sup>b</sup> Reported confirmed cases are used for these countries (as opposed to estimated cases).

<sup>a</sup> Country with no reported case investigation: India.

<sup>a</sup> Not all confirmed cases underwent case investigation in these countries.

## KEY MESSAGES

- An estimated 1.64 billion people in the WHO South-East Asia Region are at risk of malaria. The disease is endemic in nine of the region's 11 countries, accounting for nearly 50% of the burden of malaria outside the WHO African Region. In 2019, the region had 6.3 million estimated cases and 9000 estimated deaths – reductions of 73% and 74%, respectively, compared with 2000 – representing the largest decline among all regions. All countries are on target to achieve a more than 40% reduction in case incidence and mortality rate by 2020 compared with 2015, except Indonesia where the mortality rate reduced by 37%.
- Three countries accounted for 99.5% of the estimated cases in the region, India being the largest contributor (87.9%), followed by Indonesia (10.4%) and Myanmar (1.2%). Despite being the highest burden country of the region, in 2019, India recorded a 60% reduction in reported cases compared with 2017 and a 46% reduction compared with 2018. Two other countries in the region recorded substantial declines in total reported cases between 2018 and 2019: Democratic People's Republic of Korea (49% reduction) and Nepal (40% reduction).
- Continuing the declining trend, reported malaria deaths in the region dropped to 162 in 2019 – a 93% reduction compared with 2010. India, Indonesia and Myanmar accounted for 48%, 30% and 9% of the total reported deaths in the region, respectively. Bhutan, Democratic People's Republic of Korea, Nepal and Timor-Leste continue to record zero indigenous deaths.

- Three countries in this region aimed to eliminate malaria by 2020: Bhutan, Nepal and Timor-Leste. Timor-Leste continued to be free of indigenous malaria for the second successive year, while Bhutan reported just two indigenous cases in 2019. For both countries, these reductions in indigenous cases represent significant achievements compared with 2015 (100% and 94% reductions in reported cases, respectively). The majority of reported cases in these countries were imported: Bhutan at 71% (30/42), Nepal at 82% (579/710) and Timor-Leste at 100% (9/9). Maldives and Sri Lanka, which were certified as malaria free in 2015 and 2016, respectively, continue to maintain their malaria free status.
- Vector resistance to pyrethroids was confirmed in 50% of the sites, to organochlorines in 76%, to carbamates in 49% and to organophosphates in 57.5%. There remain significant gaps in standard resistance monitoring. Four countries have developed insecticide resistance monitoring and management plans.
- Challenges include decreased funding, multiple artemisinin-based combination therapy failures in the countries of the Greater Mekong subregion (GMS) and vector resistance to pyrethroids. Efforts are underway to strengthen surveillance and enhance reporting from private sector and nongovernmental organizations where relevant, and case-based surveillance and response to accelerate towards elimination. Imported malaria is an increasingly critical challenge for those countries that are on the verge of malaria elimination.

# Annex 2 – E. WHO Western Pacific Region

## EPIDEMIOLOGY

**Population denominator used to compute incidence and mortality rate:** 767 million

**Parasites:** *P. falciparum* and mixed (68%), *P. vivax* (32%) and other (<1%)

**Vectors:** *An. anthropophagus*, *An. balabacensis*, *An. barbirostris s.l.*, *An. dirus s.l.*, *An. donaldi*, *An. epirotivulus*, *An. farauti s.l.*, *An. flavirostris*, *An. jeyporiensis*, *An. koliensis*, *An. litoralis*, *An. maculatus s.l.*, *An. mangyanus*, *An. minimus s.l.*, *An. punctulatus s.l.*, *An. sinensis s.l.* and *An. sundaicus s.l.*

## FUNDING (US\$), 2010–2019

211.6 million (2010), 146.3 million (2015), 141.8 million (2019); decrease 2010–2019: 33%

**Proportion of domestic source<sup>a</sup> in 2019:** 53%

**Regional funding mechanisms:** Mekong Malaria Elimination (MME) initiative in the Greater Mekong subregion: Cambodia, China (Yunnan), Lao People's Democratic Republic and Viet Nam (supported by RAI2e Global Fund)

<sup>a</sup> Domestic source excludes patient service delivery costs and out-of-pocket expenditure.

## INTERVENTIONS, 2010–2019

**Number of people protected by IRS:** 27.9 million (2010), 3.3 million (2015), 1.8 million (2019)

**Total LLINs distributed:** 3.4 million (2010), 2.7 million (2015), 3.8 million (2019)

**Number of RDTs distributed:** 1.6 million (2010), 2.5 million (2015), 6.1 million (2019)

**Number of ACT courses distributed:** 591 000 (2010), 1.3 million (2015), 1.7 million (2019)

**Number of any antimalarial treatment courses (incl. ACT) distributed:** 963 000 (2010), 1.4 million (2015), 1.8 million (2019)

## REPORTED CASES<sup>a,b</sup> AND DEATHS IN PUBLIC SECTOR, 2010–2019

**Total (presumed and confirmed) cases:** 1.6 million (2010), 704 000 (2015), 789 000 (2019)

**Confirmed cases:** 260 000 (2010), 411 000 (2015), 769 000 (2019)

**Percentage of total cases confirmed:** 15.8% (2010), 58.3% (2015), 97.5% (2019)

**Deaths:** 910 (2010), 235 (2015), 229 (2019)

<sup>a</sup> China, Malaysia, Philippines, Republic of Korea, Vanuatu and Viet Nam included imported cases for 2019. China has had no indigenous malaria since 2017.

<sup>b</sup> In Malaysia, figures for 2015 and 2019 included indigenous *P. knowlesi* cases. All indigenous malaria cases observed since 2018 have been *P. knowlesi*.

## ESTIMATED CASES AND DEATHS, 2010–2019

**Cases:** 1.8 million (2010), 1.4 million (2015), 1.7 million (2019); decrease 2010–2019: 5%

**Deaths:** 3780 (2010), 2780 (2015), 3160 (2019); decrease 2010–2019: 16%

## ACCELERATION TO ELIMINATION

**Countries with subnational/territorial elimination programme:** Philippines

**Countries with nationwide elimination programme:** Cambodia, China, Lao People's Democratic Republic, Malaysia, Republic of Korea, Vanuatu and Viet Nam

**Zero indigenous cases for 3 consecutive years (2017, 2018 and 2019):** China


**Zero indigenous cases in 2019:** China and Malaysia

## THERAPEUTIC EFFICACY STUDIES (CLINICAL AND PARASITOLOGICAL FAILURE AMONG PATIENTS WITH *P. FALCIPARUM* MALARIA, %)

Medicine	Study years	No. of studies	Min.	Median	Max.	Percentile	
						25	75
AL	2010–2019	33	0.0	0.0	17.2	0.0	5.8
AS-MQ	2010–2019	32	0.0	0.0	12.5	0.0	0.0
AS-PY	2014–2019	15	0.0	1.6	18.0	0.0	5.1
DHA-PPQ	2010–2019	84	0.0	1.6	85.7	0.0	17.5

AL: artemether-lumefantrine; AS-MQ: artesunate-mefloquine; AS-PY: artesunate-pyronaridine; DHA-PPQ: dihydroartemisinin-piperaquine.


## STATUS OF INSECTICIDE RESISTANCE<sup>a</sup> PER INSECTICIDE CLASS (2010–2019) AND USE OF EACH CLASS FOR MALARIA VECTOR CONTROL (2019)




<sup>a</sup> Resistance is considered confirmed when it was detected to one insecticide in the class, in at least one malaria vector from one collection site.

<sup>b</sup> Number of countries that reported using the insecticide class for malaria vector control (2019).

## A. Confirmed malaria cases per 1000 population, 2019




## B. Malaria funding<sup>a</sup> by source, 2010–2019



Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; UK: United Kingdom of Great Britain and Northern Ireland; USAID: United States Agency for International Development.

<sup>a</sup> Excludes patient service delivery costs and out-of-pocket expenditure.


## C. Malaria funding<sup>a,b</sup> per person at risk, average 2017–2019



<sup>a</sup> Excludes costs related to health staff, costs at subnational level and out-of-pocket expenditure.


<sup>b</sup> Only domestic funding in Malaysia and the Republic of Korea.

## D. Share of estimated malaria cases, 2019




Note: Countries with zero cases: China and Malaysia.


## E. Percentage of *Plasmodium* species from indigenous cases, 2010 and 2019




## F. Estimated number of cases in countries on track to reduce case incidence by ≥40% by 2020




## G. Estimated number of cases in countries likely to reduce case incidence by <40% by 2020



## H. Estimated number of cases in countries with an increase in case incidence, 2015–2019




## I. Change in estimated malaria incidence and mortality rates, 2015–2019




<sup>a</sup> There have been no estimated indigenous deaths between 2015 and 2019 in these countries.

## J. Percentage of total confirmed cases investigated, 2019



Notes: Imported cases are included. No case investigation in Papua New Guinea and Solomon Islands.

## K. Reported indigenous cases in countries with national elimination activities, 2015 versus 2019



## KEY MESSAGES

- About 767 million people in 10 countries in the WHO Western Pacific Region are at risk of malaria, which is predominantly caused by *P. falciparum* (66%), followed by *P. vivax* (32%). In 2019, the region had more than 1.7 million estimated malaria cases and about 3160 estimated deaths – a 5% increase and 16% reduction from 2010, respectively. Most cases occurred in Papua New Guinea (79%) which, together with Solomon Islands (9%) and Cambodia (8%), comprised 96% of the estimated cases in the region. Almost 789 000 cases were reported in the public and private sectors and in the community, of which almost 98% were confirmed. This was a significant improvement over 2018, when only 59% of cases were confirmed. There were 229 malaria deaths reported in the region in 2019.
- Of the 10 malaria endemic countries in the region, four are on target to achieve more than a 40% reduction in case incidence by 2020, including Cambodia, China, Lao People's Democratic Republic and Malaysia, whereas the Republic of Korea and Viet Nam are on track for a 20–40% reduction. Countries that have experienced an increase in estimated cases since 2015 are Papua New Guinea (32%), the Philippines (29%), Solomon Islands (270%) and Vanuatu (20%). All countries are on track to reduce the malaria mortality rate by at least 40% by 2020, except for Papua New Guinea, the Philippines and Solomon Islands.
- China and Malaysia are on track for elimination of malaria by 2020. China has reported zero indigenous cases for 3 consecutive years (since 2017) and Malaysia has reported zero indigenous human malaria cases since 2018. However, Malaysia is facing increasing cases of zoonotic malaria due to *P. knowlesi*, which increased from 1600 cases to over 4000 between 2016 and 2018. *P. knowlesi* cases have slightly declined (to 3213 cases), but resulted in 12 deaths in 2018–2019. The Republic of Korea continues to face the challenge of malaria transmission among military

- personnel along the northern border. The Philippines has continued its subnational elimination efforts, reporting zero indigenous cases in 78 out of 81 provinces.
- Three countries of the GMS (Cambodia, Lao People's Democratic Republic and Viet Nam) – supported through a regional artemisinin-resistance initiative financed by the Global Fund – aim to eliminate *P. falciparum* by 2020 and all species of malaria by 2030. The percentage of reported cases in Cambodia due to *P. falciparum* has fallen significantly, from 61% in 2015 to 17% in 2019, owing to intensified efforts in community outreach and active case detection. Although the goal of *P. falciparum* elimination will not be met by 2020, and targets will be delayed for a few years, much progress continues to be made.
- Vector resistance to pyrethroids was confirmed in 49% of the sites, to organochlorines in 67%, to carbamates in 36% and to organophosphates in 64%. Almost no standard resistance monitoring was reported for carbamates or organophosphates other than in China, the Philippines and Solomon Islands. Five countries have developed their insecticide resistance monitoring and management plans.
- Challenges include decreased funding, some vector resistance to pyrethroids, resurgence of malaria in Solomon Islands and sustained high levels of malaria in Papua New Guinea due to challenges in health system strengthening. Recent efforts are underway to improve access to services and case-based surveillance in the Pacific Island countries, and intensified community efforts to halt malaria transmission in the GMS countries, particularly in Cambodia. Although all countries have reported minor disruptions to implementing malaria interventions due to COVID-19, no major delays to service delivery have been reported.

## ANNEX 3 – A. POLICY ADOPTION, 2019

WHO region Country/area	Insecticide-treated mosquito nets				Indoor residual spraying		Chemoprevention	
	ITNs/LLINs are distributed free of charge	ITNs/LLINs are distributed through ANC	ITNs/LLINs distributed through EPI/well baby clinic	ITNs/LLINs distributed through mass campaigns	IRS is recommended by malaria control programme	DDT is used for IRS	IPTp is used to prevent malaria during pregnancy	SMC or IPTc is used
<b>AFRICAN</b>								
Angola	●	●	●	●	●	●	●	●
Benin	●	●	●	●	●	●	●	●
Botswana	●	●	●	●	●	●	NA	NA
Burkina Faso	●	●	●	●	●	●	●	●
Burundi	●	●	●	●	●	●	●	NA
Cabo Verde	NA	NA	NA	NA	●	●	NA	NA
Cameroon	●	●	●	●	●	NA	●	●
Central African Republic	●	●	●	●	●	●	●	●
Chad	●	●	●	●	●	●	●	●
Comoros	●	●	●	●	●	●	●	NA
Congo	●	●	●	●	●	●	●	NA
Côte d'Ivoire	●	●	●	●	●	●	●	NA
Democratic Republic of the Congo	●	●	●	●	●	●	●	NA
Equatorial Guinea	●	●	●	●	●	●	●	●
Eritrea	●	●	●	●	●	●	●	●
Eswatini	●	NA	NA	●	●	●	NA	NA
Ethiopia	●	●	●	●	●	●	●	NA
Gabon	●	●	●	●	●	●	●	●
Gambia	●	●	●	●	●	●	●	●
Ghana	●	●	●	●	●	●	●	●
Guinea	●	●	●	●	●	●	●	●
Guinea-Bissau	●	●	●	●	●	●	●	●
Kenya	●	●	●	●	●	NA	●	●
Liberia	●	●	●	●	●	●	●	●
Madagascar	●	●	●	●	●	●	●	●
Malawi	●	●	●	●	●	●	●	●
Mali	●	●	●	●	●	NA	●	●
Mauritania	●	●	●	●	●	●	●	●
Mayotte	-	-	-	-	-	●	NA	NA
Mozambique	●	●	●	●	●	●	●	NA
Namibia	●	NA	NA	●	●	●	●	NA
Niger	●	●	●	●	●	●	●	●
Nigeria	●	●	●	●	●	●	●	●
Rwanda	●	●	●	●	●	●	●	●
Sao Tome and Principe	●	●	NA	●	●	●	●	●
Senegal	●	●	●	●	●	●	●	●
Sierra Leone	●	●	●	●	●	●	●	●
South Africa	●	●	●	●	●	●	●	●
South Sudan <sup>2</sup>	●	●	●	●	●	●	●	●
Togo	●	●	●	●	●	●	●	●
Uganda	●	●	●	●	●	●	●	●
United Republic of Tanzania <sup>3</sup>								
Mainland	●	●	●	●	●	●	●	NA
Zanzibar	●	●	●	●	●	●	●	●
Zambia	●	●	●	●	●	●	●	NA
Zimbabwe	●	●	●	●	●	●	●	●
<b>AMERICAS</b>								
Bolivia (Plurinational State of)	●	●	●	●	●	●	NA	NA
Brazil	●	●	●	●	●	●	●	●
Colombia	●	●	●	●	●	●	●	●
Costa Rica	●	●	●	●	●	●	NA	NA
Dominican Republic	●	●	●	●	●	●	NA	NA
Ecuador	●	●	●	●	●	●	NA	NA



## ANNEX 3 – A. POLICY ADOPTION, 2019

WHO region Country/area	Insecticide-treated mosquito nets				Indoor residual spraying		Chemoprevention	
	ITNs/LLINs are distributed free of charge	ITNs/LLINs are distributed through ANC	ITNs/LLINs distributed through EPI/well baby clinic	ITNs/LLINs distributed through mass campaigns	IRS is recommended by malaria control programme	DDT is used for IRS	IPTp is used to prevent malaria during pregnancy	SMC or IPTc is used
<b>AMERICAS</b>								
El Salvador	●	●	●	●	●	●	NA	NA
French Guiana	●	●	●	●	●	●	NA	NA
Guatemala	●	●	●	●	●	●	●	●
Guyana	●	●	●	●	●	NA	NA	NA
Haiti	●	●	●	●	●	●	NA	NA
Honduras	●	●	●	●	●	●	NA	NA
Mexico	●	●	●	●	●	●	●	●
Nicaragua	●	●	●	●	●	●	NA	NA
Panama	●	●	●	●	●	●	NA	NA
Peru	●	●	●	●	●	●	NA	NA
Suriname	●	●	●	●	●	●	●	●
Venezuela (Bolivarian Republic of)	●	●	●	●	●	●	NA	NA
<b>EASTERN MEDITERRANEAN</b>								
Afghanistan	●	●	●	●	●	●	NA	NA
Djibouti	●	●	●	●	●	●	NA	NA
Iran (Islamic Republic of)	●	●	●	●	●	●	NA	NA
Pakistan	●	●	●	●	●	●	NA	NA
Saudi Arabia	●	●	●	●	●	●	NA	NA
Somalia	●	●	●	●	●	●	●	NA
Sudan	●	●	●	●	●	NA	●	NA
Yemen	●	●	●	●	●	●	NA	NA
<b>SOUTH-EAST ASIA</b>								
Bangladesh	●	●	●	●	●	●	NA	NA
Bhutan	●	●	●	●	●	●	NA	NA
Democratic People's Republic of Korea	●	●	NA	●	●	●	NA	NA
India	●	●	●	●	●	●	NA	NA
Indonesia	●	●	●	●	●	●	NA	NA
Myanmar	●	●	●	●	●	●	NA	NA
Nepal	●	●	●	●	●	●	●	●
Thailand	●	●	●	●	●	●	NA	NA
Timor-Leste	●	●	●	●	●	●	NA	NA
<b>WESTERN PACIFIC</b>								
Cambodia	●	●	●	●	●	●	NA	NA
China	●	●	●	●	●	●	NA	NA
Lao People's Democratic Republic	●	●	●	●	●	●	NA	NA
Malaysia	●	●	●	●	●	●	NA	NA
Papua New Guinea	●	●	●	●	●	●	●	NA
Philippines	●	●	●	●	●	●	NA	NA
Republic of Korea	●	NA	NA	●	●	●	NA	NA
Solomon Islands	●	●	●	●	●	●	NA	NA
Vanuatu	●	●	●	●	●	●	NA	NA
Viet Nam	●	NA	NA	●	●	●	NA	NA

ACT: artemisinin-based combination therapy; ANC: antenatal care; DDT: dichlorodiphenyltrichloroethane; EPI: Expanded Programme on Immunization; G6PD: glucose-6-phosphate dehydrogenase; IM: intramuscular; IPTc: intermittent preventive treatment in children; IPTp: intermittent preventive treatment in pregnancy; IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; LLIN: long-lasting insecticidal net; RDT: rapid diagnostic test; SMC: seasonal malaria chemoprevention; WHO: World Health Organization.

<sup>1</sup> Single dose of primaquine (0.75 mg base/kg) for countries in the WHO Region of the Americas.

<sup>2</sup> In May 2013, South Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [https://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

<sup>3</sup> Where national data for the United Republic of Tanzania are unavailable, refer to Mainland and Zanzibar.





## ANNEX 3 – B. ANTIMALARIAL DRUG POLICY, 2019

WHO region Country/area	<i>P. falciparum</i>				<i>P. vivax</i>
	Uncomplicated unconfirmed	Uncomplicated confirmed	Severe	Prevention during pregnancy	Treatment
<b>AFRICAN</b>					
Angola	AL	AL	AS	SP(IPT)	AL
Benin	-	AL	AS	SP(IPT)	-
Botswana	-	AL+PQ	AS	SP(IPT)	AL
Burkina Faso	AL	AL	AS; QN	SP(IPT)	-
Burundi	AL	AL	AS	SP(IPT)	-
Cabo Verde	-	-	-	-	-
Cameroon	-	AL; DHA-PPQ; AS+AQ	AS	SP(IPT)	-
Central African Republic	AL	AL	AS	SP	-
Chad	-	AS+AQ; AL	AS	SP(IPT)	-
Comoros	-	-	-	-	-
Congo	AS+AQ	AS+AQ	AS	SP(IPT)	-
Democratic Republic of the Congo	AS+AQ	AS+AQ; AL	AS; QN	SP(IPT)	-
Equatorial Guinea	AS+AQ	-	AS	SP(IPT)	-
Eritrea	AS+AQ	AS+AQ	AS	-	AS+AQ
Eswatini	-	AL	AS	-	PQ
Ethiopia	AL	AL+PQ	AS	-	CQ+PQ
Gabon	AS+AQ; AL	AS+AQ; AL	AS	SP(IPT)	-
Gambia	AL	AL	AS	SP(IPT)	-
Ghana	-	-	-	-	-
Guinea	AS	AS	AS	SP	-
Guinea-Bissau	-	-	-	-	-
Kenya	AL	AL	AS	SP(IPT)	PQ
Liberia	-	-	-	-	-
Madagascar	AS+AQ	AS+AQ	AS	SP(IPT)	AS+AQ
Malawi	AL	AL	AS	SP(IPT)	-
Mali	AL	AL	AS	SP(IPT)	-
Mauritania	AS+AQ	AS+AQ	AS	SP(IPT)	AS+AQ+PQ
Mayotte	-	-	-	-	-
Mozambique	-	-	-	-	-
Namibia	-	-	-	-	-
Niger	AL	AL	AS; QN	SP(IPT)	-
Nigeria	-	-	-	-	-
Rwanda	AL	AL	AS; QN	-	-
Sao Tome and Principe	AS+AQ	AS+AQ	AS	SP(IPT)	PQ
Senegal	-	AS+AQ; AL	AS	SP(IPT)	-
Sierra Leone	AS+AQ	AL; AS+AQ	AS; AM; QN	SP(IPT)	-
South Africa	AL	AL	AS; QN	-	AL
South Sudan <sup>1</sup>	-	-	-	-	-
Togo	-	-	-	-	-
Uganda	AL	AL	AS	SP(IPT)	-
United Republic of Tanzania	-	-	-	-	-
Mainland	AL	AL	AS; AM; QN	SP(IPT)	-
Zanzibar	AS+AQ	AS+AQ	AS	-	PQ
Zambia	AL	AL	AS	SP(IPT)	-
Zimbabwe	-	AL	AS	SP(IPT)	-
<b>AMERICAS</b>					
Belize	-	CQ+PQ	AL, QN	-	CQ+PQ
Bolivia (Plurinational State of)	-	AL+PQ	AS	-	CQ+PQ
Brazil	-	AL+PQ; AS+MQ+PQ	AS	-	CQ+PQ
Colombia	-	AL	AS	-	CQ+PQ

## ANNEX 3 – B. ANTIMALARIAL DRUG POLICY, 2019

WHO region Country/area	<i>P. falciparum</i>				<i>P. vivax</i>
	Uncomplicated unconfirmed	Uncomplicated confirmed	Severe	Prevention during pregnancy	Treatment
<b>AMERICAS</b>					
Costa Rica	-	CQ+PQ	AL	-	CQ+PQ
Dominican Republic	-	CQ+PQ	AS	-	CQ+PQ
Ecuador	-	AL+PQ	AS	-	CQ+PQ
El Salvador	-	AL+PQ	AL	QN	CQ+PQ
French Guiana	-	AL+PQ	AS	-	CQ+PQ
Guatemala	-	CQ+PQ	CQ+PQ	-	CQ+PQ
Guyana	-	AL+PQ	AS; AQ	CQ	CQ+PQ
Haiti	-	CQ+PQ	AS	-	CQ+PQ
Honduras	-	CQ+PQ	AS	-	CQ+PQ
Mexico	-	AL+PQ	AM+AL	-	CQ+PQ
Nicaragua	-	CQ+PQ	AS	-	CQ+PQ
Panama	-	AL+PQ	AS	-	CQ+PQ
Peru	-	AS+MQ+PQ	AS	-	CQ+PQ
Suriname	-	AL+PQ	AS	-	CQ+PQ
Venezuela (Bolivarian Republic of)	-	AL+PQ	AS	-	CQ+PQ
<b>EASTERN MEDITERRANEAN</b>					
Afghanistan	CQ	AL+PQ	AS; AM; QN	CQ	CQ+PQ
Djibouti	AL	AL+PQ	AS	-	AL+PQ
Iran (Islamic Republic of)	-	-	-	-	-
Pakistan	CQ+PQ	AL+PQ	AS	-	CQ+PQ
Saudi Arabia	-	AS+SP+PQ	AS+AM+QN	-	CQ+PQ
Somalia	AL	AL	AS	SP(IPT)	AL+PQ
Sudan	-	AL	AS; QN	-	AL+PQ
<b>SOUTH-EAST ASIA</b>					
Bangladesh	-	AL+PQ	AS+AL+PQ	-	CQ+PQ
Bhutan	-	AL	AM; QN	-	CQ+PQ
Democratic People's Republic of Korea	-	-	-	-	CQ+PQ
India	-	AS+SP+PQ; AL+PQ	AM; AS; QN	-	CQ+PQ
Indonesia	-	DHA-PPQ	AS	-	DHA+PPQ
Myanmar	-	AL+PQ	AS; AM; QN	-	CQ+PQ
Nepal	-	AL	AS	-	CQ+PQ
Thailand	-	DHA-PPQ; AS-PYR	AS	-	CQ+PQ
Timor-Leste	AL+PQ	AL+PQ	AS; QN	-	AL+PQ
<b>WESTERN PACIFIC</b>					
Cambodia	-	AS+MQ	AS	-	AS+MQ+PQ
China	-	ART-PPQ; AS+AQ; DHA-PPQ; PYR	AM; AS; PYR	-	CQ+PQ; PQ+PPQ; ACT+PQ; PYR
Lao People's Democratic Republic	-	AL	AS	-	AL
Malaysia	-	AL	AS	-	AL+PQ
Papua New Guinea	-	AL	AS; AM	SP	AL+PQ
Philippines	-	AL+PQ	AS	-	AL+PQ
Republic of Korea	-	-	-	-	CQ+PQ
Solomon Islands	-	-	-	-	-
Vanuatu	-	AL	AS	CQ	PQ
Viet Nam	DHA-PPQ; PQ	DHA-PPQ; PQ	AS	-	CQ+PQ

Data as of 17 November 2020

ACT: artemisinin-based combination therapy; AL: artemether-lumefantrine; AM: artemether; AQ: amodiaquine; ART: artemisinin; AS: artesunate; AT: atovaquone; CL: clindamycline; CQ: chloroquine; D: doxycycline; DHA: dihydroartemisinin; IPT: intermittent preventive treatment; MQ: mefloquine; NQ: naphroquine; PG: proguanil; PPQ: piperaquine; PQ: primaquine; PYR: pyronaridine; QN: quinine; SP: sulfadoxine-pyrimethamine; T: tetracycline; WHO: World Health Organization.

<sup>1</sup> In May 2013, South Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [http://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](http://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

## ANNEX 3 – C. FUNDING FOR MALARIA CONTROL, 2017–2019

WHO region Country/area	Year	Contributions reported by donors			
		Global Fund <sup>1</sup>	PMI/USAID <sup>2</sup>	World Bank <sup>3</sup>	UK <sup>4</sup>
<b>AFRICAN</b>					
Angola	2017	15 722 726	22 888 423	0	0
	2018	12 335 146	22 383 604	0	0
	2019	5 028 041	22 000 000	0	0
Benin	2017	26 147 674	16 646 126	0	0
	2018	4 825 798	16 278 984	0	0
	2019	14 171 520	17 000 000	0	0
Botswana	2017	1 683 599	0	0	0
	2018	1 501 436	0	0	0
	2019	270 407	0	0	0
Burkina Faso	2017	9 849 157	26 009 571	11 219 204	1 432 054
	2018	33 120 195	25 435 913	9 192 622	1 008 709
	2019	33 269 764	25 000 000	9 035 082	0
Burundi	2017	28 928 791	9 363 446	0	0
	2018	1 837 003	9 156 929	0	0
	2019	31 146 086	8 000 000	0	0
Cabo Verde	2017	241 299	0	0	0
	2018	-19 345	0	0	0
	2019	0	0	0	0
Cameroon	2017	23 622 914	20 807 657	0	0
	2018	17 374 572	22 892 322	0	0
	2019	31 382 534	22 500 000	0	0
Central African Republic	2017	13 760 308	0	0	0
	2018	17 466 536	0	0	0
	2019	11 245 876	0	0	0
Chad	2017	14 521 704	0	0	0
	2018	18 642 602	0	0	0
	2019	38 076 559	0	0	0
Comoros	2017	875 331	0	0	0
	2018	2 338 882	0	0	0
	2019	1 511 064	0	0	0
Congo	2017	0	0	0	0
	2018	1 207 101	0	0	0
	2019	10 283 939	0	0	0
Côte d'Ivoire	2017	31 951 007	26 009 571	0	0
	2018	27 954 008	25 435 913	0	0
	2019	56 987 087	25 000 000	0	0
Democratic Republic of the Congo	2017	131 093 509	52 019 143	0	6 336 451
	2018	78 970 598	50 871 826	0	4 463 262
	2019	117 949 473	50 000 000	0	747 665
Equatorial Guinea	2017	0	0	0	0
	2018	0	0	0	0
	2019	-218 638	0	0	0
Eritrea	2017	13 533 044	0	0	0
	2018	4 875 453	0	0	0
	2019	8 942 830	0	0	0
Eswatini	2017	1 715 924	0	0	0
	2018	589 889	0	0	0
	2019	836 280	0	0	0
Ethiopia	2017	74 957 424	38 494 165	0	0
	2018	37 121 554	36 627 715	0	0
	2019	26 668 897	36 000 000	0	0

Contributions reported by countries

Government (NMP)	Global Fund	World Bank	PMI/USAID	Other bilaterals	WHO	UNICEF	Other contributions <sup>7</sup>
9 020 546	12 023 625		18 000 000		139 995		
46 457 232 <sup>5</sup>	9 578 147		22 000 000		88 217		
1 754 960	2 864 156		20 000 000				
4 395 380	33 122 938	0	9 642 332	3 140	158 723	5 400	
611 841	2 235 811	0	1 419 738	0	21 292	75 628	0
10 889 600	4 670 273	0	4 217 593	0	0	0	0
1 092 695	1 079 069	0	0	0	0	0	0
2 124 880	2 087 088	0	0	0	0	0	0
2 447 859	219 328	0	0	0	0	0	0
15 573 795	9 474 402	5 608 893	13 053 101		164 363	163 431	5 570 878
123 337	14 880 669	5 321 114	16 646 476		431 795	228 084	2 900 368
97 208 057	66 864 802	6 473 917	20 960 657		107 706	546 944	
3 070 872	21 228 086		9 000 000		37 832	4 967 372	869 962
1 157 984	4 734 738		9 000 000		68 488	433 441	4 664 286
4 328 977	24 301 509		4 734 719		159 500	372 925	
4 627 843	466 244				29 000		
621 612	221 609				25 641		
519 158	116 809				82 598		
2 288 193 <sup>5</sup>	28 008 486				882 650	1 105 377	9 477
10 607 209 <sup>5</sup>	47 200 683		29 913 228				
61 194 530 <sup>5</sup>	33 828 144	0	21 148 951	0	0	0	0
530 000	443 466				70 419		
675 455	8 399 445				50 000	306 968	
0	16 631 715				199 800	656 890	
652 320 <sup>6</sup>	34 927 891				416	540 870	867 119
543 725 <sup>6</sup>							
0 <sup>6</sup>							
114 684	852 996	0	0	0	54 000		0
114 684		0	0	0	60 000		0
114 684 <sup>6</sup>	824 954						
122 182	0	0	0	0	15 000	0	10 000
50 509	9 090 909	0	0	0	0	0	9 090
1 290 322	6 689 800	0	0	0	67 741	0	15 000
34 806 734	44 798 740	667 580	0	487 446	17 698	76 943	238 890
32 071 401	28 330 619	877 696	9 151 372	27 724 798	47 903	32 090	435 865
6 097 961	60 947 905		21 342 862		5 984	60 980	2 500 000
683 314	75 183 622	0	46 738 755	4 694 136	2 265 298	82 857	0
1 948 241	92 444 112	0	49 075 000	0	636 951	0	0
1 427 241	112 504 296	0	41 897 052		148 208	802 250	
3 208 473 <sup>6</sup>							
3 208 473 <sup>6</sup>							
3 153 487 <sup>6</sup>							
408 557 <sup>6</sup>	9 150 700	0	0	0	80 450	0	
408 557 <sup>6</sup>	2 748 778	0	0	0	82 500	0	0
401 555 <sup>6</sup>	4 788 233	0	0	0	120 000	0	0
10 019 754	20 910 608	0	0	0	620 000	0	0
989 110	1 376 660	0	0	0		0	0
838 430	2 652 105	0	0	0	10 613	0	0
19 401 447	31 604 918		7 150 000		0	30 000	13 500 000
20 758 465	44 800 000		26 358 971				14 000 000
22 907 737	26 083 562		18 000 000				122 344 828

## ANNEX 3 – C. FUNDING FOR MALARIA CONTROL, 2017–2019

WHO region Country/area	Year	Contributions reported by donors			
		Global Fund <sup>1</sup>	PMI/USAID <sup>2</sup>	World Bank <sup>3</sup>	UK <sup>4</sup>
<b>AFRICAN</b>					
Gabon	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Gambia	2017	10 584 939	0	0	0
	2018	8 128 184	0	0	0
	2019	3 412 032	0	0	0
Ghana	2017	41 546 764	29 130 720	0	1 183 127
	2018	44 934 700	28 488 223	0	833 369
	2019	35 771 452	28 000 000	0	1 462 425
Guinea	2017	14 656 590	15 605 743	568 210	0
	2018	12 752 728	15 261 548	1 174 368	0
	2019	28 982 985	15 000 000	1 154 242	0
Guinea-Bissau	2017	6 856 945	0	0	0
	2018	7 821 002	0	0	0
	2019	4 814 351	0	0	0
Kenya	2017	61 554 420	36 413 400	0	1 031 373
	2018	12 659 098	35 610 278	0	726 478
	2019	33 425 267	35 000 000	0	0
Liberia	2017	14 361 899	14 565 360	0	0
	2018	20 506 609	14 244 111	0	0
	2019	6 394 175	14 000 000	0	0
Madagascar	2017	14 559 438	27 049 954	0	0
	2018	41 069 905	26 453 350	0	0
	2019	6 399 993	26 000 000	0	0
Malawi	2017	12 134 701	22 888 423	0	0
	2018	31 075 220	24 418 477	0	0
	2019	14 464 267	24 000 000	0	0
Mali	2017	23 608 912	26 009 571	5 920 105	0
	2018	31 009 912	25 435 913	11 576 495	0
	2019	21 096 259	25 000 000	11 378 101	0
Mauritania	2017	4 672 266	0	0	0
	2018	4 090 649	0	0	0
	2019	73 220	0	0	0
Mayotte	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Mozambique	2017	64 693 665	30 171 103	2 118 290	7 986 026
	2018	36 396 779	29 505 659	0	5 625 187
	2019	50 895 946	29 000 000	0	0
Namibia	2017	2 754 765	0	0	0
	2018	755 622	0	0	0
	2019	618 414	0	0	0
Niger	2017	25 143 511	18 726 891	6 869 723	0
	2018	28 810 711	18 313 857	5 764 992	0
	2019	21 031 872	18 000 000	5 666 193	0
Nigeria	2017	123 616 145	78 028 714	0	0
	2018	67 768 812	71 220 557	37 237 036	0
	2019	115 283 739	70 000 000	36 598 879	2 522 480
Rwanda	2017	17 364 322	18 726 891	0	0
	2018	10 104 603	18 313 857	0	0
	2019	34 528 150	18 000 000	0	0

Contributions reported by countries

Government (NMP)	Global Fund	World Bank	PMI/USAID	Other bilaterals	WHO	UNICEF	Other contributions <sup>7</sup>
142 296	0	0	0	0	12 616	0	0
0	0	0	0	0	128 016	0	49 674
145 505 <sup>6</sup>	0						
621 025 <sup>6</sup>	9 557 650				14 400	33 839	117 749
1 327 049	8 376 620				39 000	50 414	176 987
1 203 441 <sup>5</sup>	3 940 063				68 000	90 000	288 646
683 179	40 951 105	0	22 445 306		140 000	0	0
140 392 544	47 579 039	0	30 634 694	7 560 000	300 000	0	0
151 756 820	28 442 224	0	22 448 510	0	300 000	0	0
14 796 <sup>5</sup>	9 251 505	125 000	12 500 000		65 000		
6 438 381	12 000 000	156 000	14 000 000		45 000		
951 075	25 261 667		15 000 000		39 000		
1 655 769	9 086 476	0	0	0		0	256 659
651 820	3 199 732	0	0	0		0	0
121 117 307 000	540 184 296						
1 677 914 <sup>6</sup>							
1 677 915 <sup>6</sup>							
6 568 505	14 497 642		34 000 000				
313 801 <sup>6</sup>	18 526 566		14 000 000				
313 801 <sup>6</sup>							
19 621 989	11 500 991	0	12 000 000	0	0	0	0
37 214	43 205 989	0	26 000 000	0	220 000	0	0
13 007	33 200 289	0	26 000 000		46 000		
0 <sup>6</sup>	18 378 714		26 000 000		50 000		
291 194 <sup>5</sup>	16 282 087		22 000 000				
282 401	33 049 389		20 000 000				
317 711	12 768 682				150 000		
4 382 069	19 288 748	3 226 759	25 500 000	0	140 713	854 199	
14 329 420	54 053 651	6 406 499	25 000 000			337 884	
1 273 817	19 414 667	1 085 642	25 000 000	0	24 083	2 420	7 224
605 079 <sup>5</sup>	6 957 945				47 950		13 944
2 191 549	164 778						
124 788	175 296						
76 074	58 222 077		29 000 000		240 000	3 848 028	10 995
2 136 147	45 915 417		29 000 000			1 590 000	4 361 414
1 848 592	62 708 218		29 000 000	39 548 431	414 944	1 102 707	17 667 110
5 166 667	1 096 657	0	0	0	100 000	0	789 566
11 216 160	908 515	0	0	0	100 000	100 000	1 148 515
11 123 042	3 377 753	0	0	0	100 000	0	150 000
4 454 320	22 404 758	2 177 698	220 000	0	328 594	805 598	476 444
7 363 777	20 159 800	4 490 567	18 000 000	0	220 356	674 811	0
1 332 407 <sup>5</sup>	16 329 651	6 319 943	18 000 000	0	86 206	693 054	0
107 005 355	198 176 039		75 000 000				
2 271 631 <sup>6</sup>	43 206 463		70 000 000				
261 799 700	131 373 863		70 000 000				
13 704 611	11 440 292		18 000 000		270 000		
13 460 220	27 505 974		18 000 000				
0 <sup>6</sup>	20 091 346		18 000 000				

## ANNEX 3 – C. FUNDING FOR MALARIA CONTROL, 2017–2019

WHO region Country/area	Year	Contributions reported by donors			
		Global Fund <sup>1</sup>	PMI/USAID <sup>2</sup>	World Bank <sup>3</sup>	UK <sup>4</sup>
<b>AFRICAN</b>					
Sao Tome and Principe	2017	3 030 269	0	0	0
	2018	0	0	0	0
	2019	-509 553	0	0	0
Senegal	2017	6 045 167	26 009 571	0	0
	2018	12 617 208	24 418 477	0	0
	2019	11 572 039	24 000 000	0	0
Sierra Leone	2017	1 548 151	15 605 743	0	1 316 498
	2018	1 467 366	15 261 548	0	927 313
	2019	1 216 641	15 000 000	0	0
South Africa	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
South Sudan <sup>8</sup>	2017	23 629 994	0	0	13 904 529
	2018	11 313 364	0	0	9 794 056
	2019	12 385 841	0	0	12 051 666
Togo	2017	18 522 276	0	2 477 906	0
	2018	6 679 079	0	1 070 093	0
	2019	8 664 142	0	1 051 754	0
Uganda	2017	55 050 846	34 332 634	0	7 595 938
	2018	65 879 046	33 575 405	0	5 350 418
	2019	39 302 893	33 000 000	0	11 811 455
United Republic of Tanzania <sup>9</sup>	2017	75 098 408	45 776 845	0	0
	2018	29 252 693	44 767 207	0	0
	2019	54 867 790	44 000 000	0	92 923
Mainland	2017	69 674 305	0	0	0
	2018	28 751 369	0	0	0
	2019	0	0	0	0
Zanzibar	2017	2 509 129	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Zambia	2017	41 082 748	31 211 486	643 939	0
	2018	22 492 101	30 523 096	870 986	0
	2019	23 722 752	30 000 000	856 060	188 909
Zimbabwe	2017	17 808 245	15 605 743	0	0
	2018	13 178 560	15 261 548	0	0
	2019	17 303 041	15 000 000	0	0
<b>AMERICAS</b>					
Belize	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Bolivia (Plurinational State of)	2017	2 854 289	0	0	0
	2018	3 406 162	0	0	0
	2019	822 768	0	0	0
Brazil	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Colombia	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0

Contributions reported by countries

Government (NMP)	Global Fund	World Bank	PMI/USAID	Other bilaterals	WHO	UNICEF	Other contributions <sup>7</sup>
2 044 439	3 296 207	0	0	0	89 244	0	0
0 <sup>6</sup>							
117 201	517 594	0	0	3 322 449	126 121	52 141	0
4 931 741	3 039 725	0	24 000 000	0	0	0	4 500 000
4 931 741	11 602 821	0	24 000 000	11 602 821	0	0	0
9 420 000	9 005 006	0	24 000 000	0	0	0	14 567 962
821 674 <sup>6</sup>	19 300 000				72 812	3 464 362	
65 189 <sup>5</sup>	8 728 599		15 000 000		70 000	148 214	2 742
128 621	7 522 931		15 000 000		70 000	2 059	4 779
10 656 029	27 226 495	0	0	0	20 000	0	0
16 954 533	4 197 290	0	0	0	50 000	0	
19 251 230	6 591 498	0	0	0	45 000	0	1 132 611
2 603 242 <sup>5</sup>	16 478 112	0	6 000 000	6 654 000	200 000		5 249 000
2 704 995 <sup>6</sup>							
1 069 896	17 047 017	3 124 679	0	3 755 637	0		
1 847 898	24 435 381	1 014 708	0	0	7 765	556 712	5 238 461
64 103	23 830 061	440 567	0	0	4 715	553 567	0
1 889 574 <sup>6</sup>							
7 280 412	150 649 446	0	34 000 000	8 974 881		743 791	4 335 860
7 243 128	47 530 743	0	33 000 000	14 073 138		743 791	0
7 283 521	58 333 000		33 000 000	14 389 262		1 254 438	705 940
6 510 796 <sup>6</sup>							
6 682 225 <sup>6</sup>							
6 682 225 <sup>6</sup>							
70 274 555	70 274 555				42 000		
145 258 808	145 258 808		713 228				12 168
4 898 342	25 110 093	0	8 774 918	0	57 875	0	
8 894	2 960 586	0	978 962		10 000		
79 708	1 508 555	0	15 391 465	0	14 574	0	0
100 434	2 035 288	0	1 096 204	10 000	0	0	0
27 928 587	45 468 736		25 000 000		200 000		
18 159 340	24 605 077		3 000 000		200 000		3 692 991
15 340 495	17 019 922		30 000 000		300 000		5 330 000
782 250	17 407 287		15 120 000				224 970
2 786 540	16 973 379	0	11 000 000	0	118 000	0	0
3 765 250	25 931 599		11 208 498		140 000		
250 000	0	0	9 778	0	0	0	0
252 000	11 122	0	3 234	0	5 609	0	0
252 000	0	0	11 058	0	0	0	0
451 993		0	0	0		0	0
416 666							
292 852	1 191 940	0	0	0	27 891	0	0
54 904 745 <sup>5</sup>	0	0		0		0	
23 923 126 <sup>5</sup>	0	0	82 861	0		0	
53 733 857 <sup>5</sup>	0	0	154 641				
10 897 170	0	0	2 872	0	0	0	0
3 237 708	0	0	70 647				
5 999 473	0	0	269 661	0		0	



## ANNEX 3 – C. FUNDING FOR MALARIA CONTROL, 2017–2019

WHO region Country/area	Year	Contributions reported by donors			
		Global Fund <sup>1</sup>	PMI/USAID <sup>2</sup>	World Bank <sup>3</sup>	UK <sup>4</sup>
<b>AMERICAS</b>					
Costa Rica	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Dominican Republic	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Ecuador	2017	-608 606	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
El Salvador	2017	0	0	0	0
	2018	647 719	0	0	0
	2019	217 471	0	0	0
French Guiana	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Guatemala	2017	2 336 448	0	0	0
	2018	2 228 927	0	0	0
	2019	619 705	0	0	0
Guyana	2017	774 658	0	0	0
	2018	59 439	0	0	0
	2019	75 693	0	0	0
Haiti	2017	10 853 040	0	0	0
	2018	5 576 626	0	0	0
	2019	6 038 170	0	0	0
Honduras	2017	1 252 813	0	0	0
	2018	1 134 584	0	0	0
	2019	1 544 876	0	0	0
Mexico	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Nicaragua	2017	2 534 883	0	0	0
	2018	2 329 152	0	0	0
	2019	2 974 752	0	0	0
Panama	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Peru	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Suriname	2017	1 189 182	0	0	0
	2018	834 200	0	0	0
	2019	655 335	0	0	0
Venezuela (Bolivarian Republic of) <sup>10</sup>	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0

Contributions reported by countries

Government (NMP)	Global Fund	World Bank	PMI/USAID	Other bilaterals	WHO	UNICEF	Other contributions <sup>7</sup>
4 980 000 <sup>5</sup>	0	0	0	0	9 770	0	0
5 000 000 <sup>5</sup>	0	0	0	0	12 155	0	0
5 000 000 <sup>5</sup>	0	0	7 991	0	22 842	0	0
1 149 368	125 543	0	0	0	824	0	27 987
367 647	9 949 957	0	0	0	143 176	0	48 938
2 560 753	0	0	313 661	0	322 922	0	98 488
5 835 716 <sup>5</sup>	0	0	0	0	69 039	0	0
6 898 763 <sup>5</sup>	0	0	0	0	85 733	0	0
2 675 521 <sup>5</sup>	0	0	71 420	0	76 400	0	0
2 662 869	538 732	0	0	0	73 758	0	0
3 950 441	707 436	0	0	0	15 156	0	0
0 <sup>6</sup>			34 787		3 773		
0 <sup>6</sup>	0	0	0	0	0	0	0
0 <sup>6</sup>							
3 374 612	2 231 020		75 981				
3 492 749	1 724 076	0	138 643	0		0	580 000
1 277 993	520 837		76 014		110 535		
1 473 101	1 009 615	0	8 015	0	9 793	0	0
1 503 535	340 471	0	211 698	0	0	0	0
732 166	299 843	0	1 000 000	0	140 000	0	0
388 104 <sup>6</sup>	12 540 295	0	17 956	500 000	227 455		196 777
408 174 <sup>5</sup>	7 384 832	0	0	0	275 872		514 271
2 284 758 <sup>5</sup>	6 006 513	0	10 445	0	266 004		203 638
543 312	2 594 856	0	54 475	0	0	0	554 378
543 312	1 929 881	0	46 855	0	36 961	0	714 145
543 312	1 511 759		67 612	595 460	2 613		621 496
40 661 276	0	0	0	0		0	0
37 544 836	0	0	0	0		0	0
37 024 233			41 177		59 429		
3 984 944	1 826 934		23 971		98 131		
3 263 970	1 986 357		13 254		83 000		401 133
6 154 533	2 313 411		100	400 000	13 408		15 020
3 671 002			49 705		100 000		181 109
8 000 000 <sup>5</sup>	0	0	85 165	0	18 823	0	147 827
6 383 374	475 156		32 085	668 596	62 342		
1 704 408 <sup>5</sup>			39 886		128 851		
2 381 660 <sup>5</sup>			90 000				
3 711 574 <sup>5</sup>	0	0	193 079	0		0	0
806 069	1 041 205	0	52 213	0	12 920	0	0
1 034 627	922 115	0	22 037	0	8 861	0	49 344
1 286 407	695 291		46 808		5 000		30 000
29 452 393 982 <sup>5</sup>			0		85 193		
573 136 589			0		435 366		
0 <sup>6</sup>					147 419		

## ANNEX 3 – C. FUNDING FOR MALARIA CONTROL, 2017–2019

WHO region Country/area	Year	Contributions reported by donors			
		Global Fund <sup>1</sup>	PMI/USAID <sup>2</sup>	World Bank <sup>3</sup>	UK <sup>4</sup>
<b>EASTERN MEDITERRANEAN</b>					
Afghanistan	2017	7 166 347	0	0	0
	2018	9 723 132	0	0	0
	2019	10 199 127	0	0	0
Djibouti	2017	2 662 775	0	244 338	0
	2018	663 592	0	74 153	0
	2019	1 055 614	0	72 882	0
Iran (Islamic Republic of)	2017	1 132 770	0	0	0
	2018	0	0	0	0
	2019	-105 258	0	0	0
Pakistan	2017	16 898 605	0	0	0
	2018	13 827 697	0	0	0
	2019	14 883 169	0	0	0
Saudi Arabia	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Somalia	2017	16 612 625	0	0	0
	2018	7 632 763	0	0	0
	2019	4 246 685	0	0	0
Sudan	2017	10 668 769	0	0	0
	2018	35 329 302	0	0	0
	2019	44 291 755	0	0	0
Yemen	2017	3 728 150	0	47 407 385	0
	2018	-7 374	0	17 395 815	0
	2019	-56 405	0	17 097 691	0
<b>SOUTH-EAST ASIA</b>					
Bangladesh	2017	13 182 596	0	0	0
	2018	7 061 234	0	0	0
	2019	5 406 054	0	0	0
Bhutan	2017	582 622	0	0	0
	2018	332 675	0	0	0
	2019	383 556	0	0	0
Democratic People's Republic of Korea	2017	1 549 812	0	0	0
	2018	2 354 899	0	0	0
	2019	0	0	0	0
India	2017	68 981 923	0	0	0
	2018	275 345	0	0	0
	2019	22 045 007	0	0	0
Indonesia	2017	23 964 363	0	0	0
	2018	10 161 943	0	0	0
	2019	17 489 764	0	0	0
Myanmar	2017	41 491 550	10 403 829	0	4 075 391
	2018	17 304 512	10 174 365	0	2 870 619
	2019	29 430 941	10 000 000	0	536 279
Nepal	2017	5 255 284	0	0	0
	2018	1 433 137	0	0	0
	2019	1 526 228	0	0	0
Thailand	2017	11 147 475	3 000 000	0	0
	2018	6 146 057	3 000 000	0	0
	2019	11 523 833	3 000 000	0	0

Contributions reported by countries

Government (NMP)	Global Fund	World Bank	PMI/USAID	Other bilaterals	WHO	UNICEF	Other contributions <sup>7</sup>
937 596 <sup>6</sup>	1 053 356				85 814		
203 487 <sup>6</sup>	10 556 626				26 571		
	7 759 216				80 885		
3 222 506 <sup>5</sup>		0		0	51 000	0	
3 352 640 <sup>6</sup>	871 414			0	30 000	0	
1 457 180 <sup>5</sup>	171 627				406 776		
2 700 000					48 000		
3 300 000	0	0	0		38 286		
2 930 000	0	0	0	0	38 000	0	0
18 664 597 <sup>6</sup>	22 635 097				130 000		
3 774 306	9 615 605				196 378		
2 443 594	14 600 000				296 000		
30 000 000	0	0	0	0	100 000	0	0
30 000 000	0	0	0	0	10 000	0	0
30 000 000	0	0	0	0	0	0	0
85 350	20 986 170	0	0	0	147 000		0
90 726	5 534 919	0	0	0	56 000		0
120 100	9 474 797	0	0	0	73 840	0	0
19 087 941	31 496 505	0	0	0	3 084	0	0
16 726 945	21 485 294	0	0	0	60 000	203 000	9 619
0	7 933 620				2 080 000	473 627	
0 <sup>5</sup>	1 890 037				1 427 948		
	6 123 238						
1 493 690	8 821 888	0	0	0	210 000	0	0
2 496 429	6 835 307	0	0	0	250 000	0	0
2 634 763	7 082 673	0	0	0	100 000	0	0
179 470	586 015	0	0	0	35 212	0	121 212
176 791	577 403	0	0	0	34 687	0	0
251 860	418 069	0	0	0	40 391	0	121 212
2 151 000	3 426 508	0	0	0	35 000	0	0
2 181 000	3 219 957	0	0	0		0	
2 211 100	0	0	0	0	700 000	0	0
145 564 257	94 474 099	0	0	0		0	
46 783 323	34 958 663	0	0	0		0	
107 930 657	31 242 857	0	0	0		0	
17 686 075 <sup>5</sup>	30 336 061				147 033	1 505 774	
21 683 909 <sup>5</sup>	12 272 515				260 738	933 224	
24 594 057 <sup>5</sup>	25 652 636				100 000	782 076	
6 780 092 <sup>5</sup>	53 056 520	0	10 000 000	6 532 464	25 000	0	3 462 068
6 780 092 <sup>5</sup>	29 581 578		9 000 000	6 607 886	25 000		
11 123 879 <sup>5</sup>	40 110 516		10 000 000	610 000	50 000		
263 262	102 424				24 509		
613 873	1 107 196	0	120 482	0	31 214	0	0
613 873	2 727 909	0	621 652	0	40 000	0	0
7 664 899	15 622 625	0			188 686		49 859
7 131 736	8 337 877	0	1 308 800	0	78 056	0	93 546
5 695 904	8 872 808		1 047 408		70 000		37 710

## ANNEX 3 – C. FUNDING FOR MALARIA CONTROL, 2017–2019

WHO region Country/area	Year	Contributions reported by donors			
		Global Fund <sup>1</sup>	PMI/USAID <sup>2</sup>	World Bank <sup>3</sup>	UK <sup>4</sup>
<b>SOUTH-EAST ASIA</b>					
Timor-Leste	2017	2 735 744	0	0	0
	2018	2 469 564	0	0	0
	2019	2 306 893	0	0	0
<b>WESTERN PACIFIC</b>					
Cambodia	2017	14 619 179	10 403 829	0	0
	2018	10 561 499	10 174 365	0	0
	2019	18 108 881	10 000 000	0	0
China	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Lao People's Democratic Republic	2017	3 731 157	0	0	0
	2018	3 969 853	0	0	0
	2019	6 152 594	0	0	0
Malaysia	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Papua New Guinea	2017	10 747 518	0	0	0
	2018	7 403 211	0	0	0
	2019	10 203 124	0	0	0
Philippines	2017	7 470 423	0	0	0
	2018	3 250 897	0	0	0
	2019	3 062 223	0	0	0
Republic of Korea	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Solomon Islands	2017	1 043 802	0	0	0
	2018	1 759 795	0	0	0
	2019	1 959 252	0	0	0
Vanuatu	2017	0	0	0	0
	2018	0	0	0	0
	2019	0	0	0	0
Viet Nam	2017	16 078 339	0	0	0
	2018	9 458 697	0	0	0
	2019	16 462 619	0	0	0

Global Fund: Global Fund to Fight AIDS, Tuberculosis and Malaria; NGO: nongovernmental organization; NMP: national malaria programme; PMI: United States President's Malaria Initiative; UK: United Kingdom of Great Britain and Northern Ireland government; UNICEF: United Nations Children's Fund; USAID: United States Agency for International Development; WHO: World Health Organization.

"–" refers to data not available.

<sup>1</sup> Source: Global Fund to Fight AIDS, Tuberculosis and Malaria.

<sup>2</sup> Source: [www.foreignassistance.gov](http://www.foreignassistance.gov).

<sup>3</sup> Source: Organisation for Economic Co-operation and Development (OECD) creditor reporting system (CRS) database.

<sup>4</sup> Source: Final UK aid spend.

<sup>5</sup> Budget not expenditure.

Contributions reported by countries

Government (NMP)	Global Fund	World Bank	PMI/USAID	Other bilaterals	WHO	UNICEF	Other contributions <sup>7</sup>
1 115 484	4 039 622	0	0	0	42 456	0	20 000
1 121 287	1 573 936	0	0	0	26 600	0	5 000
441 287	2 281 466				40 000		256 000
663 526	8 045 144	0	6 000 000	0	579 738	0	
83 636	3 181 783	0	10 000 000	0	628 297	0	
65 788	4 388 138	0	10 000 000	0		0	0
19 448 382 <sup>6</sup>							
19 944 390 <sup>6</sup>							
19 602 589 <sup>6</sup>							
1 008 060	1 728 818	0	604 000	0	256 734	0	1 066 089
1 914 750	3 725 427	0	500 000	0	288 108	0	1 783 267
928 955	5 327 000	0	686 183	0	1 039 774	0	1 301 618
48 365 863	0	0	0	0	0	0	0
49 561 180	0	0	0	0	0	0	0
48 817 455	0	0	0	0	0	0	0
753 771	10 330 449	0	0	0	95 000	0	911 770
108 100	7 407 034	0	0	0	86 500	0	1 083 168
48 600	8 831 155			1 474 700	95 000		
7 012 009	6 471 549	0	0	0	0	0	0
3 548 266	4 190 984	0	0	0	0	0	0
2 453 765	3 412 622	0	0	0	0	0	0
475 173	0	0	0	0	0	0	0
433 726	0	0	0	0	0	0	0
719 992	0	0	0	0	0	0	0
858 256	977 025	0	0	0	736 892	0	0
979 891	1 494 080				79 770		
299 919	717 728	0	0	455 000	37 607	0	0
139 254	285 333	0	0	206 575	21 918	0	0
128 194	131 786	0	0	92 363	9 367	0	0
181 350	182 877	0	0	0	178 245	0	0
3 022 523	9 324 657	0	0	0	200 000	0	500 000
1 813 863	7 901 624	0	0	0	105 045	0	315 396
1 620 317	10 221 830	0	0	0	333 000	0	385 000

Data as of 17 November 2020

<sup>6</sup> WHO NMP funding estimates.

<sup>7</sup> Other contributions as reported by countries: NGOs, foundations, etc.

<sup>8</sup> South Sudan became an independent state on 9 July 2011 and a Member State of WHO on 27 September 2011. South Sudan and Sudan have distinct epidemiological profiles comprising high-transmission and low-transmission areas, respectively. For this reason, data up to June 2011 from the Sudanese high-transmission areas (10 southern states which correspond to contemporary South Sudan) and low-transmission areas (15 northern states which correspond to contemporary Sudan) are reported separately.

<sup>9</sup> Where national totals for the United Republic of Tanzania are unavailable, refer to the sum of Mainland and Zanzibar.

Note: Negative disbursements reflect recovery of funds on behalf of the financing organization.

Note: All contributions reported by donors are displayed in US 2019 constant dollars.

**ANNEX 3 – D. COMMODITIES DISTRIBUTION AND COVERAGE, 2017-2019**

WHO region Country/area	Year	No. of LLINs sold or delivered	Modelled percentage of population with access to an ITN	No. of people protected by IRS	No. of RDTs distributed	Any first-line treatment courses delivered (including ACT)	No. of malaria cases treated with any first-line treatment courses (including ACT)	ACT treatment courses delivered	No. of malaria cases treated with ACT
<b>AFRICAN</b>									
Algeria	2017	0	-	-	36	453	-	-	-
	2018	0	-	-	0	1 242	-	-	-
	2019	-	-	-	0	-	1 014	-	-
Angola	2017	2 924 769	50.8	-	397 882	3 090 761	-	3 090 761	-
	2018	3 863 521	52.4	-	2 000 350	1 950 000	-	1 950 000	-
	2019	1 545 055	27.5	-	-	-	-	-	5 575 259
Benin	2017	6 771 009	54	853 221	2 171 867	1 530 617	-	1 530 617	-
	2018	504 501	59.3	1 321 758	2 016 745	1 815 236	-	1 815 236	-
	2019	505 670	31.8	1 077 411	3 984 677	4 455 581	2 353 657	4 455 581	2 353 657
Botswana	2017	3 000	-	139 244	2 645	4 429	-	4 429	-
	2018	-	-	83 488	3 141	1 954	-	1 954	-
	2019	0	-	154 663	2 526	3 198	272	3 198	-
Burkina Faso	2017	986 164	67.3	-	12 853 861	10 457 752	-	10 457 752	-
	2018	1 946 047	49	766 374	13 026 870	11 968 368	-	11 968 368	-
	2019	-	66.5	587 248	-	-	-	-	11 223 002
Burundi	2017	6 717 994	59.4	848 441	10 046 047	7 978 264	-	7 613 646	-
	2018	986 025	71.2	1 754 679	7 012 203	5 149 436	-	5 032 209	-
	2019	7 528 556	51.2	725 449	-	9 338 611	-	9 271 032	-
Cabo Verde	2017	80	-	495 313	16 573	420	-	420	-
	2018	21	-	-	9 588	21	-	21	-
	2019	-	-	302 520	0	40	40	40	0
Cameroon	2017	362 629	71	-	1 589 218	879 039	-	785 765	-
	2018	573 843	62.9	-	1 739 286	1 064 668	-	918 505	-
	2019	8 860 653	69	-	2 082 527	-	1 834 114	-	1 157 011
Central African Republic	2017	857 198	62.1	-	806 218	947 205	-	947 205	-
	2018	753 889	75.5	-	1 189 881	1 773 072	-	1 773 072	-
	2019	103 848	74.3	-	2 764 293	5 753 501	2 654 215	5 640 687	2 602 171
Chad	2017	6 886 534	51.7	-	1 287 405	1 486 086	-	1 486 086	-
	2018	461 667	48.3	-	-	-	-	-	-
	2019	613 700	19.1	0	1 788 730	-	1 665 212	-	1 595 351
Comoros	2017	34 590	81.6	-	21 988	2 794	-	2 794	-
	2018	31 012	67.2	-	-	-	-	-	-
	2019	-	48.1	-	-	-	-	-	-
Congo	2017	2 223	32.1	-	0	0	-	0	-
	2018	4 641	30.2	-	0	0	-	0	-
	2019	2 648 456	72.5	-	0	200 000	427 959	200 000	233 389
Côte d'Ivoire	2017	13 216 468	73	-	6 986 825	5 373 545	-	5 373 545	-
	2018	16 703 932	74.3	-	6 069 250	6 799 565	-	6 799 565	-
	2019	1 410 391	60.5	-	6 456 625	4 657 570	5 200 350	4 657 570	5 200 350
Democratic Republic of the Congo	2017	8 412 959	66.2	232 181	18 994 861	17 250 728	-	17 250 728	-
	2018	16 919 441	58.8	111 735	18 549 327	16 917 207	-	16 917 207	-
	2019	20 710 146	64.8	-	26 963 687	18 853 209	18 853 210	18 853 209	18 853 209
Equatorial Guinea	2017	42 317	40.8	64 617	60 798	15 341	-	15 341	-
	2018	120 376	45.4	74 416	78 695	15 633	-	15 633	-
	2019	14 843	44.8	61 561	54 340	15 769	-	15 769	-
Eritrea	2017	1 724 972	54.1	375 696	481 600	296 399	-	296 399	-
	2018	60 083	61.7	376 143	400 900	301 525	-	301 525	-
	2019	124 225	54.8	437 194	388 395	207 150	-	207 150	-
Eswatini	2017	0	-	21 316	59 760	900	-	861	-
	2018	0	-	39 144	61 974	631	-	579	-
	2019	-	-	15 055	72 369	-	586	-	580
Ethiopia	2017	5 854 497	25.4	17 860 356	6 400 000	8 470 000	-	7 300 000	-
	2018	11 466 972	20.2	9 153 163	4 053 200	3 773 179	-	3 036 690	-
	2019	15 754 582	26.4	7 441 150	8 190 815	11 931 656	1 015 792	5 070 567	836 293

## ANNEX 3 – D. COMMODITIES DISTRIBUTION AND COVERAGE, 2017-2019

WHO region Country/area	Year	No. of LLINs sold or delivered	Modelled percentage of population with access to an ITN	No. of people protected by IRS	No. of RDTs distributed	Any first-line treatment courses delivered (including ACT)	No. of malaria cases treated with any first-line treatment courses (including ACT)	ACT treatment courses delivered	No. of malaria cases treated with ACT
<b>AFRICAN</b>									
Gabon	2017	-	17.7	-	0	0	-	0	-
	2018	4 582	16.6	-	71 787	-	-	208 953	-
	2019	-	14.9	-	-	-	-	0	117 126
Gambia	2017	1 051 391	58.2	396 546	767 984	174 556	-	174 166	-
	2018	115 801	57	426 788	678 621	113 563	-	113 563	-
	2019	1 115 780	45.1	507 872	505 895	-	53 386	-	53 385
Ghana	2017	3 059 363	71.1	1 868 861	7 051 875	4 522 410	-	4 522 410	-
	2018	16 839 135	84.5	1 855 326	13 119 275	5 253 298	-	5 253 298	-
	2019	2 924 717	78.6	1 986 408	12 866 700	4 208 875	6 164 160	4 208 875	6 164 159
Guinea	2017	523 328	50.7	-	2 920 298	2 673 947	-	2 673 947	-
	2018	658 976	30.9	-	2 741 607	1 886 685	-	1 886 685	-
	2019	8 964 940	69.3	-	2 857 744	-	1 646 493	-	-
Guinea-Bissau	2017	1 222 428	64	-	303 651	136 507	-	110 508	-
	2018	93 859	62.5	-	320 217	162 773	-	147 927	-
	2019	102 586	37.5	-	325 690	155 848	155 848	140 478	140 478
Kenya	2017	15 621 773	69.8	906 388	11 337 850	10 696 827	-	10 696 827	-
	2018	2 673 730	70	1 833 860	-	-	-	-	-
	2019	1 797 075	59.8	2 011 860	4 179 875	8 285 622	5 259 988	7 247 430	5 004 487
Liberia	2017	157 954	26.1	-	-	-	-	-	-
	2018	2 500 796	59.4	-	-	994 008	-	994 008	-
	2019	197 736	54.7	-	536 915	-	1 004 895	2 108 721	732 322
Madagascar	2017	764 022	39.7	2 008 963	2 465 600	1 620 050	-	1 620 050	-
	2018	13 520 356	61.2	-	4 731 125	2 165 450	-	2 165 450	-
	2019	1 078 541	67.8	1 640 183	2 899 007	975 587	-	975 587	-
Malawi	2017	994 136	62	-	15 060 625	10 177 530	-	10 177 530	-
	2018	11 805 392	72.6	-	13 003 518	8 948 286	-	9 186 040	-
	2019	1 064 495	71.3	1 456 138	-	-	4 108 225	-	112 411
Mali	2017	4 148 911	64.5	823 201	4 164 041	3 746 616	-	3 746 616	-
	2018	4 993 868	67.3	665 581	6 105 500	3 558 964	-	3 558 964	-
	2019	4 005 010	73.7	690 793	3 656 317	2 846 438	2 846 438	2 826 112	2 826 112
Mauritania	2017	921 245	46.9	-	234 520	101 450	-	-	-
	2018	479 637	44.6	-	117 000	25 890	-	25 890	-
	2019	-	11.8	-	0	-	-	-	-
Mayotte	2017	-	-	-	-	-	-	-	-
	2018	-	-	-	-	44	-	44	-
	2019	-	-	-	-	-	-	-	-
Mozambique	2017	15 482 093	72.1	5 349 948	19 662 975	15 996 892	-	15 996 892	-
	2018	1 337 905	64.9	4 211 138	21 180 223	16 293 318	-	16 293 318	-
	2019	6 614 068	53.2	6 303 792	21 365 400	16 867 851	10 742 632	16 867 851	10 742 632
Namibia	2017	0	-	753 281	914 175	79 316	-	79 316	-
	2018	35 000	-	549 243	49 852	35 355	-	1 721	-
	2019	-	-	149 306	247 425	3 404	3 404	3 404	0
Niger	2017	981 423	69.7	0	3 909 600	2 697 115	-	2 161 440	-
	2018	4 024 529	74.4	-	5 149 981	3 536 000	-	3 536 000	-
	2019	-	76.1	-	5 831 287	3 211 243	3 015 081	3 211 243	-
Nigeria	2017	21 978 907	49.4	-	9 701 771	7 752 372	-	7 752 372	-
	2018	27 675 958	50.1	-	18 662 105	32 707 785	-	32 707 785	-
	2019	30 417 404	48.1	-	26 312 300	38 240 771	21 252 650	38 240 771	21 252 650
Rwanda	2017	2 816 586	75.4	1 753 230	4 960 020	6 300 445	-	6 265 890	-
	2018	974 847	53.5	1 621 955	5 364 990	5 233 680	-	5 214 330	-
	2019	536 637	36.4	4 532 103	4 904 370	4 231 880	3 566 544	4 215 120	3 545 251
Sao Tome and Principe	2017	15 151	-	138 000	96 826	2 410	-	2 410	-
	2018	142 894	-	-	-	-	-	-	-
	2019	16 260	-	53 401	221 450	2 457	2 457	2 457	2 457



## ANNEX 3 – D. COMMODITIES DISTRIBUTION AND COVERAGE, 2017-2019

WHO region Country/area	Year	No. of LLINs sold or delivered	Modelled percentage of population with access to an ITN	No. of people protected by IRS	No. of RDTs distributed	Any first-line treatment courses delivered (including ACT)	No. of malaria cases treated with any first-line treatment courses (including ACT)	ACT treatment courses delivered	No. of malaria cases treated with ACT
<b>AFRICAN</b>									
Senegal	2017	448 305	73.6	619 578	2 391 311	958 473	-	958 473	-
	2018	617 470	47.7	0	2 646 144	1 606 813	-	1 490 147	-
	2019	9 373 577	72.1	51 652	2 552 381	-	354 432	-	339 598
Sierra Leone	2017	4 654 654	66.3	-	2 611 550	2 504 960	-	2 504 960	-
	2018	502 834	69	-	4 316 420	3 415 480	-	3 415 480	-
	2019	492 622	49.5	-	3 930 606	4 751 000	2 813 086	4 751 000	2 404 286
South Africa	2017	0	-	1 550 235	865 050	72 439	-	72 439	-
	2018	0	-	1 600 747	887 300	51 142	-	51 142	-
	2019	-	-	1 477 420	879 625	10 592	-	10 592	13 833
South Sudan <sup>1</sup>	2017	1 902 020	59.7	153 285	1 945 875	12 188 601	-	12 188 601	-
	2018	963 092	42.4	-	-	2 680 776	-	2 680 776	-
	2019	713 717	34.2	344 242	-	-	-	-	-
Togo	2017	4 706 417	77.9	-	1 613 393	1 355 640	-	1 196 518	-
	2018	224 265	79.5	-	2 485 086	1 988 845	-	2 055 831	-
	2019	407 911	66.7	-	2 957 298	2 284 746	2 284 746	1 499 012	2 266 412
Uganda	2017	23 797 483	83.1	3 223 800	24 620 100	27 396 300	-	27 396 300	-
	2018	11 220 492	79.9	4 436 156	28 200 125	25 606 514	-	25 606 514	-
	2019	1 855 163	60.6	4 478 754	20 979 775	17 706 390	-	17 706 390	-
United Republic of Tanzania <sup>2</sup>	2017	5 335 910	-	2 759 641	35 109 007	20 903 686	-	20 903 686	-
	2018	6 378 169	-	2 842 635	30 263 725	16 425 610	-	16 425 210	-
	2019	6 968 606	-	2 989 048	26 058 455	8 487 473	6 963	8 485 301	6 385 075
Mainland	2017	5 335 910	59.6	2 568 522	34 649 050	20 895 180	-	20 895 180	-
	2018	6 200 375	59.4	2 507 920	29 906 950	16 420 560	-	16 420 560	-
	2019	6 745 132	-	2 507 920	25 699 255	8 479 635	-	8 479 635	6 378 890
Zanzibar	2017	0	-	191 119	459 957	8 506	-	8 506	-
	2018	177 794	-	334 715	356 775	5 050	-	4 650	-
	2019	223 474	-	481 128	359 200	7 838	6 963	5 666	6 185
Zambia	2017	10 759 947	70.3	7 717 767	18 884 600	17 460 232	-	17 460 232	-
	2018	689 288	64.1	6 436 719	17 868 550	27 071 994	-	27 071 994	-
	2019	1 024 635	47.3	11 767 404	17 737 525	19 134 471	19 134 471	19 134 471	-
Zimbabwe	2017	513 300	46.4	3 673 311	875 713	549 083	-	553 953	-
	2018	1 015 246	36.3	3 020 032	1 484 134	607 379	-	615 359	-
	2019	2 160 175	36.6	3 164 344	1 445 007	-	304 309	-	304 309
<b>AMERICAS</b>									
Argentina	2017	0	-	4 208	0	39	-	9	-
	2018	0	-	155	0	213	-	92	-
	2019	-	-	-	-	-	-	-	-
Belize	2017	0	-	37 466	0	9	-	1	-
	2018	2 619	-	36 688	0	7	-	0	-
	2019	0	-	43 497	0	-	2	0	0
Bolivia (Plurinational State of)	2017	23 500	-	20 000	3 500	0	-	0	-
	2018	23 500	-	2 000	-	-	-	-	-
	2019	27 000	-	29 228	36 800	8 600	-	-	9 357
Brazil	2017	0	-	83 990	72 200	651 274	-	69 960	-
	2018	300 000	-	99 321	114 775	634 935	-	79 200	-
	2019	0	-	84 126	102 275	491 126	491 126	74 360	74 360
Colombia	2017	295 250	-	153 690	265 250	95 570	-	56 030	-
	2018	0	-	60 000	13 252	46 217	-	26 507	-
	2019	78 320	-	143 320	25 000	97 324	-	59 100	-
Costa Rica	2017	104	-	8 479	0	25	-	7	-
	2018	3 100	-	4 095	700	108	-	5	-
	2019	-	-	-	-	-	-	-	-

## ANNEX 3 – D. COMMODITIES DISTRIBUTION AND COVERAGE, 2017-2019

WHO region Country/area	Year	No. of LLINs sold or delivered	Modelled percentage of population with access to an ITN	No. of people protected by IRS	No. of RDTs distributed	Any first-line treatment courses delivered (including ACT)	No. of malaria cases treated with any first-line treatment courses (including ACT)	ACT treatment courses delivered	No. of malaria cases treated with ACT
<b>AMERICAN</b>									
Dominican Republic	2017	0	-	30 361	48 850	398	-	-	-
	2018	5 052	-	36 891	42 425	484	-	9	-
	2019	-	-	33 226	55 000	-	1 314	-	7
Ecuador	2017	72 015	-	667 111	-	1 380	-	371	-
	2018	50 000	-	775 884	51 200	1 806	-	191	-
	2019	31 271	-	698 292	73 425	5 030	1 909	2 650	265
El Salvador	2017	2 925	-	19 167	0	4	-	0	-
	2018	4 817	-	32 691	0	2	-	1	-
	2019	1 813	-	23 512	-	-	-	-	-
French Guiana	2017	-	-	-	-	-	-	-	-
	2018	-	-	-	-	-	-	-	-
	2019	-	-	-	-	-	-	-	-
Guatemala	2017	83 258	-	6 245	170 325	9 995	-	0	-
	2018	310 218	-	15 358	75 300	3 246	-	-	-
	2019	128 982	-	4 091	61 275	-	-	-	2
Guyana	2017	5 534	-	-	-	13 936	-	5 141	-
	2018	43 181	-	-	-	11 767	-	3 073	-
	2019	1 759	-	-	37 800	16 913	-	6 622	-
Haiti	2017	709 720	-	-	261 600	18 772	-	-	-
	2018	1 919	-	42 130	207 800	8 083	-	-	-
	2019	19 063	-	-	293 200	22 172	10 687	-	-
Honduras	2017	24 092	-	225 027	29 710	-	-	-	-
	2018	53 944	-	338 730	15 000	14	-	45	-
	2019	32 091	-	-	17 580	14	-	-	2
Mexico	2017	5 695	-	87 772	0	765	-	14	-
	2018	17 891	-	85 812	0	803	-	10	-
	2019	19 001	-	83 581	0	641	-	-	12
Nicaragua	2017	103 676	-	182 602	46 500	49 085	-	50	-
	2018	47 301	-	183 098	117 350	86 195	-	-	-
	2019	228 589	-	139 795	63 500	35 649	13 226	-	-
Panama	2017	-	-	3 921	16 000	689	-	144	-
	2018	0	-	19 500	20 000	715	-	3	-
	2019	3 952	-	12 806	30 000	-	-	-	3
Paraguay	2017	0	-	631	5 000	2 498	-	408	-
	2018	-	-	-	-	-	-	-	-
	2019	-	-	-	-	-	-	-	-
Peru	2017	-	-	62 804	-	-	-	-	-
	2018	83 220	-	23 420	180 000	65 000	-	14 500	-
	2019	-	-	59 438	204 000	51 289	-	-	4 724
Suriname	2017	6 022	-	-	14 325	-	-	-	-
	2018	15 000	-	-	13 575	-	-	-	-
	2019	6 847	-	-	20 625	-	-	-	-
Venezuela (Bolivarian Republic of)	2017	5 000	-	3 900	-	-	-	-	-
	2018	81 402	-	-	48 117	404 924	-	97 293	-
	2019	256 311	-	-	250 000	398 285	-	-	90 153
<b>EASTERN MEDITERRANEAN</b>									
Afghanistan	2017	2 372 354	-	-	514 875	27 850	-	27 850	-
	2018	649 383	-	-	28 915	-	-	47 665	-
	2019	1 336 070	-	-	714 700	-	169 504	-	-
Djibouti	2017	134 701	20.7	-	63 488	14 212	-	-	-
	2018	109 500	31.5	-	91 416	46 380	-	98 380	-
	2019	218 650	20.9	37 663	335 625	148 890	47 691	148 890	47 691

## ANNEX 3 – D. COMMODITIES DISTRIBUTION AND COVERAGE, 2017-2019

WHO region Country/area	Year	No. of LLINs sold or delivered	Modelled percentage of population with access to an ITN	No. of people protected by IRS	No. of RDTs distributed	Any first-line treatment courses delivered (including ACT)	No. of malaria cases treated with any first-line treatment courses (including ACT)	ACT treatment courses delivered	No. of malaria cases treated with ACT
<b>EASTERN MEDITERRANEAN</b>									
Iran (Islamic Republic of)	2017	4 218	-	126 111	-	-	-	-	-
	2018	4 500	-	117 174	128 650	-	-	-	-
	2019	-	-	64 365	7 737	4 984	159	8 139	159
Pakistan	2017	1 048 037	-	776 650	1 826 221	800 000	-	63 566	-
	2018	2 762 975	-	2 937 767	2 584 675	1 000 000	-	65 000	-
	2019	2 405 841	-	1 657 670	3 895 000	290 170	413 533	57 781	90 178
Saudi Arabia	2017	127 800	-	253 222	-	1 915	-	1 915	-
	2018	127 801	-	242 009	-	1 908	-	1 908	-
	2019	0	-	225 510	13 500 000	25 000	2 152	15 000	1 515
Somalia	2017	2 571 923	21.9	1 267 526	468 750	322 260	-	322 260	-
	2018	357 569	22	2 038 381	755 750	260 580	-	260 580	-
	2019	388 766	16.8	82 720	974 700	174 030	-	174 030	-
Sudan	2017	5 741 449	63	3 683 031	3 498 425	4 507 838	-	4 507 838	-
	2018	3 454 519	61.7	3 830 195	4 117 300	4 195 600	-	4 195 600	-
	2019	8 565 841	58	3 886 652	7 246 975	4 297 167	4 297 167	4 297 167	4 297 167
Yemen	2017	433 266	-	1 338 585	148 935	138 494	-	77 115	-
	2018	1 461 760	-	995 328	571 175	440 265	-	38 420	-
	2019	612 884	-	1 982 284	907 425	458 103	-	42 698	-
<b>SOUTH-EAST ASIA</b>									
Bangladesh	2017	2 242 527	-	-	373 138	29 916	-	24 790	-
	2018	1 559 423	-	72 000	500 440	10 762	-	8 609	-
	2019	727 253	-	98 786	756 573	17 225	17 225	15 099	15 099
Bhutan	2017	137 000	-	71 690	21 650	132	62	132	10
	2018	29 770	-	76 809	12 300	293	54	293	17
	2019	13 906	-	118 730	29 100	42	235	235	11
Democratic People's Republic of Korea	2017	0	-	1 147 548	176 612	17 038	-	0	-
	2018	500 815	-	169 841	657 050	3 698	-	0	-
	2019	30 928	-	0	458 743	4 000	1 869	0	0
India	2017	16 340 000	-	39 341 409	1 064 000	104 110	-	62 650	-
	2018	9 648 400	-	34 290 886	10 500 000	1 400 000	-	1 100 000	-
	2019	22 410 000	-	30 363 425	-	-	338 494	-	156 940
Indonesia	2017	4 376 636	-	3 320	1 783 498	607 965	-	607 965	-
	2018	340 074	-	305 493	255 300	670 603	-	670 603	-
	2019	-	-	164 192	1 980 775	-	234 381	-	234 381
Myanmar	2017	5 835 192	-	-	2 053 525	108 364	-	108 364	-
	2018	775 251	-	14 017	1 761 775	57 144	-	57 144	-
	2019	11 046 312	-	4 361	2 652 010	51 779	53 003	51 779	23 623
Nepal	2017	324 156	-	300 000	100 000	3 070	-	238	-
	2018	319 046	-	230 000	132 065	3 949	-	120	-
	2019	162 409	-	263 000	205 636	13 621	710	3 522	63
Sri Lanka	2017	18 019	-	10 317	27 500	57	-	27	-
	2018	21 759	-	15 707	11 150	48	-	15	-
	2019	29 941	-	3 467	20 035	53	-	24	-
Thailand	2017	358 400	-	207 250	173 425	21 540	-	7 540	-
	2018	131 425	-	165 580	30 550	25 292	-	9 892	-
	2019	80 000	-	489 105	303 613	31 276	3 904	11 976	536
Timor-Leste	2017	334 471	-	102 891	115 115	5 593	30	5 593	30
	2018	35 367	-	154 410	144 061	5 633	8	5 633	8
	2019	97 586	-	175 654	249 750	1 070	9	1 070	9

## ANNEX 3 – D. COMMODITIES DISTRIBUTION AND COVERAGE, 2017–2019

WHO region Country/area	Year	No. of LLINs sold or delivered	Modelled percentage of population with access to an ITN	No. of people protected by IRS	No. of RDTs distributed	Any first-line treatment courses delivered (including ACT)	No. of malaria cases treated with any first-line treatment courses (including ACT)	ACT treatment courses delivered	No. of malaria cases treated with ACT
<b>WESTERN PACIFIC</b>									
Cambodia	2017	1 994 150	-	-	503 250	145 518	-	145 518	-
	2018	1 624 507	-	-	-	-	-	-	-
	2019	-	-	-	923 375	98 965	32 197	98 965	32 197
China	2017	11 349	-	352 731	-	-	-	-	-
	2018	5 987	-	161 224	-	-	-	-	-
	2019	1 807	-	206 599	-	-	2 487	-	2 125
Lao People's Democratic Republic	2017	242 405	-	-	333 675	42 972	-	39 272	-
	2018	50 403	-	2 052	34 387	8 931	-	34 765	-
	2019	1 085 527	-	3 333	1 371 367	21 071	6 551	21 071	6 550
Malaysia	2017	278 104	-	539 029	0	4 114	-	3 443	-
	2018	213 073	-	-	0	4 630	-	3 891	-
	2019	112 054	-	323 208	0	3 933	3 933	3 933	3 923
Papua New Guinea	2017	1 694 315	-	-	1 135 577	832 532	-	832 532	-
	2018	1 480 705	-	-	2 268 750	1 385 940	-	1 385 940	-
	2019	1 476 976	-	-	2 454 525	13 230 420	-	1 610 240	-
Philippines	2017	814 984	-	490 640	145 325	23 400	-	23 400	-
	2018	1 156 837	-	1 015 672	168 300	4 318	-	4 318	-
	2019	695 691	-	731 696	370 700	49 359	4 845	16 857	5 435
Republic of Korea	2017	0	-	-	0	515	-	-	-
	2018	0	-	-	0	576	-	-	-
	2019	-	-	0	20 000	-	196	-	0
Solomon Islands	2017	85 976	-	0	374 850	238 665	-	238 665	-
	2018	150 248	-	-	386 975	233 917	-	233 917	-
	2019	297 010	-	-	484 750	230 880	83 733	230 880	83 364
Vanuatu	2017	91 028	-	0	56 150	27 409	1 075	20 853	-
	2018	27 151	-	0	50 850	0	640	0	-
	2019	80 623	-	0	4 490	7 235	571	579	571
Viet Nam	2017	752 000	-	151 153	921 897	87 225	-	40 000	-
	2018	1 193 024	-	319 866	576 930	45 040	-	40 000	-
	2019	31 740	-	696 751	472 173	31 348	5 892	3 134	3 134

Data as of 17 November 2020

ACT: artemisinin-based combination therapy; IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; LLIN: long-lasting insecticidal net; RDT: rapid diagnostic test; WHO: World Health Organization.

“-” refers to data not available.

<sup>1</sup> In May 2013, South Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [http://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](http://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

<sup>2</sup> Where national data for the United Republic of Tanzania are unavailable, refer to Mainland and Zanzibar.

**ANNEX 3 - Ea. HOUSEHOLD SURVEY RESULTS, 2015-2019, COMPILED BY STATCOMPILER**

WHO region Country/area	Source	% of households					% of population	
		with at least one ITN	with at least one ITN for every two persons who stayed in the household the previous night	with IRS in the past 12 months	with at least one ITN and/or IRS in the past 12 months	with at least one ITN for every two persons and/or IRS in the past 12 months	with access to an ITN	who slept under an ITN last night
<b>AFRICAN</b>								
Angola	2015-16 DHS	30.9	11.3	1.6	31.8	12.5	19.7	17.6
Benin	2017-18 DHS	91.5	60.5	8.7	92.0	63.8	77.2	71.1
Burkina Faso	2017-18 MIS	75.3	32.8	-	-	-	54.5	44.1
Burundi	2016-17 DHS	46.2	17.1	1.0	46.8	17.9	32.3	34.7
Cameroon	2018 DHS	73.4	40.7	-	-	-	58.5	53.9
Ethiopia	2016 DHS	-	-	-	-	-	-	-
Ghana	2016 MIS	73.0	50.9	8.1	74.1	53.6	65.8	41.7
Ghana	2019 MIS	73.7	51.8	5.8	75.0	54.7	66.7	43.2
Guinea	2018 DHS	43.9	16.7	-	-	-	30.7	22.7
Kenya	2015 MIS	62.5	40.0	-	62.5	39.7	52.5	47.6
Liberia	2016 MIS	61.5	25.2	1.2	62.1	25.9	41.5	39.3
Madagascar	2016 MIS	79.5	44.4	6.9	80.9	47.9	62.1	68.2
Malawi	2015-16 DHS	56.9	23.5	4.9	58.6	27.0	38.8	33.9
Malawi	2017 MIS	82.1	41.7	-	-	-	63.1	55.4
Mali	2015 MIS	93.0	39.3	4.0	93.6	41.8	69.5	63.9
Mali	2018 DHS	89.8	54.8	-	-	-	75.2	72.9
Mozambique	2015 AIS	66.0	38.9	11.2	68.7	45.3	53.8	45.4
Mozambique	2018 MIS	82.2	51.2	-	-	-	68.5	68.4
Nigeria	2015 MIS	68.8	34.9	1.3	69.0	35.5	54.7	37.3
Nigeria	2018 DHS	60.6	29.8	-	-	-	47.5	43.2
Rwanda	2014-15 DHS	80.6	42.6	-	80.6	42.5	63.8	61.4
Rwanda	2017 MIS	84.1	55.1	19.6	89.2	66.9	71.9	63.9
Senegal	2015 DHS	76.8	40.5	4.8	77.1	43.0	66.0	51.0
Senegal	2016 DHS	82.4	56.4	5.3	82.9	58.0	75.7	63.1
Senegal	2017 DHS	84.2	50.4	4.2	84.5	52.3	72.8	56.9
Senegal	2018 DHS	76.6	39.0	2.1	76.8	40.1	62.2	51.6
Sierra Leone	2016 MIS	60.3	16.2	1.7	61.1	17.7	37.1	38.6
South Africa	2016 DHS	-	-	-	-	-	-	-
Togo	2017 MIS	85.2	71.4	-	-	-	82.3	62.5
Uganda	2016 DHS	78.4	51.1	-	-	-	64.6	55.0
Uganda	2018-19 MIS	83.0	53.9	10.1	84.2	58.7	71.5	59.2
United Republic of Tanzania	2015-16 DHS	65.6	38.8	5.5	66.2	41.0	55.9	49.0
United Republic of Tanzania	2017 MIS	77.9	45.4	-	-	-	62.5	52.2
Zambia	2018 DHS	78.3	40.9	35.3	83.3	60.4	59.9	46.4
Zimbabwe	2015 DHS	47.9	26.4	21.3	54.9	39.4	37.2	8.5

% of ITNs	% of pregnant women		% of children <5 years				% of children <5 years with fever in the past 2 weeks			
that were used last night	who slept under an ITN	who took 3+ doses of IPTp	who slept under an ITN	with moderate or severe anaemia	with a positive RDT	with a positive microscopy blood smear	for whom advice or treatment was sought	who had blood taken from a finger or heel for testing	who took antimalarial drugs	who took an ACT among those who received any antimalarial
71.0	23.0	20.0	21.7	34.0	13.5	-	51.8	34.3	18.1	76.7
73.4	79.3	13.7	76.3	43.8	36.3	39.1	53.1	17.7	17.5	37.0
76.0	58.2	57.7	54.4	50.1	20.2	16.9	73.5	48.8	51.1	79.4
86.9	43.9	12.9	39.9	36.3	37.9	26.8	69.6	66.4	47.0	11.3
76.2	61.0	31.9	59.8	31.0	24.0		61.0	21.4	32.7	21.2
-	-	-	-	32.0	-	-	35.3		7.7	11.5
47.7	50.0	59.6	52.2	35.2	27.9	20.6	71.9	30.3	50.1	58.8
50.1	48.7	61.0	54.1	27.9	23.0	14.1	69.0	34.1	45.9	84.5
64.0	28.1	35.7	26.6	43.8	-	-	62.3	20.5	24.8	18.2
75.2	57.8	22.9	56.1	16.2	9.1	5.0	72.4	39.2	27.1	91.6
71.2	39.5	23.1	43.7	49.2	44.9	-	78.2	49.8	65.5	81.1
78.7	68.5	10.6	73.4	20.5	5.1	6.9	59.0	15.5	10.1	17.0
73.3	43.9	30.4	42.7	36.1	-	-	67.0	52.0	37.6	91.8
76.8	62.5	41.1	67.5	37.1	36.0	24.3	54.4	37.6	29.4	96.4
90.7	77.9	21.0	71.2	63.0	32.4	35.7	50.0	14.2	28.7	28.9
88.7	83.7	28.3	79.1	56.7	18.9	-	52.8	16.4	18.7	31.0
70.9	52.1	23.3	47.9	36.7	40.2	-	63.2	39.6	38.4	92.6
85.4	76.4	40.6	72.7	55.2	38.9	-	68.6	47.9	32.7	98.6
60.8	49.0	21.4	43.6	43.1	45.1	27.4	66.9	12.6	41.2	37.6
80.6	58.0	16.6	52.2	41.1	36.2	22.6	72.8	13.8	43.5	52.0
77.4	72.9	-	67.7	15.7	7.8	2.2	57.0	36.1	11.4	98.7
71.0	68.5	-	68.0	-	11.8	7.2	55.6	38.1	19.6	98.7
70.0	51.8	11.2	55.4	38.0	0.6	0.3	49.6	9.5	3.4	12.5
68.2	69.0	22.1	66.6	36.7	0.9	0.9	49.9	13.0	1.7	85.0
68.6	61.8	22.0	60.7	41.8	0.9	0.4	51.4	16.1	4.7	65.5
74.5	55.7	22.4	56.4	-	-	-	52.8	13.8	5.1	24.0
89.0	44.0	31.1	44.1	49.2	52.7	40.1	71.7	51.1	57.0	96.0
-	-	-	-	37.0	-	-	-	-	-	-
52.3	69.0	41.7	69.7	47.8	43.9	28.3	55.9	29.3	31.1	76.3
74.0	64.1	17.2	62.0	29.2	30.4	-	81.2	49.0	71.5	87.8
74.3	65.4	41.0	60.3	25.0	18.2	9.8	87.0	50.7	62.5	87.7
69.4	53.9	8.0	54.4	31.2	14.4	5.6	81.2	35.9	51.1	84.9
66.7	51.4	25.8	54.6	30.5	7.3	-	75.4	43.1	36.2	89.4
64.2	48.9	58.7	51.6	29.5	-	-	77.2	63.0	34.9	96.9
18.8	6.1	-	9.0	14.9	-	-	50.7	12.7	1.0	-

**ANNEX 3 – Ea. HOUSEHOLD SURVEY RESULTS, 2015–2019, COMPILED BY STATCOMPILER**

WHO region Country/area	Source	% of households					% of population	
		with at least one ITN	with at least one ITN for every two persons who stayed in the household the previous night	with IRS in the past 12 months	with at least one ITN and/or IRS in the past 12 months	with at least one ITN for every two persons and/or IRS in the past 12 months	with access to an ITN	who slept under an ITN last night
<b>AMERICAS</b>								
Guatemala	2014–15 DHS	-	-	-	-	-	-	-
Haiti	2016–17 DHS	30.7	12.3	2.6	32.2	14.5	19.9	13.0
<b>EASTERN MEDITERRANEAN</b>								
Afghanistan	2015 DHS	26.0	2.9	-	-	-	13.2	3.9
Pakistan	2017–18 DHS	3.6	0.6	5.1	8.4	5.7	2.0	0.2
<b>SOUTH-EAST ASIA</b>								
India	2015–16 DHS	7.6	3.3	-	-	-	5.3	4.2
Myanmar	2015–16 DHS	26.8	14.1	-	-	-	21.2	15.6
Nepal	2016 DHS	-	-	-	-	-	-	-
Timor-Leste	2016 DHS	64.0	32.8	-	-	-	48.3	47.6
<b>WESTERN PACIFIC</b>								
Papua New Guinea	2016–18 DHS	68.5	45.2	-	-	-	57.9	46.0

ACT: artemisinin-based combination therapy; AIS: AIDS indicator survey; DHS: demographic and health survey; IPTp: intermittent preventive treatment in pregnancy; IRS: indoor residual spraying; ITN: insecticide-treated mosquito net; MIS: malaria indicator survey; RDT: rapid diagnostic test; WHO: World Health Organization.

“-” refers to not applicable or data not available.

Sources: Nationally representative household survey data from DHS and MIS, compiled through STATcompiler – <https://www.statcompiler.com/>.

% of ITNs that were used last night	% of pregnant women		% of children <5 years				% of children <5 years with fever in the past 2 weeks			
	who slept under an ITN	who took 3+ doses of IPTp	who slept under an ITN	with moderate or severe anaemia	with a positive RDT	with a positive microscopy blood smear	for whom advice or treatment was sought	who had blood taken from a finger or heel for testing	who took antimalarial drugs	who took an ACT among those who received any antimalarial
-	-	-	-	12.1	-	-	-	-	-	-
62.3	16.0	-	18.2	37.5	-	-	46.8	15.8	1.1	-
21.4	4.1	-	4.6	-	-	-	63.5	7.9	11.8	4.4
11.6	0.4	-	0.4	-	-	-	81.4		9.2	3.3
68.9	4.7	-	5.0	30.8	-	-	81.1	10.8	20.1	8.5
58.3	18.4	-	18.6	26.7	-	-	66.7	3.0	0.8	-
-	-	-		26.5	-	-	-	-	-	-
79.9	60.1	-	55.7	12.6	-	-	57.6	24.5	10.0	11.1
67.9	49.0	23.5	51.5	-	-	-	49.5	24.6	21.3	3.3

Data as of 17 November 2020



### ANNEX 3 – Eb. HOUSEHOLD SURVEY RESULTS, 2015–2019, COMPILED THROUGH WHO CALCULATIONS

WHO region Country/area	Survey	Fever prevalence	Health sector where treatment was sought							Diagnostic testing coverage in each health sector	
		Overall	Public excluding community health workers	Community health workers	Formal medical private excluding pharmacies	Pharmacies or accredited drug stores	Informal private	No treatment seeking	Trained provider	Public excluding community health workers	Community health workers
Angola	2015 DHS	15 (14, 16)	47 (44, 50)	0 (0, 0)	5 (4, 7)	1 (1, 2)	2 (2, 3)	45 (42, 48)	53 (50, 56)	59 (54, 63)	-
Benin	2017 DHS	20 (18, 21)	22 (20, 24)	0 (0, 0)	9 (8, 11)	9 (8, 11)	14 (12, 16)	46 (43, 49)	40 (37, 43)	52 (47, 57)	-
Burkina Faso	2017 MIS	20 (19, 22)	71 (67, 75)	1 (0, 1)	1 (1, 4)	0 (0, 1)	2 (1, 3)	26 (22, 30)	73 (69, 76)	66 (61, 70)	-
Burundi	2016 DHS	40 (38, 41)	54 (51, 56)	3 (2, 4)	10 (8, 12)	5 (4, 5)	1 (0, 1)	30 (28, 32)	69 (67, 71)	87 (86, 89)	95 (89, 98)
Cameroon	2018 DHS	16 (14, 17)	20 (17, 23)	1 (0, 1)	12 (9, 15)	12 (9, 14)	21 (18, 24)	37 (33, 41)	43 (39, 47)	52 (44, 61)	-
Ethiopia	2016 DHS	14 (13, 16)	26 (23, 30)	0 (0, 0)	9 (7, 12)	0 (0, 0)	2 (1, 3)	63 (60, 67)	35 (31, 38)	-	-
Ghana	2019 MIS	30 (27, 33)	34 (30, 38)	0 (0, 1)	20 (17, 24)	14 (10, 18)	3 (1, 5)	30 (26, 35)	67 (63, 71)	78 (72, 83)	-
Guinea	2018 DHS	17 (16, 19)	40 (36, 43)	0 (0, 1)	5 (3, 6)	0 (0, 0)	24 (21, 27)	32 (29, 36)	45 (41, 48)	42 (37, 47)	-
Kenya	2015 MIS	36 (34, 39)	51 (46, 55)	1 (0, 2)	15 (12, 19)	5 (3, 7)	3 (2, 4)	27 (24, 31)	70 (67, 74)	52 (46, 57)	-
Liberia	2016 MIS	39 (36, 43)	46 (41, 52)	0 (0, 0)	13 (10, 17)	14 (11, 18)	8 (6, 11)	22 (18, 26)	71 (67, 76)	77 (72, 82)	-
Madagascar	2016 MIS	16 (15, 18)	36 (31, 41)	7 (5, 10)	10 (8, 14)	1 (1, 2)	7 (5, 10)	40 (36, 44)	53 (49, 58)	31 (25, 39)	37 (22, 54)
Malawi	2017 MIS	40 (38, 43)	38 (34, 43)	3 (2, 5)	6 (4, 8)	2 (1, 4)	7 (5, 10)	46 (41, 51)	48 (43, 52)	76 (70, 82)	-
Mali	2018 DHS	16 (15, 17)	24 (21, 27)	3 (2, 5)	2 (1, 3)	7 (5, 9)	23 (19, 27)	42 (38, 46)	36 (33, 39)	46 (39, 53)	37 (22, 56)
Mozambique	2018 MIS	31 (28, 35)	64 (57, 70)	4 (2, 7)	0 (0, 1)	0 (0, 1)	1 (1, 3)	31 (26, 37)	68 (62, 73)	72 (67, 76)	41 (13, 76)
Nigeria	2018 DHS	24 (23, 25)	27 (25, 29)	1 (1, 1)	38 (36, 40)	5 (4, 6)	4 (3, 5)	26 (25, 28)	70 (68, 72)	35 (32, 38)	9 (4, 18)
Rwanda	2017 MIS	31 (28, 34)	33 (29, 37)	18 (15, 22)	3 (2, 4)	5 (3, 7)	1 (1, 2)	44 (40, 48)	55 (51, 59)	73 (65, 80)	74 (65, 82)
Senegal	2018 DHS	20 (18, 22)	40 (37, 44)	1 (0, 1)	2 (1, 4)	8 (5, 13)	2 (1, 3)	47 (41, 53)	51 (46, 57)	29 (24, 35)	-
Sierra Leone	2016 MIS	27 (25, 29)	63 (60, 66)	0 (0, 0)	4 (3, 6)	4 (3, 5)	2 (1, 3)	28 (25, 31)	70 (67, 74)	74 (71, 77)	-
South Africa	2016 DHS	21 (19, 23)	41 (36, 45)	0 (0, 1)	12 (9, 16)	15 (12, 20)	2 (1, 3)	31 (27, 36)	67 (62, 72)	-	-
Togo	2017 MIS	24 (22, 27)	26 (22, 31)	5 (4, 8)	7 (5, 9)	3 (2, 5)	16 (12, 21)	43 (37, 49)	42 (37, 47)	78 (71, 84)	76 (60, 87)
Uganda	2018 MIS	27 (24, 30)	33 (29, 37)	7 (5, 9)	38 (34, 41)	12 (10, 15)	1 (1, 1)	13 (11, 15)	86 (84, 88)	84 (79, 88)	77 (68, 83)
United Republic of Tanzania	2017 MIS	21 (19, 22)	46 (43, 50)	0 (0, 1)	13 (11, 16)	17 (15, 20)	1 (1, 2)	25 (22, 28)	75 (71, 78)	66 (60, 71)	-
Zambia	2018 DHS	16 (15, 17)	69 (66, 72)	3 (2, 5)	4 (3, 6)	0 (0, 1)	1 (0, 2)	23 (20, 26)	76 (73, 79)	78 (73, 82)	83 (64, 93)
Zimbabwe	2015 DHS	14 (13, 16)	35 (30, 40)	1 (1, 3)	9 (7, 13)	0 (0, 0)	6 (4, 8)	49 (45, 54)	45 (40, 50)	26 (20, 32)	-

Notes:

The analysis is presented as: point estimate (95% confidence interval).

Figures with fewer than 30 children in the denominator were removed.

“-” refers to not applicable or data not available.

Diagnostic testing coverage in each health sector				Antimalarial treatment coverage in each health sector							ACT use among antimalarial treatment in each health sector		
Formal medical private pharmacies	Pharmacies or accredited drug stores	Informal private	Trained provider	Public excluding community health workers	Community health workers	Formal medical private excluding pharmacies	Pharmacies or accredited drug stores	Self-treatment	No treatment seeking	Trained provider	Public	Private	Informal private
82 (74, 88)	27 (11, 52)	23 (13, 37)	60 (55, 64)	27 (23, 32)	-	40 (27, 54)	23 (11, 42)	10 (4, 21)	28 (24, 33)	7 (5, 10)	74 (67, 81)	84 (73, 91)	-
30 (23, 38)	9 (6, 14)	8 (5, 12)	37 (33, 40)	38 (34, 44)	-	34 (27, 41)	23 (17, 30)	12 (9, 17)	34 (30, 37)	7 (5, 9)	44 (36, 52)	31 (24, 39)	40 (26, 55)
-	-	-	66 (61, 70)	69 (64, 73)	-	-	-	-	68 (64, 72)	10 (7, 14)	80 (76, 83)	-	-
86 (82, 89)	36 (30, 44)	54 (26, 79)	84 (82, 86)	69 (66, 71)	93 (87, 97)	55 (49, 62)	32 (26, 40)	-	66 (63, 68)	9 (8, 11)	12 (10, 14)	10 (6, 15)	-
54 (43, 65)	11 (7, 19)	8 (5, 15)	42 (36, 48)	58 (49, 66)	-	48 (38, 58)	33 (24, 43)	46 (38, 54)	48 (43, 54)	12 (9, 16)	25 (17, 35)	21 (15, 27)	15 (8, 27)
-	-	-	-	16 (11, 23)	-	19 (10, 34)	-	-	17 (13, 23)	4 (2, 6)	14 (4, 38)	-	-
30 (22, 39)	8 (4, 16)	-	50 (45, 55)	63 (56, 70)	-	55 (46, 63)	57 (44, 69)	-	59 (54, 65)	18 (14, 24)	88 (80, 93)	86 (77, 92)	-
37 (23, 53)	-	4 (2, 8)	42 (37, 47)	41 (35, 47)	-	52 (33, 70)	-	24 (17, 31)	42 (37, 48)	10 (7, 14)	22 (15, 31)	15 (8, 25)	7 (3, 18)
57 (45, 67)	9 (2, 28)	25 (12, 45)	49 (44, 54)	31 (25, 37)	-	30 (21, 40)	44 (23, 66)	29 (16, 47)	31 (26, 37)	19 (14, 25)	93 (88, 96)	90 (77, 96)	-
82 (71, 89)	35 (25, 47)	14 (5, 33)	70 (64, 75)	84 (80, 88)	-	75 (65, 83)	76 (63, 85)	62 (49, 73)	81 (77, 84)	21 (15, 29)	87 (82, 91)	72 (64, 78)	80 (56, 93)
7 (3, 14)	-	3 (1, 13)	27 (22, 33)	13 (8, 19)	19 (10, 33)	13 (6, 25)	-	18 (9, 35)	14 (10, 19)	5 (3, 9)	9 (3, 26)	-	-
76 (61, 86)	-	4 (1, 14)	73 (67, 78)	55 (48, 62)	-	55 (39, 69)	-	21 (9, 41)	54 (48, 61)	7 (5, 11)	98 (94, 99)	97 (81, 99)	-
-	8 (3, 17)	5 (3, 9)	36 (30, 42)	61 (54, 68)	56 (38, 72)	-	17 (9, 30)	5 (2, 10)	50 (44, 57)	4 (3, 6)	35 (27, 43)	20 (9, 38)	-
-	-	-	70 (65, 74)	47 (40, 53)	57 (44, 70)	-	-	-	47 (41, 53)	10 (6, 17)	98 (97, 99)	-	-
8 (6, 9)	11 (7, 16)	3 (1, 5)	18 (17, 20)	64 (61, 66)	57 (39, 73)	51 (48, 53)	37 (32, 43)	23 (17, 31)	55 (53, 56)	19 (17, 21)	54 (50, 57)	50 (46, 53)	35 (22, 50)
70 (51, 84)	14 (4, 37)	-	67 (61, 73)	30 (22, 40)	60 (51, 68)	13 (4, 39)	31 (15, 54)	-	37 (31, 44)	2 (1, 4)	99 (95, 100)	-	-
-	4 (1, 9)	-	25 (20, 30)	12 (8, 19)	-	-	6 (1, 33)	-	12 (7, 19)	0 (0, 1)	22 (14, 32)	-	-
72 (56, 84)	13 (6, 26)	-	71 (67, 74)	77 (74, 81)	-	77 (61, 88)	41 (28, 55)	-	75 (71, 79)	19 (14, 24)	98 (96, 98)	93 (80, 97)	-
-	-	-	-	0 (0, 0)	-	1 (0, 5)	1 (0, 4)	-	0 (0, 1)	1 (0, 7)	-	-	-
45 (31, 60)	-	4 (2, 11)	66 (60, 72)	70 (60, 79)	83 (69, 91)	54 (37, 70)	-	10 (5, 17)	66 (59, 73)	7 (4, 10)	82 (74, 88)	56 (38, 73)	-
48 (43, 53)	20 (15, 28)	-	58 (54, 62)	72 (66, 76)	90 (84, 93)	72 (67, 77)	54 (42, 66)	-	70 (66, 74)	30 (23, 37)	89 (84, 93)	87 (82, 91)	-
76 (68, 83)	13 (8, 21)	-	55 (51, 60)	34 (28, 40)	-	49 (41, 57)	57 (48, 66)	-	42 (36, 47)	24 (19, 30)	96 (92, 98)	83 (73, 90)	-
79 (65, 89)	-	-	78 (73, 82)	42 (37, 47)	86 (72, 93)	54 (41, 67)	-	-	44 (40, 49)	10 (7, 13)	97 (95, 98)	94 (76, 99)	-
13 (6, 26)	-	9 (3, 25)	23 (18, 28)	2 (1, 5)	-	1 (0, 7)	-	0 (0, 0)	1 (1, 4)	1 (0, 3)	-	-	-

Data as of 17 November 2020

ACT: artemisinin-based combination therapy; DHS: demographic and health survey; MIS: malaria indicator survey; WHO: World Health Organization.

Sources: Nationally representative household survey data from DHS and MIS, compiled through WHO calculations.

























### ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>AMERICAS</b>								
Argentina <sup>1,2,3</sup>	2000	184 353	-	440	-	-	0	-
	2001	186 378	-	215	-	-	0	-
	2002	188 408	-	125	-	-	0	-
	2003	190 439	-	122	-	-	1	-
	2004	192 459	-	115	-	-	0	-
	2005	194 464	-	252	-	-	0	-
	2006	196 449	-	212	-	-	0	-
	2007	198 421	-	387	-	-	0	-
	2008	200 400	-	130	-	-	0	-
	2009	202 413	-	86	-	-	0	-
	2010	204 478	-	14	-	-	0	-
	2011	206 602	-	0	-	-	0	-
	2012	208 775	-	0	-	-	0	-
	2013	210 980	-	0	-	-	0	-
	2014	213 187	-	0	-	-	0	-
	2015	215 377	-	0	-	-	0	-
	2016	217 542	-	0	-	-	0	-
	2017	219 685	-	0	-	-	0	-
	2018	221 805	-	0	-	-	0	-
	2019	223 903	-	0	-	-	0	-
Belize <sup>1,2</sup>	2000	170 643	-	1 486	-	-	0	-
	2001	175 996	-	1 162	-	-	0	-
	2002	181 047	-	1 134	-	-	0	-
	2003	185 905	-	1 084	-	-	0	-
	2004	190 796	-	1 068	-	-	1	-
	2005	195 820	-	1 549	-	-	0	-
	2006	201 023	-	844	-	-	1	-
	2007	206 331	-	845	-	-	0	-
	2008	211 707	-	540	-	-	0	-
	2009	217 111	-	256	-	-	0	-
	2010	222 500	-	150	-	-	0	-
	2011	227 862	-	72	-	-	0	-
	2012	233 220	-	33	-	-	0	-
	2013	238 537	-	20	-	-	0	-
	2014	243 822	-	19	-	-	0	-
	2015	249 038	-	9	-	-	0	-
	2016	254 195	-	4	-	-	0	-
	2017	259 284	-	7	-	-	0	-
	2018	264 318	-	3	-	-	0	-
	2019	269 342	-	0	-	-	0	-
Bolivia (Plurinational State of)	2000	3 819 116	34 000	45 647	58 000	7	24	43
	2001	3 892 599	17 000	22 330	28 000	3	9	18
	2002	3 966 356	15 000	19 768	25 000	3	9	16
	2003	4 040 303	22 000	27 568	34 000	4	11	21
	2004	4 114 353	16 000	20 206	25 000	3	9	16
	2005	4 188 417	29 000	37 189	46 000	6	17	30
	2006	4 262 433	27 000	34 862	43 000	5	19	33
	2007	4 336 376	15 000	19 799	24 000	3	11	20
	2008	4 410 333	10 000	13 210	16 000	2	7	12
	2009	4 484 432	10 000	13 379	16 000	2	6	11
	2010	4 558 747	15 000	18 659	23 000	2	10	18
	2011	4 633 309	7 600	9 680	12 000	1	4	8
	2012	4 708 040	8 600	10 972	14 000	1	4	8
	2013	4 782 759	8 400	10 804	13 000	1	6	12
	2014	4 857 225	8 600	10 952	13 000	1	4	8
	2015	4 931 271	7 300	9 315	11 000	1	3	6
	2016	5 004 806	5 900	7 510	9 200	0	2	5
	2017	5 077 861	4 800	6 195	7 600	0	2	4
	2018	5 150 579	5 700	7 239	8 900	0	2	4
	2019	5 223 148	9 900	12 654	16 000	1	4	8
Brazil <sup>2</sup>	2000	35 482 438	643 000	760 760	886 000	-	245	-
	2001	35 970 799	407 000	467 114	533 000	-	142	-
	2002	36 446 116	363 000	407 200	457 000	-	95	-
	2003	36 907 274	425 000	465 651	513 000	-	104	-
	2004	37 353 321	481 000	516 739	561 000	-	102	-
	2005	37 783 803	624 000	658 276	706 000	-	123	-
	2006	38 197 972	561 000	584 183	623 000	-	110	-
	2007	38 596 477	478 000	534 516	578 000	-	93	-
	2008	38 982 161	322 000	335 694	358 000	-	68	-
	2009	39 358 958	316 000	328 858	351 000	-	85	-
	2010	39 729 868	348 000	389 809	422 000	-	76	-
	2011	40 095 452	273 000	284 024	303 000	-	70	-
	2012	40 455 322	248 000	258 095	276 000	-	60	-
	2013	40 810 285	176 000	196 793	213 000	-	40	-
	2014	41 161 038	142 000	148 071	158 000	-	36	-
	2015	41 507 764	154 000	162 341	173 000	-	35	-
	2016	41 851 100	129 000	134 862	144 000	-	35	-
	2017	42 190 269	193 000	201 475	215 000	-	34	-
	2018	42 522 271	207 000	217 900	232 000	-	44	-
	2019	42 843 057	169 000	177 967	190 000	-	36	-

## ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>AMERICAS</b>								
Colombia <sup>2</sup>	2000	8 773 678	154 000	210 720	270 000	-	124	-
	2001	8 912 266	246 000	333 024	424 000	-	168	-
	2002	9 049 394	218 000	291 432	368 000	-	162	-
	2003	9 184 115	192 000	254 224	319 000	-	118	-
	2004	9 315 195	151 000	197 464	245 000	-	126	-
	2005	9 441 780	129 000	166 899	206 000	-	87	-
	2006	9 564 247	127 000	165 418	205 000	-	77	-
	2007	9 683 047	136 000	177 615	221 000	-	68	-
	2008	9 797 609	86 000	111 811	139 000	-	54	-
	2009	9 907 215	84 000	110 555	138 000	-	28	-
	2010	10 011 853	125 000	164 479	206 000	-	42	-
	2011	10 109 278	64 000	84 072	105 000	-	23	-
	2012	10 200 703	64 000	84 176	105 000	-	24	-
	2013	10 293 636	55 000	72 310	90 000	-	10	-
	2014	10 398 179	43 000	57 024	71 000	-	17	-
	2015	10 520 599	51 000	66 603	83 000	-	18	-
	2016	10 665 474	88 000	115 550	144 000	-	36	-
	2017	10 828 150	56 000	73 861	92 000	-	19	-
	2018	10 994 461	71 000	93 468	117 000	-	9	-
	2019	11 144 649	91 000	119 302	149 000	-	3	-
Costa Rica <sup>1,2</sup>	2000	1 386 829	-	1 879	-	-	0	-
	2001	1 411 925	-	1 363	-	-	0	-
	2002	1 435 322	-	1 021	-	-	0	-
	2003	1 457 418	-	718	-	-	0	-
	2004	1 478 804	-	1 289	-	-	0	-
	2005	1 499 926	-	3 541	-	-	0	-
	2006	1 520 897	-	2 903	-	-	0	-
	2007	1 541 619	-	1 223	-	-	0	-
	2008	1 562 093	-	966	-	-	0	-
	2009	1 582 258	-	262	-	-	1	-
	2010	1 602 079	-	110	-	-	0	-
	2011	1 621 580	-	10	-	-	0	-
	2012	1 640 801	-	6	-	-	0	-
	2013	1 659 738	-	0	-	-	0	-
	2014	1 678 386	-	0	-	-	0	-
	2015	1 696 731	-	0	-	-	0	-
	2016	1 714 767	-	4	-	-	0	-
	2017	1 732 484	-	12	-	-	0	-
	2018	1 749 805	-	70	-	-	0	-
	2019	1 766 646	-	95	-	-	0	-
Dominican Republic	2000	4 666 170	1 300	1 524	1 800	-	6	-
	2001	4 736 280	1 100	1 315	1 600	-	17	-
	2002	4 805 890	1 400	1 685	2 000	-	11	-
	2003	4 874 931	1 600	1 983	2 400	-	12	-
	2004	4 943 303	2 500	3 046	3 600	-	16	-
	2005	5 010 953	4 000	4 950	5 900	-	16	-
	2006	5 077 833	3 700	4 535	5 400	-	10	-
	2007	5 144 028	2 900	3 478	4 100	-	17	-
	2008	5 209 699	1 900	2 365	2 800	-	11	-
	2009	5 275 057	1 700	2 115	2 500	-	14	-
	2010	5 340 264	2 600	3 202	3 800	-	15	-
	2011	5 405 317	1 700	2 088	2 500	-	10	-
	2012	5 470 147	1 000	1 232	1 500	-	8	-
	2013	5 534 763	610	751	900	-	5	-
	2014	5 599 185	480	566	650	-	4	-
	2015	5 663 352	660	779	900	-	3	-
	2016	5 727 282	720	851	990	-	2	-
	2017	5 790 831	420	491	570	-	1	-
	2018	5 853 645	640	750	870	-	1	-
	2019	5 915 232	1 400	1 592	1 800	-	4	-
Ecuador <sup>1,2</sup>	2000	369 527	-	104 528	-	-	66	-
	2001	376 333	-	108 903	-	-	84	-
	2002	383 000	-	86 757	-	-	64	-
	2003	389 592	-	52 065	-	-	46	-
	2004	396 198	-	28 730	-	-	37	-
	2005	402 884	-	17 050	-	-	22	-
	2006	409 690	-	9 863	-	-	9	-
	2007	416 601	-	8 464	-	-	8	-
	2008	423 571	-	4 891	-	-	5	-
	2009	430 526	-	4 120	-	-	6	-
	2010	437 423	-	1 871	-	-	0	-
	2011	444 206	-	1 219	-	-	0	-
	2012	450 915	-	544	-	-	0	-
	2013	457 715	-	368	-	-	0	-
	2014	464 836	-	242	-	-	0	-
	2015	472 418	-	618	-	-	0	-
	2016	480 551	-	1 191	-	-	0	-
	2017	489 125	-	1 275	-	-	1	-
	2018	497 838	-	1 653	-	-	0	-
	2019	506 268	-	1 803	-	-	0	-







## ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>AMERICAS</b>								
Panama <sup>2</sup>	2000	2 931 635	1 000	1 091	1 200	-	1	-
	2001	2 989 011	940	977	1 000	-	1	-
	2002	3 046 625	2 300	2 363	2 500	-	2	-
	2003	3 104 537	4 600	4 739	5 100	-	4	-
	2004	3 162 873	5 200	5 365	5 700	-	2	-
	2005	3 221 756	3 700	3 861	4 100	-	1	-
	2006	3 281 206	1 700	1 751	1 900	-	1	-
	2007	3 341 184	1 300	1 349	1 400	-	1	-
	2008	3 401 681	750	783	830	-	1	-
	2009	3 462 639	790	819	870	-	0	-
	2010	3 524 048	420	440	470	-	1	-
	2011	3 585 758	360	372	400	-	0	-
	2012	3 647 825	860	888	950	-	1	-
	2013	3 710 526	720	751	800	-	0	-
	2014	3 774 245	960	1 007	1 100	-	0	-
	2015	3 839 236	550	575	610	-	0	-
	2016	3 905 585	780	809	860	-	0	-
	2017	3 973 006	760	801	860	-	0	-
	2018	4 040 827	750	786	840	-	0	-
	2019	4 108 133	1 500	1 578	1 700	-	0	-
Paraguay <sup>1,2,3</sup>	2000	191 635	-	6 853	-	-	0	-
	2001	195 423	-	2 710	-	-	0	-
	2002	199 150	-	2 778	-	-	0	-
	2003	202 787	-	1 392	-	-	0	-
	2004	206 300	-	694	-	-	0	-
	2005	209 667	-	376	-	-	0	-
	2006	212 875	-	823	-	-	0	-
	2007	215 943	-	1 341	-	-	0	-
	2008	218 926	-	348	-	-	0	-
	2009	221 902	-	91	-	-	0	-
	2010	224 928	-	18	-	-	0	-
	2011	228 023	-	1	-	-	0	-
	2012	231 174	-	0	-	-	0	-
	2013	234 369	-	0	-	-	0	-
	2014	237 582	-	0	-	-	0	-
	2015	240 794	-	0	-	-	0	-
	2016	244 003	-	0	-	-	0	-
	2017	247 214	-	0	-	-	0	-
	2018	250 418	-	0	-	-	0	-
	2019	253 607	-	0	-	-	0	-
Peru <sup>2</sup>	2000	10 392 407	72 000	94 271	117 000	14	96	170
	2001	10 525 688	83 000	105 067	128 000	16	89	160
	2002	10 644 174	105 000	128 960	154 000	20	108	180
	2003	10 750 711	93 000	111 816	132 000	17	95	160
	2004	10 849 691	98 000	115 387	133 000	18	98	160
	2005	10 944 705	92 000	108 134	125 000	17	80	130
	2006	11 037 363	68 000	80 054	93 000	13	52	85
	2007	11 128 088	53 000	62 633	73 000	10	43	71
	2008	11 218 137	47 000	54 608	63 000	8	34	56
	2009	11 308 606	45 000	52 035	60 000	8	31	51
	2010	11 400 911	33 000	37 847	43 000	6	20	32
	2011	11 493 851	26 000	30 924	36 000	4	19	31
	2012	11 589 086	33 000	40 437	48 000	6	24	41
	2013	11 694 030	51 000	62 669	75 000	10	45	76
	2014	11 818 294	69 000	83 936	100 000	14	60	100
	2015	11 967 687	77 000	93 936	113 000	15	76	130
	2016	12 146 509	60 000	72 836	86 000	10	69	120
	2017	12 350 062	59 000	72 518	86 000	10	64	110
	2018	12 564 103	48 000	58 455	70 000	9	48	80
	2019	12 768 809	33 000	45 729	64 000	6	35	66
Suriname <sup>1,2</sup>	2000	69 558	-	11 361	-	-	24	-
	2001	70 389	-	16 003	-	-	23	-
	2002	71 225	-	12 837	-	-	15	-
	2003	72 068	-	10 982	-	-	18	-
	2004	72 916	-	8 378	-	-	7	-
	2005	73 770	-	9 131	-	-	1	-
	2006	74 631	-	3 289	-	-	1	-
	2007	75 501	-	1 741	-	-	1	-
	2008	76 378	-	2 709	-	-	0	-
	2009	77 263	-	2 380	-	-	0	-
	2010	78 151	-	1 712	-	-	1	-
	2011	79 045	-	771	-	-	1	-
	2012	79 942	-	356	-	-	0	-
	2013	80 835	-	729	-	-	1	-
	2014	81 719	-	401	-	-	1	-
	2015	82 584	-	81	-	-	0	-
	2016	83 433	-	76	-	-	0	-
	2017	84 262	-	40	-	-	1	-
	2018	85 073	-	29	-	-	0	-
	2019	85 867	-	95	-	-	0	-

### ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>AMERICAS</b>								
Venezuela (Bolivarian Republic of)	2000	12 096 224	31 000	35 530	42 000	5	26	44
	2001	12 323 235	21 000	23 843	28 000	3	15	25
	2002	12 550 203	31 000	35 041	41 000	5	19	31
	2003	12 775 812	33 000	37 524	44 000	5	27	45
	2004	12 998 297	49 000	55 005	65 000	8	30	50
	2005	13 216 222	47 000	52 999	62 000	8	34	55
	2006	13 425 095	39 000	43 654	51 000	6	33	53
	2007	13 623 800	44 000	48 852	57 000	7	37	61
	2008	13 817 913	33 000	37 496	44 000	5	26	43
	2009	14 015 505	37 000	41 943	49 000	6	36	58
	2010	14 219 971	48 000	57 926	73 000	8	53	91
	2011	14 443 936	48 000	53 584	62 000	8	47	76
	2012	14 680 413	55 000	61 873	72 000	9	56	92
	2013	14 890 523	82 000	92 159	106 000	13	104	170
	2014	15 021 486	95 000	106 079	122 000	16	110	180
	2015	15 040 913	142 000	159 661	184 000	25	150	240
	2016	14 925 624	251 000	281 897	327 000	44	261	420
	2017	14 701 240	429 000	482 617	556 000	79	424	690
	2018	14 443 558	423 000	475 212	547 000	82	426	690
2019	14 257 914	415 000	467 421	538 000	75	403	650	
<b>EASTERN MEDITERRANEAN</b>								
Afghanistan	2000	16 017 398	846 000	1 319 942	2 044 000	220	971	2 030
	2001	16 654 885	852 000	1 319 942	2 015 000	220	971	2 010
	2002	17 420 902	913 000	1 391 183	2 098 000	220	1 141	2 220
	2003	18 253 452	819 000	1 248 701	1 910 000	210	801	1 560
	2004	19 059 579	481 000	717 358	1 076 000	120	351	680
	2005	19 774 570	313 000	535 476	866 000	89	259	520
	2006	20 374 865	225 000	418 218	717 000	64	222	460
	2007	20 889 368	237 000	452 625	785 000	69	236	500
	2008	21 368 611	199 000	381 132	668 000	57	187	400
	2009	21 887 000	149 000	274 469	455 000	44	139	290
	2010	22 496 483	165 000	290 333	460 000	49	164	320
	2011	23 214 801	198 000	362 321	571 000	57	192	380
	2012	24 019 501	122 000	220 650	354 000	29	92	190
	2013	24 873 724	109 000	181 194	283 000	28	84	170
	2014	25 722 550	174 000	254 108	360 000	42	121	220
	2015	26 526 347	231 000	354 933	512 000	59	167	310
	2016	27 273 591	451 000	641 459	883 000	110	306	550
	2017	27 977 406	437 000	574 672	737 000	100	271	470
	2018	28 652 489	495 000	646 248	824 000	110	297	520
2019	29 322 964	328 000	424 653	539 000	66	175	310	
Djibouti <sup>1,2</sup>	2000	538 125	1 400	1 832	2 300	0	0	1
	2001	549 705	1 400	1 872	2 300	0	0	1
	2002	560 150	1 500	1 907	2 300	0	0	1
	2003	569 668	1 500	1 940	2 400	0	0	1
	2004	578 637	1 500	1 970	2 400	0	0	1
	2005	587 373	1 500	2 000	2 500	0	0	1
	2006	595 851	1 600	2 029	2 500	0	0	1
	2007	604 027	1 600	2 057	2 500	0	0	1
	2008	612 205	1 600	2 084	2 600	0	0	1
	2009	620 798	-	2 686	-	0	1	1
	2010	630 077	-	1 010	-	0	2	4
	2011	640 184	1 700	2 180	2 700	0	0	1
	2012	651 032	1 700	2 217	2 700	0	0	1
	2013	662 401	-	1 684	-	0	0	1
	2014	673 958	-	9 439	-	1	3	5
	2015	685 425	-	9 473	-	0	24	37
	2016	696 763	-	13 804	-	1	30	48
	2017	707 999	-	14 671	-	1	24	38
	2018	719 115	-	25 319	-	2	44	69
2019	730 089	-	49 402	-	5	97	150	

## ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>EASTERN MEDITERRANEAN</b>								
Egypt <sup>1,2</sup>	2000	68 831 561	-	0	-	-	0	-
	2001	70 152 662	-	0	-	-	0	-
	2002	71 485 044	-	0	-	-	0	-
	2003	72 826 102	-	0	-	-	0	-
	2004	74 172 073	-	0	-	-	0	-
	2005	75 523 576	-	0	-	-	0	-
	2006	76 873 670	-	0	-	-	0	-
	2007	78 232 124	-	0	-	-	0	-
	2008	79 636 081	-	0	-	-	0	-
	2009	81 134 789	-	0	-	-	0	-
	2010	82 761 244	-	0	-	-	0	-
	2011	84 529 251	-	0	-	-	0	-
	2012	86 422 240	-	0	-	-	0	-
	2013	88 404 652	-	0	-	-	0	-
	2014	90 424 668	-	0	-	-	0	-
	2015	92 442 549	-	0	-	-	0	-
	2016	94 447 071	-	0	-	-	0	-
	2017	96 442 590	-	0	-	-	0	-
	2018	98 423 602	-	0	-	-	0	-
	2019	100 388 076	-	0	-	-	0	-
Iran (Islamic Republic of) <sup>1,2</sup>	2000	670 014	-	19 716	-	-	4	-
	2001	678 445	-	19 303	-	-	2	-
	2002	686 977	-	15 558	-	-	2	-
	2003	695 535	-	23 562	-	-	5	-
	2004	703 992	-	13 821	-	-	1	-
	2005	712 273	-	18 966	-	-	1	-
	2006	720 364	-	15 909	-	-	1	-
	2007	728 345	-	15 712	-	-	3	-
	2008	736 351	-	8 349	-	-	3	-
	2009	744 562	-	4 345	-	-	0	-
	2010	753 115	-	1 847	-	-	0	-
	2011	762 022	-	1 632	-	-	0	-
	2012	771 262	-	756	-	-	0	-
	2013	780 880	-	479	-	-	0	-
	2014	790 925	-	358	-	-	0	-
	2015	801 405	-	167	-	-	1	-
	2016	812 348	-	81	-	-	0	-
	2017	823 680	-	60	-	-	1	-
	2018	835 180	-	0	-	-	0	-
	2019	846 550	-	0	-	-	0	-
Iraq	2000	3 054 686	-	1 860	-	-	0	-
	2001	3 147 063	-	1 265	-	-	0	-
	2002	3 241 149	-	952	-	-	0	-
	2003	3 333 785	-	288	-	-	0	-
	2004	3 420 798	-	148	-	-	0	-
	2005	3 499 896	-	44	-	-	0	-
	2006	3 568 256	-	23	-	-	0	-
	2007	3 628 461	-	2	-	-	0	-
	2008	3 690 146	-	2	-	-	0	-
	2009	3 766 510	-	0	-	-	0	-
	2010	3 866 457	-	0	-	-	0	-
	2011	3 994 289	-	0	-	-	0	-
	2012	4 145 701	-	0	-	-	0	-
	2013	4 310 417	-	0	-	-	0	-
	2014	4 473 553	-	0	-	-	0	-
	2015	4 624 394	-	0	-	-	0	-
	2016	4 759 382	-	0	-	-	0	-
	2017	4 881 862	-	0	-	-	0	-
	2018	4 996 368	-	0	-	-	0	-
	2019	5 110 272	-	0	-	-	0	-
Morocco <sup>1,2,3</sup>	2000	28 793 672	-	3	-	-	0	-
	2001	29 126 323	-	0	-	-	0	-
	2002	29 454 765	-	19	-	-	0	-
	2003	29 782 884	-	4	-	-	0	-
	2004	30 115 196	-	1	-	-	0	-
	2005	30 455 563	-	0	-	-	0	-
	2006	30 804 689	-	0	-	-	0	-
	2007	31 163 670	-	0	-	-	0	-
	2008	31 536 807	-	0	-	-	0	-
	2009	31 929 087	-	0	-	-	0	-
	2010	32 343 384	-	0	-	-	0	-
	2011	32 781 860	-	0	-	-	0	-
	2012	33 241 898	-	0	-	-	0	-
	2013	33 715 704	-	0	-	-	0	-
	2014	34 192 360	-	0	-	-	0	-
	2015	34 663 608	-	0	-	-	0	-
	2016	35 126 276	-	0	-	-	0	-
	2017	35 581 260	-	0	-	-	0	-
	2018	36 029 088	-	0	-	-	0	-
	2019	36 471 768	-	0	-	-	0	-





## ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>EUROPEAN</b>								
Armenia <sup>1,2,3</sup>	2000	3 069 597	-	141	-	-	0	-
	2001	3 050 686	-	79	-	-	0	-
	2002	3 033 976	-	52	-	-	0	-
	2003	3 017 938	-	29	-	-	0	-
	2004	3 000 715	-	47	-	-	0	-
	2005	2 981 262	-	7	-	-	0	-
	2006	2 958 301	-	0	-	-	0	-
	2007	2 932 615	-	0	-	-	0	-
	2008	2 907 615	-	0	-	-	0	-
	2009	2 888 094	-	0	-	-	0	-
	2010	2 877 314	-	0	-	-	0	-
	2011	2 876 536	-	0	-	-	0	-
	2012	2 884 239	-	0	-	-	0	-
	2013	2 897 593	-	0	-	-	0	-
	2014	2 912 403	-	0	-	-	0	-
	2015	2 925 559	-	0	-	-	0	-
	2016	2 936 147	-	0	-	-	0	-
	2017	2 944 789	-	0	-	-	0	-
	2018	2 951 741	-	0	-	-	0	-
	2019	2 957 728	-	0	-	-	0	-
Azerbaijan <sup>1,2</sup>	2000	186 823	-	1 526	-	-	0	-
	2001	188 537	-	1 058	-	-	0	-
	2002	190 372	-	506	-	-	0	-
	2003	192 312	-	482	-	-	0	-
	2004	194 325	-	386	-	-	0	-
	2005	196 388	-	242	-	-	0	-
	2006	198 493	-	143	-	-	0	-
	2007	200 657	-	108	-	-	0	-
	2008	202 902	-	72	-	-	0	-
	2009	205 260	-	78	-	-	0	-
	2010	207 746	-	50	-	-	0	-
	2011	210 364	-	4	-	-	0	-
	2012	213 087	-	3	-	-	0	-
	2013	215 865	-	0	-	-	0	-
	2014	218 629	-	0	-	-	0	-
	2015	221 323	-	0	-	-	0	-
	2016	223 928	-	0	-	-	0	-
	2017	226 442	-	0	-	-	0	-
	2018	228 839	-	0	-	-	0	-
	2019	231 097	-	0	-	-	0	-
Georgia <sup>1,2</sup>	2000	43 621	-	245	-	-	0	-
	2001	42 969	-	438	-	-	0	-
	2002	42 585	-	474	-	-	0	-
	2003	42 389	-	316	-	-	0	-
	2004	42 258	-	257	-	-	0	-
	2005	42 101	-	155	-	-	0	-
	2006	41 897	-	59	-	-	0	-
	2007	41 668	-	24	-	-	0	-
	2008	41 426	-	6	-	-	0	-
	2009	41 194	-	1	-	-	0	-
	2010	40 990	-	0	-	-	0	-
	2011	40 810	-	0	-	-	0	-
	2012	40 640	-	0	-	-	0	-
	2013	40 487	-	0	-	-	0	-
	2014	40 353	-	0	-	-	0	-
	2015	40 241	-	0	-	-	0	-
	2016	40 154	-	0	-	-	0	-
	2017	40 087	-	0	-	-	0	-
	2018	40 029	-	0	-	-	0	-
	2019	39 967	-	0	-	-	0	-
Kazakhstan <sup>1,2</sup>	2000	14 922 724	-	0	-	-	0	-
	2001	14 910 207	-	0	-	-	0	-
	2002	14 976 184	-	0	-	-	0	-
	2003	15 100 045	-	0	-	-	0	-
	2004	15 250 016	-	0	-	-	0	-
	2005	15 402 803	-	0	-	-	0	-
	2006	15 551 263	-	0	-	-	0	-
	2007	15 702 112	-	0	-	-	0	-
	2008	15 862 126	-	0	-	-	0	-
	2009	16 043 015	-	0	-	-	0	-
	2010	16 252 273	-	0	-	-	0	-
	2011	16 490 669	-	0	-	-	0	-
	2012	16 751 523	-	0	-	-	0	-
	2013	17 026 118	-	0	-	-	0	-
	2014	17 302 619	-	0	-	-	0	-
	2015	17 572 010	-	0	-	-	0	-
	2016	17 830 902	-	0	-	-	0	-
	2017	18 080 023	-	0	-	-	0	-
	2018	18 319 616	-	0	-	-	0	-
	2019	18 551 428	-	0	-	-	0	-

## ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>EUROPEAN</b>								
Kyrgyzstan <sup>1,2,3</sup>	2000	3 838 155	-	12	-	-	0	-
	2001	3 871 014	-	28	-	-	0	-
	2002	3 893 352	-	2 743	-	-	0	-
	2003	3 910 559	-	468	-	-	0	-
	2004	3 930 416	-	93	-	-	0	-
	2005	3 958 765	-	226	-	-	0	-
	2006	3 997 015	-	317	-	-	0	-
	2007	4 043 819	-	96	-	-	0	-
	2008	4 098 875	-	18	-	-	0	-
	2009	4 161 072	-	4	-	-	0	-
	2010	4 229 392	-	3	-	-	0	-
	2011	4 303 983	-	0	-	-	0	-
	2012	4 384 834	-	0	-	-	0	-
	2013	4 470 423	-	0	-	-	0	-
	2014	4 558 726	-	0	-	-	0	-
	2015	4 648 118	-	0	-	-	0	-
	2016	4 737 975	-	0	-	-	0	-
	2017	4 827 987	-	0	-	-	0	-
	2018	4 917 139	-	0	-	-	0	-
	2019	5 004 363	-	0	-	-	0	-
Tajikistan <sup>1,2</sup>	2000	2 076 253	-	19 064	-	-	0	-
	2001	2 110 382	-	11 387	-	-	0	-
	2002	2 146 571	-	6 160	-	-	0	-
	2003	2 184 877	-	5 428	-	-	0	-
	2004	2 225 238	-	3 588	-	-	0	-
	2005	2 267 632	-	2 309	-	-	0	-
	2006	2 312 145	-	1 344	-	-	0	-
	2007	2 358 930	-	635	-	-	0	-
	2008	2 408 114	-	318	-	-	0	-
	2009	2 459 827	-	164	-	-	0	-
	2010	2 514 150	-	111	-	-	0	-
	2011	2 570 967	-	65	-	-	0	-
	2012	2 630 195	-	18	-	-	0	-
	2013	2 691 967	-	3	-	-	0	-
	2014	2 756 444	-	2	-	-	0	-
	2015	2 823 642	-	0	-	-	0	-
	2016	2 893 634	-	0	-	-	0	-
	2017	2 966 010	-	0	-	-	0	-
	2018	3 039 682	-	0	-	-	0	-
	2019	3 113 221	-	0	-	-	0	-
Turkey <sup>1,2</sup>	2000	4 110 612	-	11 432	-	-	0	-
	2001	4 172 495	-	10 812	-	-	0	-
	2002	4 234 448	-	10 224	-	-	0	-
	2003	4 295 811	-	9 222	-	-	0	-
	2004	4 355 710	-	5 302	-	-	0	-
	2005	4 413 725	-	2 084	-	-	0	-
	2006	4 469 192	-	796	-	-	0	-
	2007	4 522 820	-	313	-	-	0	-
	2008	4 577 210	-	166	-	-	0	-
	2009	4 635 891	-	38	-	-	0	-
	2010	4 701 254	-	0	-	-	0	-
	2011	4 773 811	-	0	-	-	0	-
	2012	4 852 318	-	0	-	-	0	-
	2013	4 935 154	-	0	-	-	0	-
	2014	5 019 902	-	0	-	-	0	-
	2015	5 104 412	-	0	-	-	0	-
	2016	5 188 811	-	0	-	-	0	-
	2017	5 272 570	-	0	-	-	0	-
	2018	5 352 105	-	0	-	-	0	-
	2019	5 422 923	-	0	-	-	0	-
Turkmenistan <sup>1,2,3</sup>	2000	293 548	-	24	-	-	0	-
	2001	296 665	-	8	-	-	0	-
	2002	299 651	-	18	-	-	0	-
	2003	302 623	-	7	-	-	0	-
	2004	305 720	-	3	-	-	0	-
	2005	309 052	-	1	-	-	0	-
	2006	312 657	-	1	-	-	0	-
	2007	316 559	-	0	-	-	0	-
	2008	320 824	-	0	-	-	0	-
	2009	325 516	-	0	-	-	0	-
	2010	330 668	-	0	-	-	0	-
	2011	336 314	-	0	-	-	0	-
	2012	342 413	-	0	-	-	0	-
	2013	348 814	-	0	-	-	0	-
	2014	355 311	-	0	-	-	0	-
	2015	361 743	-	0	-	-	0	-
	2016	368 054	-	0	-	-	0	-
	2017	374 248	-	0	-	-	0	-
	2018	380 308	-	0	-	-	0	-
	2019	386 236	-	0	-	-	0	-



### ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>EUROPEAN</b>								
Uzbekistan <sup>1,2,3</sup>	2000	24 769	-	126	-	-	0	-
	2001	25 108	-	77	-	-	0	-
	2002	25 431	-	74	-	-	0	-
	2003	25 749	-	74	-	-	0	-
	2004	26 077	-	66	-	-	0	-
	2005	26 427	-	102	-	-	0	-
	2006	26 804	-	73	-	-	0	-
	2007	27 204	-	30	-	-	0	-
	2008	27 626	-	7	-	-	0	-
	2009	28 065	-	0	-	-	0	-
	2010	28 515	-	3	-	-	0	-
	2011	28 977	-	0	-	-	0	-
	2012	29 449	-	0	-	-	0	-
	2013	29 932	-	0	-	-	0	-
	2014	30 426	-	0	-	-	0	-
	2015	30 929	-	0	-	-	0	-
	2016	31 441	-	0	-	-	0	-
	2017	31 959	-	0	-	-	0	-
	2018	32 476	-	0	-	-	0	-
2019	32 981	-	0	-	-	0	-	
<b>SOUTH-EAST ASIA</b>								
Bangladesh	2000	13 727 050	42 000	78 958	128 000	7	166	390
	2001	13 988 438	43 000	80 462	131 000	7	170	400
	2002	14 245 368	42 000	81 940	132 000	7	173	410
	2003	14 494 139	43 000	83 371	135 000	7	176	420
	2004	14 730 151	44 000	84 728	137 000	7	179	420
	2005	14 950 487	46 000	85 995	140 000	7	181	440
	2006	15 153 253	41 000	68 539	103 000	6	139	290
	2007	15 340 271	63 000	75 032	88 000	8	155	270
	2008	15 517 025	96 000	121 084	151 000	12	264	480
	2009	15 691 292	109 000	128 228	149 000	15	251	430
	2010	15 868 787	59 000	68 774	80 000	6	165	290
	2011	16 051 340	54 000	63 356	73 000	5	155	270
	2012	16 237 645	31 000	35 747	41 000	3	87	150
	2013	16 426 435	23 000	25 366	29 000	2	60	100
	2014	16 615 254	49 000	54 801	61 000	4	133	220
	2015	16 802 238	41 000	46 361	52 000	4	111	190
	2016	16 987 281	29 000	32 789	37 000	2	77	130
	2017	17 170 973	30 000	34 766	40 000	3	80	140
	2018	17 352 838	11 000	12 708	15 000	0	27	48
2019	17 532 354	18 000	21 146	25 000	1	47	82	
Bhutan <sup>1,2</sup>	2000	437 350	-	5 935	-	-	15	-
	2001	446 695	-	5 982	-	-	14	-
	2002	455 858	-	6 511	-	-	11	-
	2003	464 601	-	3 806	-	-	14	-
	2004	472 718	-	2 670	-	-	7	-
	2005	480 070	-	1 825	-	-	5	-
	2006	486 478	-	1 868	-	-	7	-
	2007	492 006	-	793	-	-	2	-
	2008	496 992	-	329	-	-	2	-
	2009	501 963	-	972	-	-	4	-
	2010	507 271	-	436	-	-	2	-
	2011	513 039	-	194	-	-	1	-
	2012	519 170	-	82	-	-	1	-
	2013	525 573	-	15	-	-	0	-
	2014	532 099	-	19	-	-	0	-
	2015	538 634	-	34	-	-	0	-
	2016	545 162	-	15	-	-	0	-
	2017	551 716	-	11	-	-	0	-
	2018	558 253	-	6	-	-	0	-
2019	564 689	-	2	-	-	0	-	









## ANNEX 3 – F. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND ESTIMATED MALARIA CASES AND DEATHS, 2000–2019

WHO region Country/area	Year	Population denominator for incidence and mortality rate	Cases			Deaths		
			Lower	Point	Upper	Lower	Point	Upper
<b>WESTERN PACIFIC</b>								
Vanuatu	2000	184 964	13 000	23 167	38 000	2	34	75
	2001	189 209	13 000	18 702	27 000	1	24	49
	2002	193 927	25 000	36 655	53 000	4	52	100
	2003	198 960	28 000	42 687	64 000	5	68	140
	2004	204 123	23 000	33 102	45 000	3	49	93
	2005	209 282	17 000	25 624	39 000	3	35	71
	2006	214 379	14 000	22 943	35 000	2	30	63
	2007	219 464	15 000	27 312	117 000	3	36	170
	2008	224 700	13 000	26 771	116 000	2	37	180
	2009	230 244	8 100	14 887	25 000	1	22	51
	2010	236 216	13 000	15 669	20 000	1	20	35
	2011	242 658	8 900	11 631	16 000	1	14	27
	2012	249 505	6 400	8 394	11 000	-	0	-
	2013	256 637	4 100	5 326	7 200	-	0	-
	2014	263 888	1 900	2 427	3 300	-	0	-
	2015	271 128	680	787	920	-	0	-
	2016	278 326	3 200	4 177	5 600	-	0	-
	2017	285 499	1 700	2 268	3 100	-	0	-
	2018	292 675	900	1 167	1 600	-	0	-
	2019	299 882	800	1 047	1 400	-	0	-
Viet Nam	2000	58 893 103	158 000	201 414	270 000	22	421	800
	2001	59 506 334	148 000	185 145	241 000	21	380	700
	2002	60 089 950	105 000	131 451	172 000	14	271	500
	2003	60 655 409	78 000	96 592	125 000	11	197	360
	2004	61 216 381	47 000	56 559	72 000	6	115	210
	2005	61 783 751	34 000	40 604	51 000	4	79	140
	2006	62 362 205	37 000	43 620	54 000	4	92	160
	2007	62 953 293	24 000	28 022	34 000	2	53	92
	2008	63 560 457	16 000	17 911	22 000	1	37	65
	2009	64 186 031	21 000	22 853	26 000	1	47	81
	2010	64 831 191	21 000	22 959	26 000	2	45	75
	2011	65 497 232	19 000	20 206	23 000	2	35	58
	2012	66 183 027	22 000	23 838	27 000	2	40	66
	2013	66 883 664	19 000	20 760	23 000	2	33	55
	2014	67 592 103	18 000	19 060	21 000	2	29	47
	2015	68 301 988	10 000	11 283	13 000	1	16	25
	2016	69 011 962	4 600	5 024	5 600	0	7	12
	2017	69 719 636	5 000	5 481	6 100	0	9	15
	2018	70 416 327	5 300	5 794	6 500	0	9	16
	2019	71 091 518	8 900	9 702	11 000	0	17	29

Data as of 17 November 2020

"-" refers to not applicable.

<sup>1</sup> The number of indigenous malaria cases registered by the NMPs is reported here without further adjustments.

<sup>2</sup> The number of indigenous malaria deaths registered by the NMPs is reported here without further adjustments.

<sup>3</sup> Certified malaria free countries are included in this listing for historical purposes.

<sup>4</sup> South Sudan became an independent state on 9 July 2011 and a Member State of WHO on 27 September 2011. South Sudan and Sudan have distinct epidemiological profiles comprising high-transmission and low-transmission areas respectively. For this reason, data up to June 2011 from the Sudanese high-transmission areas (10 southern states, which correspond to South Sudan) and low-transmission areas (15 northern states which correspond to contemporary Sudan) are reported separately.

Note: Population denominator for incidence and mortality rate is based on the United Nations population, times the proportion of the population at risk at baseline.







## ANNEX 3 – G. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND REPORTED MALARIA CASES BY PLACE OF CARE, 2019

WHO region Country/area	Population			
	UN population	At risk (low + high)	At risk (high)	Number of people living in active foci
<b>AFRICAN</b>				
Angola	31 825 299	31 825 299	31 825 299	-
Benin	11 801 151	11 801 151	11 801 151	-
Botswana	2 303 703	1 527 309	97 032	486 563
Burkina Faso*	20 321 383	20 321 383	20 321 383	-
Burundi	11 530 577	11 530 577	11 530 577	-
Cabo Verde	549 936	142 983	0	4 521
Cameroon	25 876 387	25 876 387	18 372 235	-
Central African Republic	4 745 179	4 745 179	4 745 179	-
Chad	15 946 882	15 772 264	10 741 022	-
Comoros	850 891	850 891	404 854	873 724
Congo	5 380 504	5 380 504	5 380 504	-
Côte d'Ivoire	25 716 554	25 716 554	25 716 554	-
Democratic Republic of the Congo	86 790 564	86 790 564	84 186 847	-
Equatorial Guinea	1 355 982	1 355 982	1 355 982	-
Eritrea	3 497 117	3 497 117	2 482 953	-
Eswatini	1 148 133	321 477	0	-
Ethiopia	112 078 736	76 213 540	30 485 416	-
Gabon	2 172 578	2 172 578	2 172 578	-
Gambia	2 347 696	2 347 696	2 347 696	-
Ghana	30 417 858	30 417 858	30 417 858	-
Guinea	12 771 246	12 771 246	12 771 246	-
Guinea-Bissau	1 920 917	1 920 917	1 920 917	-
Kenya	52 573 968	52 573 968	36 904 297	-
Liberia	4 937 374	4 937 374	4 937 374	-
Madagascar	26 969 306	26 969 306	23 670 420	-
Malawi	18 628 749	18 628 749	18 628 749	-
Mali	19 658 023	19 658 023	17 918 681	-
Mauritania	4 525 698	4 525 698	2 917 627	-
Mayotte	-	-	-	-
Mozambique	30 366 043	30 366 043	30 366 043	-
Namibia	2 494 524	1 980 028	1 151 497	-
Niger	23 310 719	23 310 719	23 310 719	-
Nigeria	200 963 608	200 963 608	153 491 985	-
Rwanda	12 626 938	12 626 938	12 626 938	-
Sao Tome and Principe	215 048	215 048	215 048	0
Senegal	16 296 362	16 296 362	16 202 332	0
Sierra Leone	7 813 207	7 813 207	7 813 207	-
South Africa	58 558 268	5 855 827	2 342 331	-
South Sudan <sup>1</sup>	11 062 114	11 062 114	11 062 114	-
Togo	8 082 359	8 082 359	8 082 359	-
Uganda	44 269 584	44 269 584	44 269 584	-
United Republic of Tanzania <sup>2</sup>	58 005 458	58 005 458	4 234 398	-
Mainland	56 364 236	56 364 236	41 145 892	-
Zanzibar	1 641 222	1 641 222	1 005 806	-
Zambia	17 861 034	17 861 034	17 861 034	-
Zimbabwe	14 645 473	11 532 241	4 190 949	-
<b>AMERICAS</b>				
Belize	390 351	269 342	0	18 968
Bolivia (Plurinational State of)	11 513 102	5 223 149	287 597	14 869
Brazil	211 049 544	42 843 057	4 854 140	-
Colombia	50 339 446	11 144 650	5 058 108	9 710 964
Costa Rica	5 047 561	1 766 646	50 476	172 541
Dominican Republic	10 738 957	5 915 232	151 956	-
Ecuador	17 373 657	506 268	158 795	-
El Salvador	6 453 550	1 310 071	0	0
French Guiana	291 000	161 004	26 859	-
Guatemala	17 581 476	13 272 608	2 398 641	-
Guyana	782 775	782 775	85 432	-
Haiti	11 263 079	10 062 660	2 730 058	-
Honduras	9 746 115	8 828 031	2 484 090	-

Public sector		Private sector		Community level	
Presumed	Confirmed	Presumed	Confirmed	Presumed	Confirmed
475 810	6 575 539	-	-	0	479 439
73 703	2 220 806	114 944	438 805	-	806 532
0	272	-	-	-	-
358 631	5 873 053	238 970	312 409	-	86 458
19 299	7 569 303	5 011	542 498	-	1 228 631
0	40	0	0	-	-
110 739	1 505 211	80 591	1 122 980	-	191 091
252 123	1 410 908	39 414	235 859	-	770 193
223 617	1 342 071	0	13 574	54 372	388 784
0	17 697	-	-	-	-
427 959	135 947	-	-	-	-
0	5 895 048	0	40 130	0	0
-	20 263 277	-	-	0	1 670 850
17 993	65 403 <sup>3</sup>	-	-	-	-
-	68 375	-	-	-	25 503
0	378	0	212	-	-
111 297	904 496	-	-	-	-
89 735	53 113	-	-	0	0
0	50 878	-	2 258	-	250
347 037	3 274 299	175 948	1 107 581	65 435	1 733 387
-	1 783 753 <sup>5</sup>	0	50 103	0	309 369
300 936	143 457	23 957	4 343	12 116	4 150
-	4 656 702 <sup>4</sup>	-	-	-	362 687
102 155	707 630	23 800	208 215	-	-
13 476	1 016 327	-	-	-	-
21 813	4 241 905	-	-	0	866 201
360 269	2 884 919	-	-	25 433	297 905
120 251	14 869	-	-	-	-
-	-	-	-	-	-
39 436	10 864 677	-	-	7 154	870 249
0	3 404	0	0	0	12
-	3 108 340	0	77 796	0	248 027
2 923 017	17 322 638	646 861	2 032 897	-	451 380
0	1 075 010	0	449 269	0	2 048 482
0	2 457	-	-	0	285
2 849	274 467 <sup>4</sup>	-	-	1 689	80 241
2 442 191	2 407 505 <sup>5</sup>	-	-	-	-
0	13 833	-	-	-	-
2 160 920	1 863 823 <sup>3</sup>	-	-	-	-
64 879	1 261 696	109 907	349 668	0	794 727
1 141 110	10 086 845	391 764	1 681 636	77 557	2 213 881
-	-	-	-	-	-
73 626	5 970 934	16 122	318 208	-	-
0	6 963 <sup>4</sup>	-	-	0	0
212 670	5 082 697	-	-	-	-
-	186 050	-	8 761	-	122 123
-	2 <sup>5</sup>	-	-	-	-
-	9 357 <sup>5</sup>	-	-	-	-
-	157 454 <sup>4</sup>	-	-	-	-
-	80 415 <sup>5</sup>	-	-	-	-
-	145 <sup>5</sup>	-	-	-	-
0	1 314 <sup>5</sup>	-	-	-	-
7	1 909 <sup>5</sup>	-	-	-	-
0	3	0	0	0	0
0	212 <sup>4</sup>	-	-	-	-
-	2 072 <sup>5</sup>	-	-	-	-
0	18 826 <sup>5</sup>	-	-	-	-
-	6 717	-	1 968	-	2 002
-	391 <sup>5</sup>	-	-	-	-

## ANNEX 3 – G. POPULATION DENOMINATOR FOR CASE INCIDENCE AND MORTALITY RATE, AND REPORTED MALARIA CASES BY PLACE OF CARE, 2019

WHO region Country/area	Population			
	UN population	At risk (low + high)	At risk (high)	Number of people living in active foci
<b>AMERICAS</b>				
Mexico	127 575 524	2 704 601	127 576	1 975 222
Nicaragua	6 545 503	2 857 112	561 801	-
Panama	4 246 440	4 108 133	178 945	46 429
Peru	32 510 462	12 768 809	1 627 474	-
Suriname	581 363	85 867	24 685	1 157
Venezuela (Bolivarian Republic of)	28 515 829	14 257 915	5 913 755	-
<b>EASTERN MEDITERRANEAN</b>				
Afghanistan	38 041 754	29 322 964	10 358 009	-
Djibouti	973 557	730 090	341 796	-
Iran (Islamic Republic of)	82 913 888	846 551	0	72 749
Pakistan	216 565 320	212 907 532	62 624 194	-
Saudi Arabia	34 268 533	2 745 252	0	143 632
Somalia	15 442 906	15 442 906	7 859 976	-
Sudan	42 813 236	42 813 236	37 204 702	-
Yemen	29 161 922	18 801 274	11 219 175	-
<b>SOUTH-EAST ASIA</b>				
Bangladesh	163 046 168	17 532 354	2 059 273	-
Bhutan	763 094	564 690	99 202	16 742
Democratic People's Republic of Korea	25 666 158	10 022 121	1 441 668	1 671 952
India	1 366 417 920	1 276 780 904	165 760 158	-
Indonesia	270 625 584	270 625 584	17 303 800	-
Myanmar	54 045 422	32 166 754	8 545 122	-
Nepal	28 608 715	8 304 538	1 495 378	276 247
Thailand	69 625 584	13 212 151	1 541 510	274 079
Timor-Leste	1 293 120	1 216 192	437 948	0
<b>WESTERN PACIFIC</b>				
Cambodia	16 486 542	11 659 118	7 934 313	-
China	1 441 860 352	603 014 836	201 860	0
Lao People's Democratic Republic	7 169 456	3 730 555	3 730 555	-
Malaysia <sup>6</sup>	31 949 789	1 277 992	958 494	9 211
Papua New Guinea	8 776 119	8 776 119	8 249 552	-
Philippines	108 116 620	813 684	428 437	519 001
Republic of Korea	51 225 322	3 585 773	0	-
Solomon Islands	669 821	663 123	663 123	-
Vanuatu	299 882	299 882	260 672	-
Viet Nam	96 462 116	71 091 518	6 557 012	-
<b>REGIONAL SUMMARY</b>				
African	1 045 213 130	950 833 144	789 192 269	1 364 808
Americas	552 545 734	138 867 930	26 720 388	11 940 150
Eastern Mediterranean	460 181 116	323 609 805	129 607 852	216 381
South-East Asia	1 980 091 765	1 630 425 288	198 684 059	2 239 020
Western Pacific	1 763 016 019	704 912 600	28 984 018	528 212
<b>Total</b>	<b>5 801 047 764</b>	<b>3 748 648 767</b>	<b>1 173 188 586</b>	<b>16 288 571</b>

RDT: rapid diagnostic testing; UN: United Nations; WHO: World Health Organization.

“-” refers to not applicable or data not available.

\* Double counting of microscopy and RDT reported, but proportion is not indicated.

<sup>1</sup> In May 2013, South Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [https://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

<sup>2</sup> Where national data for the United Republic of Tanzania are unavailable, refer to Mainland and Zanzibar.

Public sector		Private sector		Community level	
Presumed	Confirmed	Presumed	Confirmed	Presumed	Confirmed
-	641	-	-	-	-
-	13 226 <sup>5</sup>	-	-	-	-
-	1 597 <sup>5</sup>	-	-	-	-
-	24 324	-	-	-	-
-	215 <sup>5</sup>	-	-	-	-
-	398 285 <sup>5</sup>	-	-	-	-
976	120 367	-	4 226	58	47 405
-	49 402	-	-	-	-
-	1 189 <sup>3</sup>	-	-	-	5
-	290 712	0	122 821	-	-
0	2 152 <sup>4</sup>	-	-	-	-
25 688	39 687 <sup>3</sup>	-	-	-	-
1 816 930	1 752 011	-	-	-	-
31 800	113 095	19 064	41 183	-	7 544
0	3 455	0	46	0	13 724
0	42	-	-	0	0
-	1 869 <sup>3</sup>	-	-	-	-
-	338 494 <sup>5</sup>	-	-	-	-
-	205 352	-	37 153	-	8 139
-	9 717	-	2 257	-	44 437
548	551	180	134	-	11
-	4 143	-	421	-	757
0	8	0	1	0	0
0	12 843	-	-	0	16 598
5	2 482	-	-	-	-
-	4 283	-	606	-	1 798
-	3 941	-	-	0	0
-	640 703	-	-	-	-
0	1 846	0	89	0	3 311
0	196	-	-	0	0
12 923	71 530	426	1 093	-	-
0	576 <sup>3</sup>	-	-	-	-
5 887	4 665	-	100	-	-
12 487 541	132 207 015	1 867 289	8 997 202	243 756	16 050 837
7	717 105	0	1 968	0	2 002
1 875 394	2 368 615	19 064	168 230	58	54 954
548	563 631	180	40 012	0	67 068
18 815	743 065	426	1 888	0	21 707
<b>14 382 305</b>	<b>136 599 431</b>	<b>1 886 959</b>	<b>9 209 300</b>	<b>243 814</b>	<b>16 196 568</b>

Data as of 17 November 2020

<sup>3</sup> Figures reported for the public sector include cases detected at the community level.

<sup>4</sup> Figures reported for the public sector include cases detected in the private sector.

<sup>5</sup> Figures reported for the public sector include cases detected at the community level and in the private sector.

<sup>6</sup> Figures include all imported or non-human malaria cases, none of them being indigenous malaria cases.

























## ANNEX 3 – H. REPORTED MALARIA CASES BY METHOD OF CONFIRMATION, 2010–2019

WHO region Country/area	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>REGIONAL SUMMARY (presumed and confirmed malaria cases)</b>										
African	116 141 463	110 102 595	119 380 425	132 833 076	135 963 214	139 545 767	149 056 280	154 864 155	159 845 148	188 651 462
Americas	677 230	493 823	469 385	439 700	392 491	450 101	568 941	773 486	763 232	723 028
Eastern Mediterranean	6 368 813	5 952 130	5 835 463	4 944 058	5 331 046	5 385 450	3 575 561	4 269 075	5 202 797	4 486 311
European	229	275	766	550	515	454	426	439	18	0
South-East Asia	3 114 651	2 503 527	2 110 897	1 659 380	1 645 583	1 627 837	1 454 264	1 215 771	745 313	671 606
Western Pacific	1 643 835	1 367 136	1 085 937	1 294 237	802 833	704 569	873 965	1 014 763	1 075 966	788 727
<b>Total</b>	<b>127 946 221</b>	<b>120 419 486</b>	<b>128 882 873</b>	<b>141 171 001</b>	<b>144 135 682</b>	<b>147 730 660</b>	<b>155 749 604</b>	<b>162 373 650</b>	<b>167 627 726</b>	<b>195 321 141</b>

Data as of 17 November 2020

RDT: rapid diagnostic test; WHO: World Health Organization.

“–” refers to not applicable or data not available.

\* Excluding data from Khartoum.

\*\* Microscopy and RDT examined and positives results are combined and cannot be disaggregated.

\*\*\* Case investigation is less than 100%.

<sup>1</sup> Certified malaria free countries are included in this listing for historical purposes.

<sup>2</sup> Figures reported for the public sector include cases detected at the community level.

<sup>3</sup> Figures reported for the public sector include cases detected at the community level and in the private sector.

<sup>4</sup> Figures reported for the public sector include cases detected in the private sector.

<sup>5</sup> In May 2013, South Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [https://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

<sup>6</sup> Figures include all imported or non-human malaria cases, none of them being indigenous malaria cases.

<sup>7</sup> There are no indigenous cases.

Note: Imported cases also include introduced cases.



**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>											
Algeria <sup>1</sup>	Indigenous cases	1	1	55	8	0	0	0	0	0	0
	Total <i>P. falciparum</i>	7	4	-	-	5	-	-	-	-	0
	Total <i>P. vivax</i>	4	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	1	-	-	-	13	-	-	-	-	-
	Imported cases	394	187	825	587	260	727	420	446	1 241	1 014
Angola	Indigenous cases	1 682 870	1 632 282	1 496 834	1 999 868	2 298 979	2 769 305	3 794 253	3 874 892	5 150 575	7 054 978
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	7 054 978
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Benin	Indigenous cases	-	68 745	705 839	1 090 602	1 309 238	1 721 626	1 610 790	1 933 912	1 730 005	3 299 144
	Total <i>P. falciparum</i>	-	68 745	-	-	1 044 235	1 268 347	1 324 576	1 696 777	1 768 450	3 299 144
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	294 518
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Botswana	Indigenous cases	1 046	432	193	456	1 346	326	716	1 900	585	169
	Total <i>P. falciparum</i>	1 046	432	193	456	1 346	326	703	1 891	585	169
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	2	-	0
	Total mixed cases	-	-	-	-	-	-	12	9	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	30	30	48	64	62	51	103
Burkina Faso	Indigenous cases	804 539	428 113	3 858 046	3 769 051	5 428 655	7 015 446	9 779 154	10 225 459	10 278 970	5 877 426
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	5 877 426
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Burundi	Indigenous cases	1 763 447	1 571 874	2 151 076	4 141 387	4 585 273	5 159 706	8 274 062	7 670 177	4 966 511	9 959 533
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	-
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Cabo Verde	Indigenous cases	47	7	1	22	26	7	48	423	2	0
	Total <i>P. falciparum</i>	47	7	-	-	26	7	48	423	2	1
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	29	35	24	20	21	27	23	18	39
Cameroon	Indigenous cases	-	0	-	0	-	1 162 784	1 675 264	1 191 257	1 249 705	2 318 830
	Total <i>P. falciparum</i>	-	-	-	-	-	592 351	810 367	1 191 257	1 249 705	2 318 830
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Central African Republic	Indigenous cases	-	-	46 759	116 300	295 088	598 833	1 032 764	383 309	972 119	4 063 727
	Total <i>P. falciparum</i>	-	-	-	-	295 088	598 833	1 032 764	383 309	972 119	4 063 727
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-

**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>											
Chad	Indigenous cases	200 448	181 126	7 710	754 565	914 032	787 046	1 294 768	1 962 372	1 364 706	1 632 529
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	1 364 706	1 632 529
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Comoros	Indigenous cases	36 538	24 856	49 840	53 156	2 203	1 300	1 066	2 274	15 613	17 599
	Total <i>P. falciparum</i>	33 791	21 387	43 681	45 669	2 203	1 300	1 066	2 274	15 613	17 599
	Total <i>P. vivax</i>	528	334	637	72	-	-	-	-	-	-
	Total mixed cases	-	-	-	363	-	-	-	-	-	-
	Total other species	880	557	1 189	363	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	98
Congo	Indigenous cases	-	37 744	120 319	43 232	66 323	51 529	171 847	127 939	116 903	117 837
	Total <i>P. falciparum</i>	-	37 744	120 319	43 232	66 323	51 529	171 847	127 939	116 903	117 837
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Côte d'Ivoire	Indigenous cases	62 726	29 976	1 140 627	2 506 953	3 712 831	3 375 904	3 471 024	3 274 683	4 766 477	5 935 178
	Total <i>P. falciparum</i>	-	-	-	2 506 953	3 712 831	3 375 904	3 471 024	3 274 683	4 766 477	5 935 178
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Democratic Republic of the Congo	Indigenous cases	2 417 780	4 561 981	4 791 598	6 715 223	9 968 983	11 627 473	15 330 841	15 176 927	16 972 207	21 608 681
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	21 608 681
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Equatorial Guinea	Indigenous cases	53 813	22 466	15 169	13 129	20 417	15 142	147 714	15 725	8 962	25 904
	Total <i>P. falciparum</i>	53 813	22 466	15 169	13 129	17 452	-	-	-	-	239
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	15 790
	Total mixed cases	-	-	-	-	-	-	-	-	-	2 036
	Total other species	-	-	-	-	-	-	-	-	-	0
	Imported cases	-	-	-	-	-	-	-	-	-	-
Eritrea	Indigenous cases	35 982	34 848	21 815	21 317	50 534	28 036	24 251	54 005	46 440	93 878
	Total <i>P. falciparum</i>	9 785	10 263	12 121	12 482	23 787	14 510	20 704	21 849	16 553	75 568
	Total <i>P. vivax</i>	3 989	4 932	9 204	7 361	6 780	4 780	2 999	9 185	6 108	15 790
	Total mixed cases	63	94	346	1 391	166	70	543	429	268	2 036
	Total other species	57	19	346	83	35	12	5	23	26	0
	Imported cases	-	-	-	-	-	-	-	-	-	-
Eswatini	Indigenous cases	268	549	562	962	711	157	350	724	308	239
	Total <i>P. falciparum</i>	87	189	192	253	389	157	209	724	308	239
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	1	-	-	-	-	-	-
	Imported cases***	-	170	153	234	322	282	221	403	348	338
Ethiopia	Indigenous cases	1 158 197	1 480 306	1 692 578	2 645 454	2 118 815	1 867 059	1 718 504	1 530 739	962 087	904 495
	Total <i>P. falciparum</i>	732 776	814 547	946 595	1 687 163	1 250 110	1 188 627	1 142 235	1 059 847	859 675	738 155
	Total <i>P. vivax</i>	390 252	665 813	745 983	958 291	868 705	678 432	576 269	470 892	102 412	166 340
	Total mixed cases	73 801	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-

**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>											
Gabon	Indigenous cases	13 936	-	19 753	28 982	31 900	23 867	23 915	35 244	111 719	52 811
	Total <i>P. falciparum</i>	2 157	-	-	26 432	26 117	-	23 915	35 244	111 719	52 811
	Total <i>P. vivax</i>	720	-	-	-	-	-	-	-	-	0
	Total mixed cases	55	-	-	-	-	-	-	-	-	0
	Total other species	2 015	-	-	-	1 570	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Gambia	Indigenous cases	116 353	261 967	300 363	240 792	166 229	240 382	153 774	69 931	87 448	53 386
	Total <i>P. falciparum</i>	64 108	190 379	271 038	240 792	99 976	240 382	153 685	69 931	87 448	53 386
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Ghana	Indigenous cases	1 071 637	1 041 260	3 755 166	1 639 451	3 415 912	4 319 919	4 505 442	4 348 694	4 931 448	6 115 267
	Total <i>P. falciparum</i>	926 447	593 518	3 755 166	1 629 198	3 415 912	4 319 919	4 421 788	4 266 541	4 808 163	6 075 297
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	83 654	82 153	-	28 952
	Total other species	102 937	31 238	-	-	-	-	-	-	-	11 018
	Imported cases	-	-	-	-	-	-	-	-	-	-
Guinea	Indigenous cases	20 936	95 574	317 200	211 257	660 207	810 979	992 146	1 335 323	1 214 996	2 143 225
	Total <i>P. falciparum</i>	20 936	5 450	191 421	63 353	660 207	810 979	992 146	1 335 323	1 214 996	2 143 225
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Guinea-Bissau	Indigenous cases	50 391	71 982	50 381	54 584	93 431	142 309	150 903	143 554	125 511	160 907
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	89 784	125 511	160 907
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Kenya	Indigenous cases	898 531	1 002 805	1 453 471	2 375 129	2 851 555	2 041 277	3 064 796	3 607 026	2 318 090	4 656 702
	Total <i>P. falciparum</i>	898 531	1 002 805	1 453 471	2 335 286	2 808 931	1 499 027	2 783 846	3 215 116	1 521 566	4 656 702
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Liberia	Indigenous cases	922 173	1 915 762	1 407 455	1 244 220	864 204	931 086	1 191 137	1 070 113	-	915 845
	Total <i>P. falciparum</i>	212 927	577 641	1 407 455	1 244 220	864 204	2 086 600	1 191 137	1 760 966	-	915 845
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Madagascar	Indigenous cases	202 450	224 498	359 420	385 598	377 963	744 103	475 333	800 661	972 790	970 828
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	-
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	712	1 167	1 212	-	-	-
Malawi	Indigenous cases	-	304 499	1 564 984	1 280 892	2 905 310	3 661 238	4 827 373	4 901 344	5 865 476	5 184 107
	Total <i>P. falciparum</i>	-	-	-	-	2 905 310	3 585 315	4 730 835	4 901 344	5 830 741	5 184 107
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-

**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>											
Mali	Indigenous cases	239 787	307 035	968 136	1 506 940	2 220 956	2 454 508	2 311 098	2 277 218	2 562 921	3 221 535
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	3 221 535
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Mauritania	Indigenous cases	6 367	5 991	9 037	13 085	15 835	22 631	23 042	20 105	30 609	14 869
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	-
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Mayotte	Indigenous cases	0	45	27	0	1	0	0	0	3	0
	Total <i>P. falciparum</i>	-	38	21	-	1	-	-	-	-	0
	Total <i>P. vivax</i>	-	2	2	-	-	-	-	-	-	0
	Total mixed cases	-	-	4	-	-	-	-	-	-	0
	Total other species	-	-	2	-	-	-	-	-	-	-
	Imported cases***	224	51	47	71	14	10	10	10	10	44
Mozambique	Indigenous cases	1 522 577	1 756 874	1 853 276	3 282 172	7 407 175	8 222 814	9 690 873	9 892 601	10 304 472	11 734 926
	Total <i>P. falciparum</i>	878 009	663 132	927 841	2 998 874	7 117 648	7 718 782	8 520 376	8 921 081	9 292 928	11 734 926
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Namibia	Indigenous cases	556	1 860	194	4 911	15 914	12 168	25 198	54 268	36 451	2 340
	Total <i>P. falciparum</i>	556	335	194	136	15 914	12 050	329	364	280	2 340
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	6
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	16
	Imported cases***	-	-	-	-	-	2 888	3 980	11 874	4 021	1 064
Niger	Indigenous cases	620 058	780 876	2 239 858	2 353 422	1 953 309	2 272 000	4 148 167	2 638 580	3 046 450	3 771 451
	Total <i>P. falciparum</i>	601 455	757 449	817 072	1 426 696	3 828 486	2 267 867	3 961 178	2 638 580	3 046 450	3 748 155
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	17 123	21 370	22 399	46 068	78 102	-	-	-	-	0
	Total other species	17 123	21 370	25 270	5 102	39 066	4 133	186 989	-	-	23 296
	Imported cases	-	-	-	-	-	-	-	-	-	-
Nigeria	Indigenous cases	551 187	-	-	-	7 826 954	7 100 032	13 598 282	13 087 878	14 548 024	17 774 018
	Total <i>P. falciparum</i>	523 513	-	-	-	-	-	-	-	-	-
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Rwanda	Indigenous cases	669 322	273 293	563 852	1 040 557	1 719 904	2 694 566	4 725 577	5 940 533	4 231 883	3 612 822
	Total <i>P. falciparum</i>	638 669	208 858	483 470	962 618	1 623 176	-	-	2 927 780	1 657 793	1 306 846
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	2 305 976
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Sao Tome and Principe	Indigenous cases	2 740	8 442	10 701	9 243	1 754	2 058	2 238	2 239	2 937	2 447
	Total <i>P. falciparum</i>	2 219	6 363	10 700	9 242	1 754	2 055	2 234	2 239	2 937	2 447
	Total <i>P. vivax</i>	14	4	1	1	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	6	-	-	-	1	-	-	-	-
	Imported cases***	-	-	-	-	-	2	4	2	3	10

**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>											
Senegal	Indigenous cases	330 331	274 119	280 241	366 687	268 912	492 253	349 540	395 706	530 652	354 663
	Total <i>P. falciparum</i>	343 670	277 326	281 080	345 889	265 624	491 901	347 635	395 706	530 652	354 663
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	1	-	-	-	-	-	-	0
	Total other species	-	-	1	-	-	-	-	-	-	-
	Imported cases***	-	-	-	-	-	352	1 905	0	292	45
Sierra Leone	Indigenous cases	934 028	638 859	1 537 322	1 701 958	1 374 476	1 483 376	1 775 306	1 651 236	1 733 831	2 407 505
	Total <i>P. falciparum</i>	218 473	25 511	1 537 322	1 701 958	1 374 476	1 483 376	1 775 306	1 651 236	1 733 831	2 407 505
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	0	-
South Africa	Indigenous cases	8 060	9 866	5 629	8 645	11 705	4 357	4 323	28 295	9 540	3 096
	Total <i>P. falciparum</i>	2 181	6 906	3 109	8 645	11 563	554	3 104	22 061	9 540	3 096
	Total <i>P. vivax</i>	-	14	5	-	-	-	-	-	-	0
	Total mixed cases	12	-	-	-	-	1	-	-	-	0
	Total other species	5	15	7	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	3 568	3 075	6 234	5 742	8 890
South Sudan <sup>2</sup>	Indigenous cases	900 283	112 024	225 371	262 520	71 377	24 371	7 619	1 488 005	3 242	1 903 742
	Total <i>P. falciparum</i>	-	112 024	-	-	-	-	7 619	1 488 005	3 242	1 902 505
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Togo	Indigenous cases	1 021 854	519 450	909 129	965 334	1 524 339	1 632 594	1 746 234	1 756 582	2 002 877	2 406 091
	Total <i>P. falciparum</i>	224 080	237 282	260 526	272 847	1 130 234	1 113 910	1 174 116	1 208 957	1 090 110	2 381 123
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	8	-	-	-	-	-	0
	Total other species	7	23	9	8	17	17	9 149	77	224	1 005
	Imported cases	-	-	-	-	-	-	-	-	-	-
Uganda	Indigenous cases	1 628 595	231 873	2 662 258	1 502 362	3 631 939	7 137 662	9 385 132	11 667 831	5 759 174	13 982 362
	Total <i>P. falciparum</i>	1 565 348	231 873	2 662 258	1 502 362	3 631 939	7 137 662	9 385 132	11 700 000	5 759 174	13 982 362
	Total <i>P. vivax</i>	15 812	-	-	-	-	-	-	-	-	0
	Total mixed cases	47 435	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
United Republic of Tanzania	Indigenous cases	1 278 998	2 150 761	1 986 955	1 551 777	106 991	412 608	5 188 505	5 354 486	6 051 914	5 908 168
	Total <i>P. falciparum</i>	2 338	4 489	2 730	1 475	227	412 433	-	1 733	486	1 338
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	212 837	69 511	106 764	175	-	1 606	1 020	-
	Total other species	-	-	-	-	106 609	-	-	10	26	-
	Imported cases***	-	-	-	1 438	3 166	5 100	-	-	1 754	3 286
Mainland	Indigenous cases	1 276 660	2 146 272	1 984 024	1 550 250	106 609	411 741	5 188 505	5 351 137	6 050 382	5 906 621
	Total <i>P. falciparum</i>	-	-	-	-	-	411 741	-	-	-	-
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	212 636	69 459	106 609	-	-	-	-	-
	Total other species	-	-	-	-	106 609	-	-	-	-	-
	Imported cases	-	-	-	719	1 583	2 550	-	-	-	-
Zanzibar	Indigenous cases	2 338	4 489	2 931	1 527	382	867	0	3 349	1 532	1 547
	Total <i>P. falciparum</i>	2 338	4 489	2 730	1 475	227	692	-	1 733	486	1 338
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	201	52	155	175	-	1 606	1 020	-
	Total other species	-	-	-	-	-	-	-	10	26	-
	Imported cases***	-	-	-	719	1 583	2 550	-	-	1 754	3 286

## ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>											
Zambia	Indigenous cases	-	-	-	-	4 077 547	4 184 661	4 851 319	5 505 639	5 039 679	5 147 350
	Total <i>P. falciparum</i>	-	-	-	-	4 077 547	4 184 661	4 851 319	5 505 639	5 039 679	5 147 350
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Zimbabwe	Indigenous cases	249 379	319 935	276 963	422 633	548 276	482 379	314 003	468 276	255 388	308 173
	Total <i>P. falciparum</i>	249 379	319 935	276 963	422 633	535 931	391 651	279 988	315 624	183 755	308 173
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	9 299
	Total mixed cases	-	-	-	-	-	-	-	-	-	5
	Total other species	-	-	-	-	-	-	-	-	-	8
	Imported cases***	-	-	-	-	-	180	358	768	672	-
<b>AMERICAS</b>											
Argentina <sup>1</sup>	Indigenous cases	14	0	0	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	-
	Total <i>P. vivax</i>	26	-	-	-	-	-	-	-	-	-
	Total mixed cases	-	-	-	-	-	-	-	-	-	1
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	46	18	4	4	4	8	5	15	23	-
Belize	Indigenous cases	150	72	33	20	19	9	4	7	3	0
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	1
	Total <i>P. vivax</i>	149	72	33	20	19	9	4	5	2	0
	Total mixed cases	1	-	-	-	-	-	-	-	2	-
	Total other species	-	-	-	-	-	-	-	-	-	0
	Imported cases	-	7	4	4	0	4	1	2	4	2
Bolivia (Plurinational State of)	Indigenous cases	13 769	7 143	7 415	7 301	7 401	6 874	5 542	4 572	5 342	9 338
	Total <i>P. falciparum</i>	1 557	526	385	959	325	77	4	-	-	26
	Total <i>P. vivax</i>	13 694	7 635	8 141	6 346	7 060	6 785	5 535	4 572	5 342	9 299
	Total mixed cases	35	17	11	37	16	12	3	-	-	5
	Total other species	-	-	-	-	-	-	-	-	-	8
	Imported cases	-	-	-	-	-	33	11	15	12	19
Brazil	Indigenous cases	334 483	267 146	242 758	168 862	139 272	138 229	124 178	189 503	187 693	153 296
	Total <i>P. falciparum</i>	47 406	32 100	32 437	25 928	21 297	14 762	13 160	18 614	17 852	15 138
	Total <i>P. vivax</i>	283 435	231 368	203 018	137 887	115 809	122 746	110 341	169 887	168 499	136 949
	Total mixed cases	3 642	3 606	7 722	5 015	2 139	683	669	1 032	1 331	1 189
	Total other species	183	143	0	32	38	38	8	26	11	20
	Imported cases	-	-	-	8 905	4 847	4 915	5 087	4 867	6 819	4 158
Colombia	Indigenous cases	117 589	60 105	60 179	51 696	40 768	47 616	82 609	52 805	61 195	78 109
	Total <i>P. falciparum</i>	32 900	14 650	17 612	17 650	20 067	27 875	47 232	29 558	29 953	40 074
	Total <i>P. vivax</i>	83 255	44 701	44 283	33 345	20 129	19 002	32 635	22 132	30 063	37 197
	Total mixed cases	1 434	754	672	690	567	739	2 742	1 115	1 179	838
	Total other species	48	16	9	11	5	-	-	-	-	9
	Imported cases	-	-	-	-	-	7 785	618	1 297	1 948	2 306
Costa Rica	Indigenous cases	110	10	6	0	0	0	4	12	70	95
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	9	8
	Total <i>P. vivax</i>	110	11	4	-	-	-	4	12	61	92
	Total mixed cases	-	-	1	-	-	-	-	-	-	5
	Total other species	-	-	2	-	-	-	-	-	-	10
	Imported cases	4	6	1	4	5	8	9	13	38	45
Dominican Republic	Indigenous cases	2 482	1 616	952	579	459	631	690	398	608	1 291
	Total <i>P. falciparum</i>	2 480	1 614	950	474	459	631	690	341	561	1 291
	Total <i>P. vivax</i>	2	2	2	-	-	-	-	-	29	0
	Total mixed cases	-	-	-	-	-	-	-	-	2	0
	Total other species	-	-	-	-	-	-	-	-	-	0
	Imported cases	-	-	-	105	37	30	65	57	50	23

**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AMERICAS</b>											
Ecuador	Indigenous cases	1 871	1 219	544	368	242	618	1 191	1 275	1 653	1 803
	Total <i>P. falciparum</i>	258	290	78	160	40	184	403	309	149	211
	Total <i>P. vivax</i>	1 630	929	466	208	202	434	788	963	1 504	1 592
	Total mixed cases	-	-	-	-	-	-	-	3	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	14	14	10	-	59	233	105	153	106
El Salvador	Indigenous cases	19	9	13	6	6	2	12	0	0	0
	Total <i>P. falciparum</i>	-	1	-	-	-	-	-	-	-	0
	Total <i>P. vivax</i>	17	8	-	6	6	2	12	-	-	0
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	0
	Imported cases	7	6	6	1	2	7	1	3	2	3
French Guiana	Indigenous cases	1 608	1 209	900	877	448	374	217	554	546	176
	Total <i>P. falciparum</i>	987	584	382	304	136	32	29	33	-	17
	Total <i>P. vivax</i>	476	339	257	220	129	203	99	409	-	193
	Total mixed cases	561	496	381	348	182	3	3	5	-	0
	Total other species	5	5	2	345	1	-	-	-	-	2
	Imported cases	-	-	-	-	-	60	41	43	-	36
Guatemala	Indigenous cases	7 384	6 817	5 346	6 214	5 685	6 836	4 853	3 743	3 021	2 069
	Total <i>P. falciparum</i>	30	64	54	101	24	43	4	3	-	0
	Total <i>P. vivax</i>	7 163	6 707	5 278	6 062	5 593	5 487	4 849	3 739	4 766	2 069
	Total mixed cases	5	3	14	51	67	8	-	1	-	0
	Total other species	-	-	-	-	-	-	-	0	0	0
	Imported cases	-	-	-	-	1	2	1	2	3	3
Guyana	Indigenous cases	22 935	29 471	31 610	31 479	12 354	9 984	1 461	13 936	17 038	18 826
	Total <i>P. falciparum</i>	11 244	15 945	16 722	13 655	3 943	3 219	4 046	5 141	6 032	5 737
	Total <i>P. vivax</i>	8 402	9 066	11 244	13 953	7 173	6 002	6 923	7 645	9 853	11 940
	Total mixed cases	3 157	4 364	3 607	3 770	1 197	731	930	1 078	1 089	381
	Total other species	132	96	83	101	41	32	57	72	64	38
	Imported cases	-	-	-	-	-	-	411	-	-	0
Haiti	Indigenous cases	84 153	34 350	25 423	20 957	17 696	17 926	22 718	16 733	7 075	10 687
	Total <i>P. falciparum</i>	84 153	32 969	25 423	20 957	17 696	17 583	21 998	18 843	9 112	10 687
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	319
	Total mixed cases	-	-	-	-	-	-	-	-	-	123
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Honduras	Indigenous cases	9 744	7 510	6 437	5 428	3 378	3 575	4 096	1 286	882	330
	Total <i>P. falciparum</i>	866	585	560	1 113	562	904	1 309	128	93	11
	Total <i>P. vivax</i>	8 759	7 044	5 865	4 269	2 881	2 642	2 745	1 149	763	319
	Total mixed cases	120	34	24	46	37	29	40	-	2	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases***	-	-	-	-	2	0	3	10	21	61
Mexico	Indigenous cases	1 226	1 124	833	495	656	517	551	736	803	618
	Total <i>P. falciparum</i>	-	-	-	-	-	-	-	-	-	1
	Total <i>P. vivax</i>	1 226	1 124	833	495	656	517	551	736	803	618
	Total mixed cases	-	-	-	-	-	-	-	-	-	123
	Total other species	-	-	-	-	-	-	-	-	0	0
	Imported cases	7	6	9	4	10	34	45	29	23	22
Nicaragua	Indigenous cases	692	925	1 235	1 162	1 142	2 279	6 272	10 949	15 917	13 200
	Total <i>P. falciparum</i>	154	150	236	208	155	338	1 285	1 836	1 319	2 398
	Total <i>P. vivax</i>	538	775	999	954	985	1 937	4 965	9 080	14 553	10 679
	Total mixed cases	-	-	-	-	2	4	22	33	45	123
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	34	21	29	12	3	17	26

### ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AMERICAS</b>											
Panama	Indigenous cases	418	354	844	705	864	546	769	649	684	1 554
	Total <i>P. falciparum</i>	20	1	1	-	-	-	21	1	-	25
	Total <i>P. vivax</i>	398	353	843	696	864	546	748	648	684	1 197
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	375
	Imported cases	-	-	-	9	10	16	42	40	31	43
Paraguay <sup>1</sup>	Indigenous cases	18	1	0	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	-	-	-	-	-	0	0	0	0	-
	Total <i>P. vivax</i>	18	1	-	-	-	0	0	0	0	-
	Total mixed cases	-	-	-	-	-	0	0	0	0	-
	Total other species	-	-	-	1	-	0	0	0	0	-
	Imported cases	9	9	15	11	8	8	10	5	-	-
Peru	Indigenous cases	31 545	25 005	31 436	48 719	65 252	61 865	56 623	55 367	45 443	24 324
	Total <i>P. falciparum</i>	2 291	2 929	3 399	7 890	10 416	12 569	15 319	13 173	9 438	4 724
	Total <i>P. vivax</i>	29 169	21 984	28 030	40 829	54 819	49 287	41 287	42 044	36 004	19 600
	Total mixed cases	83	89	102	213	-	-	-	148	0	0
	Total other species	3	3	7	11	17	9	17	2	0	0
	Imported cases	-	-	-	-	0	0	0	-	176	159
Suriname	Indigenous cases	1 712	771	356	729	401	81	76	40	29	95
	Total <i>P. falciparum</i>	638	310	115	322	165	17	6	1	5	0
	Total <i>P. vivax</i>	817	382	167	322	78	61	69	17	23	95
	Total mixed cases	83	21	11	85	158	3	1	1	1	0
	Total other species	36	17	2	-	-	0	0	0	0	0
	Imported cases	-	-	-	204	-	274	251	414	198	111
Venezuela (Bolivarian Republic of)	Indigenous cases	45 155	45 824	52 803	78 643	90 708	136 402	240 613	411 586	404 924	398 285
	Total <i>P. falciparum</i>	10 629	9 724	10 978	22 421	21 074	24 018	46 046	68 362	80 087	64 307
	Total <i>P. vivax</i>	32 710	34 651	39 478	49 691	62 850	100 880	178 187	314 406	342 692	308 133
	Total mixed cases	-	-	-	4 808	6 769	11 491	14 531	25 849	28 128	25 846
	Total other species	60	6	23	46	15	13	25	28	9	0
	Imported cases	-	-	-	1 677	1 210	1 594	1 948	2 941	2 125	1 848
<b>EASTERN MEDITERRANEAN</b>											
Afghanistan	Indigenous cases	69 798	77 549	54 840	52 965	106 478	119 859	241 233	311 598	248 689	173 860
	Total <i>P. falciparum</i>	6 142	5 581	1 231	1 877	3 000	4 004	6 369	6 907	6 437	2 701
	Total <i>P. vivax</i>	63 255	71 968	53 609	43 369	58 362	82 891	132 407	154 468	166 583	170 747
	Total mixed cases	-	-	-	-	1 566	-	311	403	473	232
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Djibouti	Indigenous cases	1 010	-	25	1 684	9 439	9 473	13 804	14 671	25 319	49 402
	Total <i>P. falciparum</i>	1 010	-	20	-	-	-	11 781	9 290	16 130	36 025
	Total <i>P. vivax</i>	-	-	-	-	-	-	2 041	5 381	9 189	13 377
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Iran (Islamic Republic of)	Indigenous cases	1 847	1 632	756	479	358	167	81	60	0	0
	Total <i>P. falciparum</i>	166	152	44	72	21	8	7	2	0	6
	Total <i>P. vivax</i>	1 656	1 502	711	426	351	157	87	55	0	78
	Total mixed cases	25	56	32	22	4	1	1	-	0	1
	Total other species	-	-	-	1	-	-	-	-	0	0
	Imported cases	1 184	1 529	842	853	867	632	612	868	602	1 105
Pakistan	Indigenous cases	240 591	334 589	326 211	274 648	264 867	203 859	323 510	368 519	374 511	413 533
	Total <i>P. falciparum</i>	73 857	73 925	95 095	46 067	33 391	30 075	42 011	54 467	55 832	87 169
	Total <i>P. vivax</i>	143 136	205 879	228 215	283 661	232 332	163 872	257 962	300 623	314 385	323 355
	Total mixed cases	-	-	2 901	10 506	556	8 066	24 493	14 787	4 489	2 510
	Total other species	-	-	-	-	-	-	-	-	0	-
	Imported cases	-	-	-	-	-	-	-	-	-	-



**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>EASTERN MEDITERRANEAN</b>											
Saudi Arabia	Indigenous cases	29	69	82	34	30	83	272	177	61	38
	Total <i>P. falciparum</i>	29	69	82	34	51	83	270	172	57	122
	Total <i>P. vivax</i>	-	-	-	-	-	0	2	5	4	1
	Total mixed cases	-	-	-	-	-	0	0	0	-	0
	Total other species	-	-	-	6	-	0	0	0	0	0
	Imported cases	1 912	2 719	3 324	2 479	2 254	2 537	5 110	2 974	2 517	2 029
Somalia	Indigenous cases	24 553	3 351	35 712	8 944	11 001	20 953	35 628	35 138	31 021	39 687
	Total <i>P. falciparum</i>	5 629	189	-	-	-	-	-	-	-	36 304
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	-	3 383
	Total mixed cases	-	-	-	-	-	-	-	-	-	-
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Sudan	Indigenous cases	720 557	506 806	526 931	592 383	1 068 506	586 827	566 015	720 879	1 606 833	1 752 011
	Total <i>P. falciparum</i>	-	-	-	-	-	-	333 009	580 145	1 286 915	1 363 507
	Total <i>P. vivax</i>	-	-	-	-	-	-	82 175	58 335	143 314	194 904
	Total mixed cases	-	-	-	-	-	-	32 557	82 399	187 270	193 600
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Yemen	Indigenous cases	106 697	90 954	112 359	102 778	86 707	76 259	98 701	114 004	117 652	165 899
	Total <i>P. falciparum</i>	77 271	59 689	109 504	102 369	86 428	75 898	45 469	109 849	112 823	163 941
	Total <i>P. vivax</i>	966	478	398	408	267	334	347	1 833	970	1 802
	Total mixed cases	30	7	2	-	12	27	70	2 322	63	114
	Total other species	2	33	4	-	-	-	-	-	69	42
	Imported cases	-	-	-	-	-	-	-	-	-	-
<b>EUROPEAN</b>											
Armenia <sup>1</sup>	Indigenous cases	0	0	0	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	0	0	0	0	0	0	0	0	3	-
	Total <i>P. vivax</i>	0	0	0	0	0	0	0	0	1	-
	Total mixed cases	0	-	-	-	-	0	0	0	0	-
	Total other species	0	0	0	0	0	0	0	0	2	-
	Imported cases	1	0	4	0	1	1	1	2	6	-
Azerbaijan	Indigenous cases	50	4	3	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	0	0	0	0	0	0	0	0	2	0
	Total <i>P. vivax</i>	50	4	3	0	0	0	0	0	0	0
	Total mixed cases	0	0	0	0	0	0	0	0	0	0
	Total other species	0	0	0	0	0	0	0	0	0	0
	Imported cases	2	4	1	4	2	1	1	1	2	0
Georgia	Indigenous cases	0	0	0	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	0	0	0	0	0	0	7	7	7	-
	Total <i>P. vivax</i>	0	1	1	0	0	0	0	1	2	-
	Total mixed cases	0	0	0	0	0	0	0	0	0	-
	Total other species	0	0	0	0	0	0	0	0	0	-
	Imported cases	0	5	4	7	5	5	7	8	9	-
Kazakhstan	Indigenous cases	-	-	-	-	-	0	0	0	0	0
	Total <i>P. falciparum</i>	-	-	-	-	-	0	3	1	2	2
	Total <i>P. vivax</i>	-	-	-	-	-	0	0	1	1	0
	Total mixed cases	-	-	-	-	-	0	1	1	1	0
	Total other species	-	-	-	-	-	0	0	0	0	0
	Imported cases	3	4	2	2	1	2	4	3	4	2
Tajikistan	Indigenous cases	111	65	18	3	2	0	0	0	0	0
	Total <i>P. falciparum</i>	0	0	0	0	0	0	0	2	0	-
	Total <i>P. vivax</i>	111	53	18	3	2	0	1	1	1	-
	Total mixed cases	0	0	0	0	0	0	0	0	0	-
	Total other species	0	0	0	0	0	0	0	0	0	-
	Imported cases	1	13	15	7	5	4	1	3	1	-

## ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>EUROPEAN</b>											
Turkey	Indigenous cases	0	0	0	0	0	0	0	0	-	0
	Total <i>P. falciparum</i>	-	-	-	-	-	0	0	0	-	-
	Total <i>P. vivax</i>	9	4	-	34	-	0	0	0	-	-
	Total mixed cases	-	-	-	-	-	0	0	0	-	-
	Total other species	-	-	-	-	-	0	0	0	-	-
	Imported cases	81	128	376	251	249	221	208	214	-	-
Turkmenistan <sup>1</sup>	Indigenous cases	0	-	-	-	0	0	0	0	0	0
	Total <i>P. falciparum</i>	0	0	0	0	0	0	0	0	0	-
	Total <i>P. vivax</i>	0	0	0	0	0	0	0	0	0	-
	Total mixed cases	0	-	-	-	-	0	0	0	0	-
	Total other species	0	0	0	0	0	0	0	0	0	-
	Imported cases	0	0	0	0	0	0	0	0	0	-
Uzbekistan <sup>1</sup>	Indigenous cases	3	0	0	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	0	0	0	0	0	0	0	0	0	-
	Total <i>P. vivax</i>	3	0	0	0	0	0	0	0	0	-
	Total mixed cases	0	0	0	0	0	0	0	0	0	-
	Total other species	0	0	0	0	0	0	0	0	0	-
	Imported cases	2	1	1	3	1	0	0	0	0	-
<b>SOUTH-EAST ASIA</b>											
Bangladesh	Indigenous cases	55 873	51 773	29 518	21 454	47 101	39 719	27 737	29 026	10 467	17 219
	Total <i>P. falciparum</i>	52 012	49 084	9 428	3 597	8 981	5 279	3 460	4 210	1 571	14 752
	Total <i>P. vivax</i>	3 824	2 579	396	262	489	477	418	520	277	2 126
	Total mixed cases	37	110	36	5	746	723	800	163	30	338
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases***	-	-	-	-	-	129	109	19	41	6
Bhutan	Indigenous cases	436	194	82	15	19	34	15	11	6	2
	Total <i>P. falciparum</i>	140	87	-	6	11	13	1	0	1	4
	Total <i>P. vivax</i>	261	92	-	9	8	21	13	11	5	2
	Total mixed cases	35	15	-	0	0	0	1	0	0	0
	Total other species	0	0	-	33	-	0	0	0	0	0
	Imported cases	-	-	0	23	34	70	56	38	34	30
Democratic People's Republic of Korea	Indigenous cases	-	-	-	-	-	-	-	-	0	1 869
	Total <i>P. falciparum</i>	13 520	16 760	21 850	14 407	10 535	7 022	5 033	4 603	3 698	0
	Total <i>P. vivax</i>	-	-	-	-	-	-	-	-	0	1 869
	Total mixed cases	-	-	-	-	-	-	-	-	0	0
	Total other species	-	-	0	0	0	0	0	0	0	0
	Imported cases	13 520	16 760	21 850	14 407	10 535	7 022	5 033	4 603	3 698	0
India	Indigenous cases	1 599 986	1 310 656	1 067 824	881 730	1 102 205	1 169 261	1 087 285	844 558	429 928	338 494
	Total <i>P. falciparum</i>	830 779	662 748	524 370	462 079	720 795	774 627	706 257	525 637	204 733	154 645
	Total <i>P. vivax</i>	765 622	645 652	534 129	417 884	379 659	390 440	375 783	315 028	222 730	181 514
	Total mixed cases	3 585	2 256	0	1 767	1 751	4 194	5 245	3 893	2 465	2 295
	Total other species	3 585	2 256	9 325	1 767	-	0	0	0	0	-
	Imported cases	-	-	-	-	-	-	-	-	-	-
Indonesia	Indigenous cases	465 764	422 447	417 819	343 527	252 027	217 025	217 343	258 519	222 074	205 583
	Total <i>P. falciparum</i>	226 241	205 364	203 114	164 722	125 217	105 525	118 836	144 600	116 046	142 036
	Total <i>P. vivax</i>	205 877	186 730	184 684	156 266	108 268	96 284	82 063	96 142	84 862	86 769
	Total mixed cases	32 185	29 192	28 872	21 146	16 564	13 385	16 471	18 988	18 383	4 120
	Total other species	1 281	1 161	1 149	1 393	1 978	1 831	1 080	1 887	2 794	357
	Imported cases***	-	-	-	-	-	-	-	-	11	61
Myanmar	Indigenous cases	420 808	465 294	481 204	333 871	205 658	182 616	110 146	85 019	76 518	56 411
	Total <i>P. falciparum</i>	388 464	433 146	314 676	222 770	138 311	110 449	62 917	50 730	38 483	23 092
	Total <i>P. vivax</i>	29 944	28 966	135 385	98 860	61 830	65 536	43 748	32 070	36 502	32 940
	Total mixed cases	2 054	3 020	31 040	12 216	5 511	6 624	3 476	2 214	1 530	599
	Total other species	346	162	103	25	6	7	5	5	3	4
	Imported cases	-	-	-	-	-	-	-	-	-	-

**ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019**

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>SOUTH-EAST ASIA</b>											
Nepal	Indigenous cases	3 894	3 414	3 230	1 974	832	591	507	623	619	127
	Total <i>P. falciparum</i>	550	0	20	273	81	162	61	25	1	9
	Total <i>P. vivax</i>	2 349	908	1 480	1 659	663	1 119	433	587	613	118
	Total mixed cases	216	30	0	22	58	35	13	11	5	0
	Total other species	0	0	0	0	0	0	0	0	0	-
	Imported cases***	-	1 069	592	-	667	521	502	670	539	579
Sri Lanka <sup>1</sup>	Indigenous cases	684	124	23	0	0	0	0	0	0	0
	Total <i>P. falciparum</i>	6	3	4	0	0	0	0	0	0	0
	Total <i>P. vivax</i>	669	119	19	0	0	0	0	0	0	0
	Total mixed cases	9	2	0	0	0	0	0	0	0	0
	Total other species	0	0	0	0	0	0	0	0	0	-
	Imported cases	52	51	70	95	49	36	41	57	48	53
Thailand	Indigenous cases	22 949	14 465	29 059	30 218	34 844	23 540	17 800	8 417	6 094	3 538
	Total <i>P. falciparum</i>	9 401	5 710	11 553	14 449	13 743	3 301	1 774	726	441	391
	Total <i>P. vivax</i>	13 401	8 608	17 506	15 573	20 513	810	2 671	2 025	3 000	2 752
	Total mixed cases	147	147	0	196	588	122	109	63	61	25
	Total other species	20	13	3 172	3 084	3 077	19	14	10	21	30
	Imported cases	-	-	-	-	-	9 890	5 724	4 020	1 618	1 342
Timor-Leste	Indigenous cases	48 137	19 739	5 208	1 025	347	80	81	16	0	0
	Total <i>P. falciparum</i>	28 350	14 261	1 962	373	118	33	46	4	0	0
	Total <i>P. vivax</i>	11 432	3 758	2 288	512	139	24	7	3	0	0
	Total mixed cases	468	1 720	0	140	85	23	28	9	0	0
	Total other species	0	0	0	0	0	0	0	0	0	0
	Imported cases	-	-	-	-	-	-	0	13	7	9
<b>WESTERN PACIFIC</b>											
Cambodia	Indigenous cases	96 464	106 905	69 551	44 069	69 178	68 109	43 380	76 804	62 582	32 197
	Total <i>P. falciparum</i>	8 213	7 054	14 896	7 092	8 332	17 830	12 156	20 328	10 525	4 834
	Total <i>P. vivax</i>	4 794	5 155	19 575	11 267	10 356	13 146	9 816	15 207	30 680	26 871
	Total mixed cases	1 270	1 583	4 971	2 418	6 464	2 954	1 520	1 397	1 080	442
	Total other species	-	-	4 971	-	-	2 498	-	-	-	-
	Imported cases	-	-	-	-	-	-	-	-	-	0
China	Indigenous cases	4 990	1 308	244	83	53	36	1	0	0	0
	Total <i>P. falciparum</i>	1 269	57	16	11	6	1	0	0	0	0
	Total <i>P. vivax</i>	3 675	677	179	67	45	24	1	0	0	0
	Total mixed cases	26	1	5	1	0	0	0	0	0	0
	Total other species	20	0	0	0	0	6	0	0	0	0
	Imported cases	2 118	2 819	2 474	4 051	3 026	3 240	3 149	2 672	2 511	2 486
Lao People's Democratic Republic	Indigenous cases	26 723	20 708	61 935	51 471	68 028	50 724	16 541	11 748	9 489	6 687
	Total <i>P. falciparum</i>	4 393	5 770	37 692	24 538	23 928	14 430	4 255	4 550	4 726	2 167
	Total <i>P. vivax</i>	122	442	7 634	12 537	22 625	20 804	6 795	4 590	4 077	4 441
	Total mixed cases	8	-	769	956	1 517	822	173	193	110	69
	Total other species	1	14	769	1	1	-	-	-	-	2 079
	Imported cases***	-	-	-	-	-	0	-	-	0	0
Malaysia	Indigenous cases	5 194	3 954	3 662	2 921	3 147	242	266	85	0	0
	Total <i>P. falciparum</i>	1 344	634	651	422	177	110	69	18	3	20
	Total <i>P. vivax</i>	3 387	1 750	915	385	241	84	192	59	16	47
	Total mixed cases	145	120	48	42	33	22	9	1	0	0
	Total other species	943	1 660	2 187	194	120	26	12	7	2	29
	Imported cases	831	1 044	805	865	766	435	428	423	485	630
Papua New Guinea	Indigenous cases	93 956	84 060	150 195	279 994	281 182	297 787	478 497	478 340	516 249	646 648
	Total <i>P. falciparum</i>	56 735	59 153	58 747	119 469	120 641	118 452	183 686	163 160	174 818	176 063
	Total <i>P. vivax</i>	13 171	9 654	7 108	7 579	78 846	62 228	95 328	113 561	138 006	163 237
	Total mixed cases	4 089	1 164	769	1 279	79 574	115 157	197 711	200 186	201 658	296 139
	Total other species	1 990	632	609	1 279	2 125	1 950	1 772	1 433	1 767	2 079
	Imported cases	-	-	-	-	-	-	-	-	-	-

## ANNEX 3 – I. REPORTED MALARIA CASES BY SPECIES, 2010–2019

WHO region Country/area		2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>WESTERN PACIFIC</b>											
Philippines	Indigenous cases	19 102	9 583	8 086	7 720	6 087	11 410	6 680	6 542	4 346	5 681
	Total <i>P. falciparum</i>	11 824	6 877	4 774	4 968	3 760	834	366	3 258	1 310	5 016
	Total <i>P. vivax</i>	2 885	2 380	2 189	1 357	5 881	882	1 503	551	116	535
	Total mixed cases	214	166	113	83	-	-	-	81	22	80
	Total other species	175	127	57	67	5 320	826	534	60	47	50
	Imported cases***	-	-	-	-	68	18	55	69	79	95
Republic of Korea	Indigenous cases	1 267	505	394	383	557	627	602	436	501	485
	Total <i>P. falciparum</i>	27	20	36	-	-	-	-	-	-	0
	Total <i>P. vivax</i>	1 691	754	473	383	557	627	602	436	501	485
	Total mixed cases	-	-	-	-	-	-	-	-	-	0
	Total other species	-	-	-	-	-	-	-	-	-	-
	Imported cases	54	64	46	50	78	71	67	79	75	74
Solomon Islands	Indigenous cases	39 704	26 657	24 383	25 609	18 404	23 998	54 431	52 463	59 191	72 767
	Total <i>P. falciparum</i>	22 892	14 454	14 748	13 194	9 835	10 478	16 607	15 400	15 771	15 595
	Total <i>P. vivax</i>	12 281	8 665	9 339	11 628	7 845	12 150	33 060	30 169	35 072	47 164
	Total mixed cases	200	83	232	446	724	1 370	4 718	6 881	8 341	9 584
	Total other species	200	-	232	-	-	-	46	13	4	27
	Imported cases	-	-	-	-	-	-	-	-	-	-
Vanuatu	Indigenous cases	9 817	6 179	4 532	2 883	1 314	571	2 252	1 228	632	567
	Total <i>P. falciparum</i>	1 545	770	1 257	1 039	279	150	186	273	42	36
	Total <i>P. vivax</i>	2 265	1 224	1 680	1 342	703	273	1 682	798	590	531
	Total mixed cases	193	81	470	-	-	-	-	-	-	0
	Total other species	10	2	-	-	-	-	-	-	-	0
	Imported cases	-	-	-	-	-	0	0	1	12	9
Viet Nam	Indigenous cases	17 515	16 612	19 638	17 128	15 752	9 331	4 161	4 548	4 813	3 100
	Total <i>P. falciparum</i>	12 763	10 101	11 448	9 532	8 245	4 327	2 323	2 858	2 966	3 100
	Total <i>P. vivax</i>	4 466	5 602	7 220	6 901	7 220	4 756	1 750	1 608	1 751	1 514
	Total mixed cases	286	909	970	695	287	234	73	70	83	31
	Total other species	-	-	-	-	-	14	15	12	13	10
	Imported cases***	-	-	-	-	-	-	-	-	1 681	1 565

*P.*: *Plasmodium*; WHO: World Health Organization.

"-" refers to not applicable or data not available.

\*\*\* Case investigation is less than 100%.

<sup>1</sup> Certified malaria free countries are included in this listing for historical purposes.

<sup>2</sup> In May 2013, Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [https://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

Note: Indigenous cases are reported for countries with elimination programmes and/or with >99% of total confirmed cases investigated. For countries in the WHO Region of the Americas, the number of Total: *P. falciparum*, Total: *P. vivax*, Total: mixed cases and Total: other species are indigenous cases for all years apart from Dominican Republic and Venezuela (Bolivarian Republic of) (2013 onwards), Argentina, Guatemala and Peru (2014 onwards) and Bolivia (Plurinational State of), Honduras and Suriname (2015 onwards). Indigenous cases are reported for Botswana and Eswatini from 2015 onwards. Suspected cases include indigenous and imported cases. For countries with only suspected cases shown; no species breakdown was provided.

Data as of 17 November 2020

## ANNEX 3 – J. REPORTED MALARIA DEATHS, 2010–2019

WHO region Country/area	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AFRICAN</b>										
Algeria <sup>1</sup>	1	0	0	0	0	0	0	0	0	
Angola	8 114	6 909	5 736	7 300	5 714	7 832	15 997	13 967	11 814	18 691
Benin	964	1 753	2 261	2 288	1 869	1 416	1 646	2 182	2 138	2 589
Botswana	8	8	3	7	22	5	3	17	9	14
Burkina Faso	9 024	7 001	7 963	6 294	5 632	5 379	3 974	4 144	4 294	1 060
Burundi	2 677	2 233	2 263	3 411	2 974	3 799	5 853	4 414	2 481	3 316
Cabo Verde	1	1	0	0	1	0	1	2	0	0
Cameroon	4 536	3 808	3 209	4 349	4 398	3 440	2 639	3 195	3 256	4 510
Central African Republic	526	858	1 442	1 026	635	1 763	2 668	3 689	1 292	2 017
Chad	886	1 220	1 359	1 881	1 720	1 572	1 686	2 088	1 948	3 374
Comoros	53	19	17	15	0	1	0	3	8	
Congo		892	623	2 870	271	435	733	229	131	107
Côte d'Ivoire	1 023	1 389	1 534	3 261	4 069	4 413	3 340	3 222	3 133	1 693
Democratic Republic of the Congo	23 476	23 748	21 601	30 918	25 502	39 054	33 997	27 458	18 030	13 072
Equatorial Guinea	30	52	77	66		28	109			15
Eritrea	27	12	30	6	15	12	21	8	5	3
Eswatini	8	1	3	4		5	3	20	2	3
Ethiopia	1 581	936	1 621	358	213	662	510	356	158	213
Gabon	182	74	134	273	159	309	101	218	591	314
Gambia	151	440	289	262	170	167	79	54	60	41
Ghana	3 859	3 259	2 855	2 506	2 200	2 137	1 264	599	428	336
Guinea	735	743	979	108	1 067	846	867	1 174	1 267	1 881
Guinea-Bissau	296	472	370	418	357	477	191	296	244	288
Kenya	26 017	713	785	360	472	15 061	603			858
Liberia	1 422		1 725	1 191	2 288	1 379	1 259	758		601
Madagascar	427	398	552	641	551	841	443	370	927	657
Malawi	8 206	6 674	5 516	3 723	4 490	3 799	4 000	3 613	2 967	2 341
Mali	3 006	2 128	1 894	1 680	2 309	1 544	1 344	1 050	1 001	1 454
Mauritania	60	66	106	46	19	39	315	67		
Mayotte	0	0	0	0	0	0	0			
Mozambique	3 354	3 086	2 818	2 941	3 245	2 467	1 685	1 114	968	734
Namibia	63	36	4	21	61	45	65	104	82	7
Niger	3 929	2 802	2 825	2 209	2 691	2 778	2 226	2 316	3 576	4 449
Nigeria	4 238	3 353	7 734	7 878	6 082					
Rwanda	670	380	459	409	496	516	715	376	341	224
Sao Tome and Principe	14	19	7	11	0	0	1	1	0	0
Senegal	553	472	649	815	500	526	325	284	555	260
Sierra Leone	8 188	3 573	3 611	4 326	2 848	1 107	1 345	1 298	1 949	2 771
South Africa	83	54	72	105	174	110	34	301	69	79
South Sudan <sup>2</sup>	1 053	406	1 321	1 311				3 483	1 191	4 873
Togo	1 507	1 314	1 197	1 361	1 205	1 127	847	995	905	1 275
Uganda	8 431	5 958	6 585	7 277	5 921	6 100	5 635	5 111	3 302	5 027
United Republic of Tanzania	15 915	11 806	7 828	8 528	5 373	6 315	5 046	3 685	2 753	1 171
Mainland	15 867	11 799	7 820	8 526	5 368	6 313	5 045	3 684	2 747	1 163
Zanzibar	48	7	8	2	5	2	1	1	6	8
Zambia	4 834	4 540	3 705	3 548	3 257	2 389	1 827	1 425	1 209	1 339
Zimbabwe	255	451	351	352	406	200	351	527	192	532

## ANNEX 3 – J. REPORTED MALARIA DEATHS, 2010–2019

WHO region Country/area	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>AMERICAS</b>										
Argentina <sup>1</sup>	0	0	0	0	0	0	0	0	0	0
Belize	0	0	0	0	0	0	0	0	0	0
Bolivia (Plurinational State of)	0	0	0	0	1	0	0	0	0	0
Brazil	76	70	60	40	36	35	35	34	56	36
Colombia	42	23	24	10	17	18	36	19	9	3
Costa Rica	0	0	0	0	0	0	0	0	0	0
Dominican Republic	15	10	8	5	4	3	2	1	1	4
Ecuador	0	0	0	0	0	0	0	1	0	0
El Salvador <sup>3</sup>	0	0	0	0	0	0	0	0	0	0
French Guiana	1	2	2	3	0	0	0	0	0	0
Guatemala	0	0	0	1	1	1	0	0	0	0
Guyana	24	36	35	14	11	12	13	11		15
Haiti	8	5	6	10	9	15	13	24	26	7
Honduras	3	2	1	1	2	0	0	1	1	0
Mexico	0	0	0	0	0	0	1	0	1	0
Nicaragua	1	1	2	0	0	1	2		3	1
Panama	1	0	1	0	0	0	0	0	0	0
Paraguay <sup>1</sup>	0	0	0	0	0	0	0	0	0	0
Peru	0	1	7	4	4	5	7	10	4	5
Suriname	1	1	0	1	1	0	0	1	0	0
Venezuela (Bolivarian Republic of)	18	16	10	6	5	8	105	333	257	126
<b>EASTERN MEDITERRANEAN</b>										
Afghanistan	22	40	36	24	32	49	47	10	1	0
Djibouti	0	0	0	17	28	23	5			
Iran (Islamic Republic of) <sup>3</sup>	0	0	0	0	0	1	0	1	0	0
Pakistan		4	260	244	56	34	33	113	102	0
Saudi Arabia	0	0	0	0	0	0	0	0	0	0
Somalia	6	5	10	23	14	27	13	20	31	20
Sudan	1 023	612	618	685	823	868	698	1 534	3 129	1 663
Yemen	92	75	72	55	23	14	65	37	57	5
<b>EUROPEAN</b>										
Armenia <sup>1</sup>	0	0	0	0	0	0	0	0	0	0
Azerbaijan <sup>3</sup>	0	0	0	0	0	0	0	0	0	0
Georgia <sup>3</sup>	0	0	0	0	0	0	0	0	0	0
Kyrgyzstan <sup>1</sup>									0	0
Tajikistan <sup>3</sup>	0	0	0	0	0	0	0	0	0	0
Turkey <sup>3</sup>	0	0	0	0	0	0	0	0	0	0
Turkmenistan <sup>1</sup>	0	0	0	0	0	0	0	0	0	0
Uzbekistan <sup>1</sup>	0	0	0	0	0	0	0	0	0	0

## ANNEX 3 – J. REPORTED MALARIA DEATHS, 2010–2019

WHO region Country/area	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<b>SOUTH-EAST ASIA</b>										
Bangladesh	37	36	11	15	45	9	17	13	7	9
Bhutan	2	1	1	0	0		0	0	0	0
Democratic People's Republic of Korea	0	0	0	0	0	0	0	0	0	0
India	1 018	754	519	440	562	384	331	194	96	77
Indonesia	432	388	252	385	217	157	161	47	34	49
Myanmar	788	581	403	236	92	37	21	30	19	14
Nepal	6	2	0	0	0	0	3	4	0	0
Sri Lanka <sup>1</sup>	0	0	0	0	0	0	0	0	0	
Thailand	80	43	37	47	38	33	27	15	15	13
Timor-Leste <sup>3</sup>	58	16	6	3	1	0	0	0	0	0
<b>WESTERN PACIFIC</b>										
Cambodia	151	94	45	12	18	10	3	1	0	0
China <sup>3</sup>	19	33	0	0	0	0	0	0	0	0
Lao People's Democratic Republic	24	17	44	28	4	2	1	2	6	0
Malaysia <sup>4</sup>	13	12	12	10	4	4	2	0	0	0
Papua New Guinea	616	523	381	307	203	163	306	273	216	180
Philippines	30	12	16	12	10	20	7	4	2	9
Republic of Korea	1	2	0	0	0	0	0	0	0	1
Solomon Islands	34	19	18	18	23	13	20	27	7	14
Vanuatu	1	1	0	0	0	0	0	0	0	0
Viet Nam	21	14	8	6	6	3	3	6	1	0
<b>REGIONAL SUMMARY</b>										
African	150 383	104 057	104 113	116 354	99 376	118 286	103 748	94 213	73 276	82 189
Americas	190	167	156	95	91	98	214	435	358	197
Eastern Mediterranean	1 143	736	996	1 048	976	1 016	861	1 715	3 320	1 688
European	0	0	0	0	0	0	0	0	0	0
South-East Asia	2 421	1 821	1 229	1 126	955	620	560	302	164	162
Western Pacific	910	727	524	393	268	215	342	325	232	204
<b>Total</b>	<b>155 047</b>	<b>107 508</b>	<b>107 018</b>	<b>119 016</b>	<b>101 666</b>	<b>120 235</b>	<b>105 725</b>	<b>96 990</b>	<b>77 350</b>	<b>84 440</b>

Data as of 17 November 2020

<sup>1</sup> Certified malaria free countries are included in this listing for historical purposes.

<sup>2</sup> In May 2013, South Sudan was reassigned to the WHO African Region (WHA resolution 66.21, [https://apps.who.int/gb/ebwha/pdf\\_files/WHA66/A66\\_R21-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R21-en.pdf)).

<sup>3</sup> There are no indigenous malaria deaths.

<sup>4</sup> In Malaysia, there was no local transmission of human malaria in 2018. Malaria deaths were imported non-human malaria.

Note: Deaths can be probable and confirmed or only confirmed deaths depending on the country. Malaria deaths presented in this annex are considered to be due to local transmission.

# Notes









**Imperial College  
London**

For further information please contact:

**Global Malaria Programme**

**World Health Organization**


20, avenue Appia

CH-1211 Geneva 27

Web: [www.who.int/teams/global-malaria-programme](http://www.who.int/teams/global-malaria-programme)

Email: [infogmp@who.int](mailto:infogmp@who.int)

ISBN 978 92 4 001579 1



9 789240 015791